

# Comparative *In Vitro* Antioxidative Potential of *Withania Somnifera(L)* Root Extract

<sup>1</sup>Masroof Ahmad Malik

Department of Biochemistry, Chaudhary Charan Singh University

Meerut-250001

## Abstract

**Objective:** To evaluate and compare *in vitro* antioxidant activity of methanol extract of the indigenous *Withania somnifera* with the imported root extract.

**Methods:** ABTS<sup>•+</sup> radical cation de-colorization assay, DPPH<sup>•</sup> radical scavenging activity, nitric oxide radical scavenging activity, Fe<sup>2+</sup> chelating activity assay, superoxide anion and hydrogen peroxide radical scavenging activity and ferrous reducing power were studied.

**Results:** The indigenous root extracts was observed comparatively better antioxidant potential than the imported root extract.

**Conclusion:** Indigenous root extract showed high antioxidant potential compared to the imported root extract.

**Keywords:** *Withania somnifera*, antioxidative, indigenous, foreign, *In vitro*, etc

## INTRODUCTION

The use of herbal medicine and herbal medicinal products both in developing and developed countries are increasing day by day due to their origin and less side effects. There are many sources of traditional medicines. These may be medicinal plants, minerals and organic and inorganic matter [1]. Traditional herbal medicines are new therapeutic candidates because of their, structural complexity, chemical diversity, lack of substantial toxic effects, and broad spectrum of antimicrobial activity [2].

Medicinal plant, *Withania somnifera* is a small, woody shrub of 60-200 cm height in the Solanaceae family. *Withania somnifera* is locally known as Ginseng and Ashwagandha. It can be found in the India, Mediterranean, and Africa. The roots of this plant are mainly used therapeutically [3, 4]. Withanolides, which are the active pharmaceutical ingredients, are isolated from the root and leaves of *Withania somnifera*. Recently, the plant was found to show antibacterial activities [5, 6 and 7]. Besides antibacterial activities it also shows immune modulatory and antitumor activity [8, 9 and 10]. Sehgal, 2012 described that root extract of *Withania somnifera* can reverse Alzheimer's disease pathology via the peripheral clearance of  $\beta$ -amyloid [11]. Kataria *et. al.* was found that the water extract from the leaves of *Withania somnifera* protect RA differentiated C<sub>6</sub> and IMR-32 cells against glutamate-induced excitotoxicity [12].

Ashwagandha was found to have anti-carcinogenic effects. Research on animal cell cultures has revealed that the herb reduces the intercellular tumor necrosis factor, decreases the levels of the nuclear factor kappa B and potentiates apoptotic signaling in cancerous cell lines [13]. Ashwagandha also has capacity to fight cancers by reducing tumor size [14, 15].

The reactive oxygen species (ROS) such as hydroxyl, hydrogen peroxide, superoxide, nitric oxide radical are continuously being produced during regular physiological processes. These reactive radicals may cause cellular injuries, damage bio-molecules such as carbohydrates, nucleic acids, proteins, poly unsaturated fatty acids and lipids, eventually resulting in cancer, cardiovascular diseases, and diabetes. Previous researches conducted on *Withania* species revealed the presence of steroidal lactones, alkaloids, flavonoids, tannin. However, to date no studies have been reported on the locally grown indigenous *Withania somnifera*. Therefore, the aim was to evaluate the antioxidant potential of the *Withania somnifera* and to compare it with the imported root extract.

## 1. MATERIAL AND METHODS

### 1.1. Collection of the Plant Samples

The experiment was carried out on two kinds of *Withania somnifera* root samples, one is *Withania Somnifera* Indigenous (WSI) and other is *Withania Somnifera* Foreign (WSF). WSI was collected from botanical garden, Deptt. Of Botany Chaudhary Charan Singh university, Meerut during winter season. The WSF was imported from Bangladesh National Herbarium, Sector-2, Mirpur, Dhaka-1216, Bangladesh.

### 1.2. Chemicals

All the chemicals used as methanol, petroleum ether, sodium carbonate, aluminium chloride, ferrous chloride, ferric chloride, disodium hydrogen phosphate dodeca hydrate, potassium persulfate, sodium dihydrogen phosphate dehydrate, Ethylenediaminetetraacetic acid (EDTA), potassium ferricyanide, gallic acid, Folin's-Ciocalteu reagent, quercetin, L-ascorbic acid, 2, 2-Diphenyl-1-Picryl Hydrazyl (DPPH), trichloroacetic acid (TCA), ferrozine, sodium nitroprusside, citric acid, sulfanilamide, o- phosphoric acid, naphthyl ethylenediamine dihydrochloride, 2, 2'-Azinobis-3-ethylbenzothiozoline-6- sulfonic acid (ABTS), phenazine methosulphate, ferrous chloride tetrahydrate, nitro blue tetrazolium (NBT) and hydrogen peroxide were purchased from Merck (Germany).

### 1.3. Extract Preparation

The collected roots of *Withania Somnifera* were washed in water, dried in open air for 2-3 weeks at 35-40 °C temperature. Those dried materials were pulverized in an electric grinder. 100 g dry powders were defatted with petroleum ether and successively extracted with methanol (64-66 °C). Finally, 12.0 g defatted dry extract was obtained.

### 1.4. Phytochemical Screening

Phytochemical screening of defatted methanolic extract of *Withania somnifera* was carried out according to the method used by Mbaebie *et al.*, 2012 [16].

#### 1.4.1. Estimation of Total Phenolic Content

Total phenolic content of the methanolic extracts of the root of *Withania somnifera* was determined using gallic acid equivalence (GAE) [17]. The dry extracts were diluted with methanol to obtain a concentration of 1 mg/ml. 0.5 ml of each sample was transferred to a 10 ml volumetric flask, to which 5 ml undiluted Folin's-Ciocalteu reagent was added. After one minute, 4 ml of 7.5 % (w/v) Na<sub>2</sub>CO<sub>3</sub> was added, and the volume was made up to 10 ml with distilled water. The reaction mixture was incubated at 40 °C for half an hour; the absorbance of solution was measured at 765 nm by UV- Visible spectrophotometer (Model: UV-1800, Shimadzu Corporation, Japan) and compared with a pre-prepared gallic acid calibration curve. The end point of reaction mixture was indicated by the formation of blue color in the solution.

#### 1.4.2. Determination of Total Flavonoid Content

Total flavonoid content of the methanolic extracts of *Withania somnifera* root was assessed by the method followed by Velusamy and Veerabahu, 2012 [18]. 0.5 ml of extract (1:10 g/ml) was mixed with ethanolic solution of AlCl<sub>3</sub> (10 %, 0.1ml) and potassium acetate (0.1 ml of 1 M). After half hour incubation at room temperature, a yellow color indicated the presence of flavonoid. The absorbance was measured at 415 nm. The total flavonoid content was calculated as quercetin equivalent (mg QE/g).

#### 1.4.3. DPPH Radical Scavenging Activity

The ability of *Withania somnifera* extracts to scavenge DPPH radical was assessed by using the method of Shih *et al.*, 2010 with modification [19]. Briefly, the aliquots of the extract with varying concentrations (ranging from 20-100 µg/ml) were mixed with 2.0 ml DPPH (0.1mmol/L in methanol). After 30 minute incubation at 37 °C the absorbance was recorded at 517 nm with the UV-Visible spectrophotometer. The following formula was used to determine the percentage of scavenging activity,

$$\text{Percentage of inhibition (\%)} = [(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}}] \times 100 \quad (1) \text{ Where,}$$

A<sub>control</sub> – absorbance of DPPH,

A<sub>sample</sub> – absorbance reaction of mixture (DPPH with Sample)

#### 1.4.4. Ferric Reducing Power

The reducing ability of *Withania somnifera* methanolic extracts was measured according to the method of Singh *et al.*, 2009 [20]. Methanolic extracts (1.0 ml) of varying concentrations (20-100 µg/ml) were mixed with 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide (1 %). The mixture was incubated at 50 °C for 20 minutes with TCA (10 %: 2.5 ml). Then the mixture was centrifuged at 3000 rpm for 10 minutes. The supernatant (2.5 ml) was mixed with 2.5 ml of distilled water and 0.5 ml of ferric chloride (1%). The absorbance of the reaction mixture was measured at 700 nm by using the UV-Visible spectrophotometer. The higher absorbance of the reaction mixture indicated greater reducing power. The reducing power of methanolic extract of *Withania somnifera* was compared with that of standard antioxidant L-ascorbic acid (1000 µg/ml).

#### 1.4.5. Fe<sup>2+</sup> Chelating Activity Assay

The chelating activity of *Withania somnifera* root extracts was evaluated by measuring the Fe<sup>2+</sup> chelating activity according to the method of Thambiraj *et al.*, 2012 [21]. Aliquots of extracts at various concentrations (20-100 µg/ml) were added to 0.1 ml of FeCl<sub>2</sub> (2 mM) in 1.6 ml of distilled water. After 30 s, 0.2 ml of ferrozine solution (5 mM) was added to the solution. The reaction mixture was incubated for 10 minutes at 30 °C and the absorbance of the Fe<sup>2+</sup> ferrozine complex was measured at 562 nm. A lower absorbance indicates a higher chelating power. The chelating activity of the extracts on Fe<sup>2+</sup> was compared with that of EDTA (0.01 mM) and citric acid (0.025 M). The percentage of chelating activity was calculated using the following formula:

$$\text{Chelating Activity (\%)} = (A_1 - A_2) / A_1 \times 100$$

Where, A<sub>1</sub> – absorbance of the reaction mixture without extract, A<sub>2</sub> –

absorbance of the reaction mixture with extract

#### 1.4.6. Nitric Oxide Radical Scavenging Activity

Nitric oxide radical scavenging activity measurement was carried by the method of Olabinri *et al.*, 2010 [22]. Sodium nitroprusside solution was used to produce nitric oxide radicals. 1ml of 10 mM sodium nitroprusside solution was mixed with 1 ml of methanolic extract of *Withania somnifera* aliquot of the extract of different concentrations (20-100 µg/ml) in phosphate buffer (0.2 M pH 7.4). The mixture was incubated at 30 °C for 160 minute. After incubation the reaction mixture was mixed with 1.0 ml of pre-prepared Griess reagent (1% sulphanilamide, 0.1% naphthylethylenediamine dichloride, and 2% phosphoric acid). The absorbance was measured at 546 nm and the percentage of inhibition was calculated utilizing the formula no.1. A decrease in the absorbance indicates a high nitric oxide scavenging activity.

#### 1.4.7. ABTS<sup>•+</sup> Radical Cation Decolourisation Assay

The *Withania somnifera* extracts were evaluated for their ABTS<sup>•+</sup> radical capacity following a method used by Ramesh and Sivasudha, 2012 [23]. The experiments were carried out using an improved ABTS decolorization method. ABTS<sup>•+</sup> was generated by oxidation of ABTS with potassium persulfate. 2 ml of generated ABTS<sup>•+</sup> solution was mixed with 20 ml of methanol extract solution of different concentrations (20-100 µg/ml). The decreasing absorption was measured after 6 min interval at 734 nm. The inhibition of the ABTS<sup>•+</sup> radical scavenging activity was calculated using the formula no. 1.

#### 1.4.8. Superoxide Anion Scavenging Activity

Superoxide anion scavenging activity was assessed according to the method of Ilhami *et al.*, 2005 [24]. The reaction mixture consisted of various concentrations (20-100 µg/ml) of *Withania somnifera* root extracts (1 ml in methanol), 1 ml of phenazine methosulphate (PMS) solution (60 µM PMS in phosphate buffer 0.1 M) and 1 ml of nitro blue tetrazolium (NBT) solution (150 µM NBT in phosphate buffer 0.1M) was prepared to initiate the reaction. The reaction mixture was incubated at 27 °C for 10 minute, and the absorbance was measured at 560 nm. L-ascorbic acid was used as standard. Formula no. 1 was used to calculate the ability to scavenge the superoxide radicals and the percentage inhibition.

#### 1.4.9. Hydrogen Peroxide Radical Scavenging Activity

The hydrogen peroxide radical scavenging activity was assessed by utilizing the method used by Mohammad and Ali, 2010 [25]. Methanolic extract of *Withania somnifera* root aliquots of concentrations ranging 20-100 µg/ml were added to 0.6 ml hydrogen peroxide (40 mM) solution in already prepared phosphate buffer (pH 7.4). The reaction mixtures were then incubated for 15 minutes at room temperature. After incubation, the reaction mixture absorbance reading at 230 nm against phosphate buffer was taken as blank. The percentage of inhibition calculated based on the formula no. 1 as:

$$\text{Inhibition (\%)} = (A_1 - A_2) / A_1 \times 100$$

Where, A<sub>1</sub>—absorbance of the H<sub>2</sub>O<sub>2</sub>

A<sub>2</sub>—absorbance of the reaction mixture with extract

### 1.5. Statistical Analysis

All experiments were repeated at least thrice. The results were expressed with the mean ± standard deviation.

## 2. RESULTS

### 2.1. Phytochemical Screening of *Withania somnifera*

The defatted methanolic extract of *Withania somnifera* root contains tannins, terpenoids, flavonoids, alkaloids, cardio glycosides, steroids, amino acids and proteins (Table-1).

**Table1.** Phytochemical screening of the *Withania somnifera*

| Extracts        | Major secondary metabolites present |     |     |     |   |    |    |    |    |
|-----------------|-------------------------------------|-----|-----|-----|---|----|----|----|----|
|                 | AL                                  | SA  | TER | FLA | S | G  | T  | AA | P  |
| Indigenous root | ++                                  | +++ | -   | +++ | + | ++ | ++ | ++ | ++ |
| Imported root   | +                                   | +++ | -   | ++  | - | +  | +  | ++ | ++ |

Alkaloids – AL, Saponins– SA, Terpenoids– TER, Flavonoids – FLA, Steroids – S, Glycosides –G, Tannins – T, Amino acids– AA and Protein –P

**Note:** (+) Present in low concentration, (++) Present in medium concentration, (+++) Present in high concentration, (-) Not present.

### 2.2. Estimation of Total Phenolic Content

The data of total phenolic content in methanolic extract of *Withania somnifera* root was depicted in Figure 1. The Gallic acid linear curve obtained using  $Y = 6.9104 X - 0.0937$ , ( $R^2 = 0.9972$ ). Using this gallic acid linear curve, total phenolic content values of *Withania somnifera* indigenous root and imported root were found to be 0.52 mg/ml and 0.39 mg/ml respectively. The total phenolic content was high in indigenous root extract compared to the imported root extract. It shows the *Withania somnifera* indigenous root possesses high antioxidant ability.

### 2.3. Determination of Total Flavonoid Content

The total flavonoid (mg/ml) content was obtained using the regression calibration curve  $Y = 6.9104X - 0.0937$ , ( $R^2 = 0.9972$ ) (Figure 2) with quercetin equivalent. For the indigenous *Withania somnifera* root extract, the value of total flavonoid content was found to be 0.50 mg/mL, which is comparatively higher than that found for imported root with a value of 0.38 mg/mL.




Figure 1. Total phenolic content




Figure 2. Total flavonoid content

### 2.4. DPPH Radical Scavenging Activity

The photometric evaluation of the antioxidant capacity of the methanolic extracts of *Withania somnifera* indigenous (WSI) root and imported (WSF) root showed good antioxidant capacity (Figure 3). Significant decrease was observed in the DPPH radical activity due to the scavenging ability of the extracts. The  $IC_{50}$  values of the methanolic extract of *Withania somnifera* imported (WSF) root, indigenous (WSI) root and standard antioxidant (ascorbic acid) were found as 18  $\mu\text{g/mL}$ , 12  $\mu\text{g/mL}$  and 10  $\mu\text{g/mL}$  respectively. A lower  $IC_{50}$  value indicates a higher free radical scavenging activity [26]. The ability of DPPH radical scavenging is higher in indigenous (WSI) root extract compare to the imported (WSF) root extract.




Figure 3. DPPH radical scavenging activity

## 2.5. Ferric Reducing Power

The reducing ability for both of the root extracts (WSI and WSF) was found to increase with increasing concentration of the sample. The higher absorbance value indicates a high antioxidant capacity of the extracts [27]. The result showed a higher value of absorbance for WSI root extract compared to the WSF root implying that the extracts possess ferric ions ( $\text{Fe}^{3+}$ ) reducing ability (Figure 4).




Figure 4. Ferric reducing power

## 2.6. $\text{Fe}^{2+}$ Chelating Activity Assay

The  $\text{Fe}^{2+}$  chelating activity assessment revealed that the root extracts show potent chelating power. The  $\text{IC}_{50}$  values were 18 µg/ml for WSI root, 21 µg/ml for WSF root and 17 µg/ml for standard L-ascorbic acid (Figure 5). The iron generates free radicals through the Haber-Weiss and the Fenton reactions responsible for the oxidative stresses in biological systems.




Figure 5.  $\text{Fe}^{2+}$  chelating activity

## 2.7. Nitric Oxide Radical Scavenging Activity

Nitric oxide radical scavenging activity estimation of *Withania somnifera* root extracts was performed by formation of nitric oxide using sodium nitroprusside. Sodium nitroprusside acts as major source of nitric oxide radicals. The extracts scavenge the nitric oxide formed from the sodium nitroprusside by inhibiting the chromophore formation, hence absorbance decreases with increasing the concentrations of the extracts [28]. The  $\text{IC}_{50}$  values were 27 µg/ml in WSI root, 55 µg/ml in WSF root (Figure 6) revealed that, WSI is more potent NO radical scavenger compared to its counterpart WSF.

## 2.8. ABTS<sup>•+</sup> Radical Cation Decolourisation Assay

The ABTS<sup>•+</sup> radical scavenging ability was found to be higher in *Withania somnifera* indigenous (WSI) root extract compared to the *Withania somnifera* imported (WSF) root extract. The IC<sub>50</sub> values were found to be 20, 18 and 16 µg/ml in WSF root, WSI root and *L*-ascorbic acid standard respectively (Figure 7).




Figure 6. Nitric oxide radical scavenging activity




Figure 7. ABTS radical cation decolourisation assay

## 2.9. Superoxide Anion Scavenging Activity

The phenomenon of reduction of the yellow dye (NBT<sup>2+</sup>) to produce the blue formazan was utilized to determine the superoxide scavenging activity of the root. The *Withania somnifera* imported (WSF) and *Withania somnifera* indigenous (WSI) root extracts have potent antioxidants capacity. The decrease in absorbance at 560 nm indicates the high antioxidant power and the IC<sub>50</sub> values of WSF root and WSI root were found to be 30 µg/ml and 19 µg/ml respectively (Figure 8).




Figure 8. Superoxide anion scavenging activity

## 2.10. Hydrogen Peroxide Radical Scavenging Activity

Hydrogen peroxide scavenging abilities of the *Withania somnifera* imported (WSF) root and *Withania somnifera* indigenous (WSI) root extracts are shown in Figure 9. *L*-ascorbic acid was used as standard antioxidant and found the IC<sub>50</sub> value to be 36 µg/ml. The *Withania somnifera* indigenous (WSI) root possesses higher scavenging activity compared to the *Withania somnifera* imported (WSF) root and IC<sub>50</sub> values were calculated to be 64 µg/ml and 94 µg/ml respectively.




Figure 9. Hydrogen peroxide radical scavenging activity

## 3. DISCUSSION

The free radicals are very unstable chemical species containing one or more unpaired electrons that causes damage to other molecules by removing electrons from them with a specific end goal to achieve stability. Free radicals assume the double part as both pernicious and valuable species since they can be either harmful or beneficial to the living system. It is apparent from this study, that *Withania somnifera* root extract possesses effective antioxidant activity. *In vitro* antioxidant assay of the methanolic extract of *Withania somnifera* root was explored in the present study by total flavonoid content, total phenolic content, nitric oxide radical, DPPH<sup>•</sup> radical, hydrogen peroxide radical and superoxide anion scavenging activities, ABTS<sup>•+</sup> radical cation de-colourisation, Fe<sup>2+</sup> chelating activity assay and ferric reducing power. The antioxidant activities of plant extracts can be ascribed to the presence of respective phytochemicals like phenolics, flavonoids, alkaloids, saponins, steroids, glycosides, tannins, protein and amino acids in that species [29, 30]. Reactive oxygen species (ROS) readily combine and oxidize biomolecules thus making them inactive subsequently causing damage to cells, tissues and organs. Antioxidants mainly work by neutralizing the free radicals produced in the biological systems [31, 32].

Higher phenolic contents such as flavonoids, polyphenols and mono-phenols are some features of medicinal plants [33]. The flavonoid and phenolic compounds are responsible for the antioxidant ability of the medicinal plants [34] and these are abundant in some vegetables, fruits and flowers [35]. Antioxidants, derived from plant origin, especially flavonoids and polyphenols have been used to treat various disease such as aging, diabetic, cancer and prevention of cardiovascular diseases.

The DPPH<sup>•</sup>, a stable free radical, is destroyed by a free radical scavenger, as the scavenger accepts an electron or hydrogen radical to become a stable diamagnetic molecule [39]. From the result, a dose-dependent relationship on the DPPH<sup>•</sup> radical scavenging activity can be observed. It has been found that, glutathione, cysteine, tocopherols, ascorbic acid, tannins, flavonoids and aromatic amines reduce and decolorize the DPPH<sup>•</sup> by their hydrogen donating ability since they produce hydrazine by producing paired electrons [40]. Flavonoids and phenolic compounds of methanolic extracts of both



indigenous and imported *Withania somnifera* root are possibly involved in this antiradical activity .

The reducing power of the extract was explored by the transformation of  $Fe^{3+}$  to  $Fe^{2+}$  through the electron transferring ability that serves as a significant indicator of its antioxidant activity. The reductive activity of the extract and the standard increased with increasing concentration. The antioxidant activity of *Withania somnifera* root extract was slightly lower than that of the standard used in this present study. Findings from this study showed that the antioxidant activity is well advocated by the observation of the antioxidant activity of *Schotia latifolia* bark extract by Mbaebie *et al.*, 2012 [16].

Stable Iron chelation formation was utilized to evaluate the chelating ability of the extracts. The high chelating power by ferrozine reduces the free ferrous ion concentration resulting in the decrease in the Fenton reaction which is implicated in many diseases [42]. The extracts showed significant metal ion scavenging effect with an increasing trend with the increase in the concentrations of the extracts. The methanolic root extracts of *Withania somnifera* were found to possess higher metal ion scavenging activity than that of the standard. It may be due to the formation of sigma bonds with the metal and effective as secondary antioxidants because they reduce the redox potential [43]. Similar trend of metal ion scavenging activity was observed in the species *Leucas indica* var *nagalapuram lana* and *Leucas indica* var *lavandulifolia* [44].

The nitric oxide plays a pivotal role in human body, particularly in various types of inflammatory processes [40] physiological process and also has its fair share of importance as the chemical mediator in endothelial cells, macrophages, neurons [40]. However, nitric oxide in excess concentration can cause several diseases. Nitrate and peroxynitrite anions are generated when excess nitric oxide reacts with oxygen [45]. The free radical scavengers in the extract compete with oxygen leading to the decrease in the concentration of nitrite ion. Both the extracts e.g. indigenous (WSI) root and imported (WSF) root have shown good ability to scavenge nitric oxide. Vijayabaskar and Shiyamala, 2012 have reported similar tendency in nitric oxide scavenging activities by *Turbinaria ornate* extract [46].

ABTS<sup>•</sup> radical is often used for the screening of complex antioxidant mixtures such as beverages, biological fluids and plant extracts for their antioxidant activities because of its ability in both the organic and aqueous media and the stability in a wide pH range [47]. The extract showed potent antioxidant activity in ABTS<sup>•</sup> method which is consistent with that reported by, Srikanth *et al.*, 2012 and is comparable with the standard used [37]. Here, the extract's radical scavenging activity is for the direct role of its phenolic compounds in free radical scavenging.

Superoxide is produced by the oxidative enzyme of a body from molecular oxygen through non-enzymatic reaction. The superoxide generates too much harmful oxygen species in a human body such as singlet oxygen and hydroxyl radicals and these may cause the peroxidation of lipids [48].

Final result of superoxide anions in the body may be the reaction with biological macromolecules leading to tissue damage [49]. The extracts were found to be efficient scavengers of superoxide radical. Raja *et al.*, 2012 has reported similar tendency in superoxide scavenging activities by *Phyllanthus acidus* leaf extracts [49]. These results and literature survey clearly indicate that the indigenous *Withania somnifera* root extract has a noticeably higher superoxide scavenging activity compared to that of the imported one.

The presence of  $H_2O_2$  in the cell culture may lead to DNA oxidative damage. Hence removal of hydrogen peroxide is very essential for antioxidant defense in cells. Ramalingam *et al.*, 2012 reported in a recent report that the methanolic extract of *Withania somnifera* root shows high degree of  $H_2O_2$  scavenging activity [44]. In our present study we observed that the *Withania somnifera* indigenous (WSI) root extract possesses higher scavenging activity compared to the imported one (WSF).

#### 4. CONCLUSION

Antioxidant activity was correlated with the ABTS<sup>•+</sup> radical cation de-colorization assay, DPPH<sup>•</sup> radical scavenging activity, ferrous reducing power, nitric oxide radical scavenging activity,  $Fe^{2+}$  chelating activity assay, superoxide anion and hydrogen peroxide radical scavenging activities. The data obtained from these studies were high in indigenous root compared to the imported root extracts which indicate *Withania somnifera* indigeneous root show higher antioxidant activity. So the isolation

of bioactive compounds from *Withania somnifera* will definitely serve as a good phytotherapeutic agent.

#### CONFLICT OF INTEREST STATEMENT

I declare that I have no conflict of interest.

#### ACKNOWLEDGEMENT

I gratefully acknowledge Department of Biochemistry, Chaudhary charan Singh, University-Meerut for their kind cooperation and help to record different instrumental data during this research work.

#### REFERENCES

- [1] Modak M., Dixit P., Londhe J., Ghaskadbi S., and Devasagayam T.P.A., Indian herbs and herbal drugs used for the treatment of diabetes, *Journal of Clinical Biochemistry and Nutrition*. 40(3), 163–173 (2007).
- [2] Mukhtar M., Rshad M., Ahmad M., Pomerantz R.J., Wigdahl B., and Parveen Z., Anti viral potentials of medicinal plants, *Virus Research*. 131(2), 111–120 (2008).
- [3] Girdhari L., and Rana A., *Withania somnifera* (Ashwagandha): a review, *Pharmacognosy Reviews*. (1), 129–136 (2007).
- [4] Fawzy T.K.H., *Medicinal Plants in Libya*, Arab Encyclopedia House, Hazmieh, Lebanon, 1<sup>st</sup> edition. (1983).
- [5] Arora S., Dhillon S., Rani G., and Nagpal A., The *in vitro* antibacterial/synergistic activities of *Withania somnifera* extracts, *Fitoterapia*. 75(3-4), 385–388 (2004).
- [6] Owais M., Sharad K.S., Shehbaz A., and Saleemuddin M., Antibacterial efficacy of *Withania somnifera* (Ashwagandha) an indigenous medicinal plant against experimental murine salmonellosis, *Phytomedicine*. 12(3), 229–235 (2005).
- [7] Teixeira S.T., Valadares M.C., Goncalves S.A., de Melo A., and Queiroz M.L.S., Prophylactic administration of *Withania somnifera* extract increases host resistance in *Listeria monocytogenes* infected mice, *International Immunopharmacology*. 6(10), 1535–1542 (2006).
- [8] Ziauddin M., Phansalkar N., Patki P., Diwanay S., and Patwardhan B., Studies on the immunomodulatory effects of Ashwagandha, *Journal of Ethnopharmacology*. 50(2) 69–76 (1996).
- [9] Dhuley J.N., Effect of some Indian herbs on macrophage functions in ochratoxin A treated mice, *Journal of Ethnopharmacology*. 58(1), 15–20 (1997).
- [10] Davis L. and Kuttan G., Immunomodulatory activity of *Withania somnifera*, *Journal of Ethnopharmacology*. 71(1-2), 193–200 (2000).
- [11] Sehgal, N. *Withania somnifera* reverses Alzheimer's disease pathology by enhancing low-density lipoprotein receptor-related protein in liver, *Proc Natl Acad Sci of USA*. 109, 3510–3515 (2012).
- [12] Kataria H., Wadhwa Renu., Sunil C. and Kaul G.K., Water extract from the leaves of *Withania somnifera* protect RA differentiated C<sub>6</sub> and IMR-32 cells against glutamate-induced excitotoxicity, *PLoS ONE*. 7(5), 7080(2012).
- [13] Ichikawa H., Takada Y., Shishodia S., Jayaprakasam B., Nair M.G., and Aggarwal B.B. "Withanolides potentiate apoptosis, inhibit invasion, and abolish osteoclastogenesis through suppression of nuclear factor-kappa B (NF-Kappa B) activation and NF-Kappa B-regulated gene expression", *Molecular Cancer Therapeutics*. 5(6), 1434-45 (2006).
- [14] Prakash J., Gupta S.K., and Dinda A.K., *Withania somnifera* root extract prevents DMBA-induced squamous cell carcinoma of skin in Swiss albino mice, *Nutr Cancer*. 42, 91-97 (2002).
- [15] Jayaprakasam B., Zhang Y., Seeram N., and Nair M., Growth inhibition of human tumor cell lines by withanolides from *Withania somnifera* leaves, *Life Sci*. 74, 125-132 (2003).
- [16] Mbaebie B.O., Edeoga H.O., and Afolayan A.J., Phytochemical analysis and antioxidants activities of aqueous stem bark extract of *Schotia latifolia* Jacq, *Asian Pac J Trop Biomed*. 2(2), 118-124 (2012).

- [17] Končić M.Z., Kremer D., Gruz J., Strnad M., Bisevac G., and Kosalec I., Antioxidant and antimicrobial properties of *Moltkia petraea* (tratt.) Griseb. flower, leaf and stem infusions, *Food Chem Toxicol.* 48(6), 1537-1542 (2010).
- [18] Velusamy K., and Veerabahu R.M., *In vitro* antioxidant studies of *Begonia malabarica* Lam. and *Begonia floccifera* Bedd, *Asian Pac J Trop Biomed.* 2, 1572-1577 (2012).
- [19] Shih C.L., Jau T.L., Chin K.W., Hsin Y.C., and Deng J.Y., Antioxidant properties of various solvent extracts from lychee (*Litchi chinensis* Sonn.) flowers, *Food Chemistry*, 114, 577-581 (2010).
- [20] Singh *et al.*, 2009 Singh R., Singh B., Singh S., Kumar N., Kumar S. and Arora S., Investigation of ethyl acetate extract/fractions of *Acacia nilotica* wild. Ex. Del as potent antioxidant, *Rec Nat Prod.* 3, 131-138 (2009).
- [21] Thambiraj J., and Paulsamy S., *In vitro* antioxidant potential of methanol extract of the medicinal plant, *Acacia caesia* (L.) Willd", *Asian Pac J Trop Biomed.* 2, 732-736 (2012).
- [22] Olabinri B.M., Odedire O.O., Olaleye M.T., Adekunle A.S., Ehigie L.O., and Olabinri P.F., *In vitro* evaluation of hydroxyl and nitric oxide radical scavenging activities of artemether, *Research Journal of Biological Science.* 5(1), 102-105 (2010).
- [23] Ramesh Kumar A., and Sivasudha T., *In vitro* antioxidant and antibacterial activity of aqueous and methanolic extract of *Mollugo nudicaulis* Lam Leaves, *Asian Pac J Trop Biomed.* 2(2), 895-900 (2012).
- [24] İlhami G.I., Hacı A.A., and Mehmet C., Determination of *in vitro* antioxidant and radical scavenging activities of propofol, *Chem Pharm Bull.* 53(3), 281-285 (2005).
- [25] Mohammad A.E., and Ali S., Antioxidant, free radical scavenging activities of *Salvia brachyantha* and its protective effect against cardiac cell injury, *Food and Chemical Toxicology.* 48, 846-853 (2010).
- [26] Ozsoy *et al.*, 2008 Ozsoy N., Can A., Yanardag R., and Akev N., Antioxidant activity of *Smilax excelsa* L. leaf extracts, *Food Chem.* 110, 571-583 (2008).
- [27] Francis M.A., Peter N.U., Julius O.O., John R., Patrick I., and Xiao-Jian Y., Free radical scavenging activity and immunomodulatory effect of *Stachytarpheta angustifolia* leaf extract, *Food Chem.* 119(4), 1409-1416 (2010).
- [28] Halliwell B., and Gutteridge J.M., Role of free radicals and catalytic metal ions in human disease: an overview, *Methods Enzymol.* 18, 1-85 (1990).
- [29] Akter R., Raquibul Hasan S.M., Mokarram Hossain M., Jamila M., Sultana S., Chowdhury Mazumder M.E.H., and Rahman S., Antidiarrhoeal and antioxidant properties of *Curcuma alismatifolia* leaves, *Australian Journal of Basic and Applied Sciences*, 4(3), 450-456 (2010).
- [30] Armoskaite V., Ramanauskiene K., Maruska A., Razukas A., Dagilyte A., Baranauskas A., and Briedis V., The analysis of quality and antioxidant activity of green tea extracts, *Journal of Medicinal Plants Research.* 5(5), 811-816 (2011).
- [31] Siddique A. N., Mujeeb M., Kalam Najmi A., and Akram M., Evaluation of antioxidant activity, quantitative estimation of phenols and flavonoids in different parts of *Aeglemarmelos*, *African Journal of Plant Science.* 4(1), 1-5 (2010).
- [32] Koksai E., Bursal E., Dikici E., Tozoglu F., and Gulcin I., Antioxidant activity of *Melissa officinalis* leaves, *Journal of Medicinal Plants Research.* 5(2), 217-222 (2011).
- [33] Zhao-Jian G., Jian-Bing L., and Xing-Guo X., Purification and characterization of polyphenol oxidase from leaves of *Cleome gynandra* L., *Food Chem.* 129(3), 1012-1018 (2011).
- [34] Sangilimuthu A., Lukmanul H.F., and Sathish K.R., Antioxidant activity of *Withania somnifera* (L.) Dunal by different solvent extraction methods, *J Pharm Res.* 4(5), 1428-1430 (2011).
- [35] Onanong K., Sirithon S., Natthida W., and Naret M., Phenolic compounds and antioxidant activities of edible flowers from Thailand, *J Funct Food.* 3, 88-99 (2011).
- [36] Periyar R.R.S., Toda S., and Kandasamy M., Phytochemical screening and *in-vitro* anti-oxidant activity of methanolic root extract of *Erythrina indica*, *Asian Pac J Trop Biomed.* 2, 1696-1700 (2012).

- Srikanth G., Manohar B.S., Kavitha C.H.N., Bhanoji Rao M.E., Vijaykumar N., and Pradeep C.H. (2010) Senthil R.K., Raj Kapoor B. and Perumal P., Antioxidant activities of *Indigofera cassioides* Rottl. Ex. DC. Using various *in vitro* assay models, *Asian Pac J Trop Biomed.* 2(4), 256-261(2012).
- [37] Sridevi M., Senthil S. and Pugalendi K.V. "In vitro antioxidant activities of *Solanum surattense* leaf extract", *Asian Pac J Trop Biomed.* 3(1), 28-34 (2013).
- [38] Kalaivani T., and Mathew L., Free radical scavenging activity from leaves of *Acacia nilotica* (L.) Wild ex Delile, an Indian medicinal tree, *Food Chem Toxicol.* 48, 298-305 (2010).
- [39] Filomena, C., Mariangela, M., Colica, C., Federica, M., Perri, V., and Dimitar, U. "Bioactive phytonutrients (omega fatty acids, tocopherols, polyphenols), *in vitro* inhibition of nitric oxide production and free radical scavenging activity of non-cultivated Mediterranean vegetables, *Food Chem.* 129(4), 1413-1419 (2011).
- [40] Halliwell B., and Gutteridge J.M.C., *Free radicals in biology and medicine*, 4<sup>th</sup> ed. Oxford: Oxford University Press (2007).
- [41] Ilhami G., Antioxidant activity of *L*-adrenaline: A structure-activity insight, *Chem Biol Interact*, 179, 71-80 (2009).
- [42] Gulcin I.S., Beydemir H., Alici A., Elmasta M. and Buyukokuroglu M.E., *In vitro* antioxidant properties of morphine, *Pharmacol Res.* 49, 59-66 (2004).
- [43] Ramalingam R., Sushruthi S., Bindu M.B., and Ravinder N.A., Antioxidant, free radical scavenging and *in vitro* cytotoxic studies of ethanolic extract of *Leucas indica* var *lavandulifolia* and *Leucas indica* var *nagalapuramlana*, *Asian Pac J Trop Biomed.* 2, 1637-1642 (2012).
- [44] Manmohan *et al.*, 2011 Manmohan S., Arindam P. and Hemendra P.S., Nitric oxide scavenging activity of synthesized methyl semicarbazone derivatives, *Der Pharm Chem.* 3(2), 460-467(2011).
- [45] Vijayabaskar P., and Shiyamala V., Antioxidant properties of seaweed polyphenol from *Turbinaria ornate* (Turner), *Asian Pac J Trop Biomed.* 2(1), 90-98 (2012).
- [46] Huang M.H., Huang S.S., Wang B.S., Wu C.H., Sheu M.J., and Hou W.C., Antioxidant and anti-inflammatory properties of *Cardiospermum halicacabum* and its reference compound *ex vivo* and *in vivo*, *J Ethnopharmacol.* 133, 743-750 (2011).
- [47] Srikanth G., Manohar B.S., Kavitha C.H.N., Bhanoji Rao M.E., Vijaykumar N., and Pradeep C.H., Studies on *in-vitro* antioxidant activities of *Carica papaya* aqueous leaf extract. *RJPBCS.* 1(2), 59 (2010).
- [48] Raja, C., Biplab, D., Nayakanti, D., and Saikat, S. Antiinflammatory, antinociceptive and antioxidant activities of *Phyllanthus acidus* L. extracts, *Asian Pac J Trop Biomed.* 2, 953- 961(2012).