

Allelopathic Influence of *Celosia argentea* L. on activity of α -Amylase during Seed Germination of *Cyamopsis tetragonoloba* (L.) Taub

Dilipkumar T Patil¹ and Shankar K. Khade²

¹Department of Botany, Smt. K.R.P. Kanya Mahavidyalaya, Urun-Islampur, Dist. Sangli, Maharashtra, India, 415409.

²Department of Botany, Smt. Meenalben Mehta College (Arts, Commerce and Science), Panchagani, Tal. Mahabaleshwara, Dist. Satara, Maharashtra, India.

ABSTRACT

Celosia argentea L. is dominant weed reported in crop fields of Islampur of Sangli district of Maharashtra, India. It has been scrutinized for its allelopathic potentiality against guar (*Cyamopsis tetragonoloba* L. Taub.). The laboratory experiments were conducted to assess activity of α -amylase during seed germination of guar after treating different concentrations (5, 20, 40, 60 and 80%) of stem, leaves and inflorescence (flower) aqueous extracts separately. There is positively correlation between the aqueous leachate and activity of α -amylase. The activity increased in germinating seeds of guar after treatment of aqueous leachates of all plant parts. The treatment of 40 to 80% inflorescence, leaf and root aqueous extract recorded doubled α -amylase activity as compared to control in germinating seeds. The activity of α -amylase was more pronounced in leaf leachate treated seedlings as compared to other leachate treatments. It indicated that allelochemicals are more in leaf than other plant parts. This study indicates that some allelochemicals are present in aqueous extract of *C. argentea* and regulated the activity of enzyme α -amylase.

KEY WORDS: Allelochemicals, α -amylase, *Cyamopsis tetragonoloba* (L.) Taub.

Celosia argentea L. etc

INTRODUCTION:

Weeds are a unwanted unplanted redundant plant that affect the growth of main crop in field through releasing chemicals called as allelochemicals (Batish *et al.*, 2007). They often affect growth dynamics crop (Kadiolgue *et al.*, 2005) and metabolic functions including photosynthesis, respiration, mineral nutrition and such others (Saxena *et al.*, 2004) through allelopathic mechanism (Benyas *et al.*, 2010). Allelopathy functions either negatively or positive interaction between the plants, results in to stimulatory or inhibitory actions on neighboring plants.

The weed, *Celosia argentea* L. is an exotic flowering herb belonging to Amaranthaceae predominately interfere in crop field of legumes (Inamdar and Kamble,2009). Guar (*Cyamopsis tetragonoloba* (L.) Taub., family Leguminosae, sub family Fabaceae), multipurpose crop and grown in India since ancient time for its green pods, is used as vegetable, and grains as pulse and green plants as fodder. India is first rank producer

of guar comprise 83% of world production (NRAA, 2014) but its field is affecting by weed *Celosia argentea* L. in western part of Maharashtra, India.

In this connection the attempt has made to study the influence of aqueous extracts plant parts of *Celosia argentea* L. on this activity of enzyme α -amylase, to evaluate the allelopathic effect of *Celosia argentea* L. on carbohydrate metabolism during seed germination of guar. This attempt signified for understanding weed crops interactions and open new area for further research on this background.

MATERIALS AND METHODS:

Preparation of aqueous leaf extracts

The weed, *C. argentea* was collected from guar fields of Islampur, Sangli district of Maharashtra, India [17° 15' - 18° 01' N latitude and 74° 12' - 74° 74' E longitude] and washed with tap water to remove soil particles. The plant parts such as leaves, roots and inflorescence were separated and shade dried for 10 days. Dried parts were powdered with the help of grinder and stored in polythene bag. The extract were prepared by taking 10 gm of fine powder of each part and poured in 100ml distilled water as pure extract, stock solution. From this extract, the different (5, 20, 40, 80%) concentrations were prepared for treatments while distilled water used as control (0%). The extract was filtered after 24h through a double layered muslin cloth; the filtrate was used as leachates, for further analysis.

Seed treatment with aqueous Leachates:

Healthy uniform seeds of guar variety Navbahar were selected and procured from authorized shop of Shetkari Sahakari Sangh Pvt. Ltd, Kolhapur. The seeds were surface sterilized with 1% sodium hypochloride for 10 min, then rinsed with distilled water for several times to remove excess of chemical. Then surface sterilized seeds were soaked for treatments in 20 to 80% concentrations of plant extracts for 6h. The seeds soaked in distilled water were used as a control. These treated seeds were placed in petriplate ((9.0 cm diameter) containing wet blotting paper and covered with a lid. At each concentration and incubation period, triplicate sets were arranged and placed in the laboratory under normal temperature for germination, for 72h. The analysis of carbohydrates and bioassay for enzyme amylase was carried out after 72h of germination.

Bioassay for enzyme Alpha amylase was carried out through a modified method of Katsumi and Fukuhara (1969).

Statistical analysis

The analysis was carried out in three replicates for all determinations and the mean were calculated.

RESULTS AND DISCUSSION:

Qualitative and quantitative changes were involved in several metabolic pathways during seed germination and seedling growth (Kengar *et al.*, 2014). Seed germination is linked with degradation and mobilization of food accumulated during seed maturation (Borisjuk *et al.*, 2004 & Penfield *et al.*, 2005). These carbohydrates are utilized by developing seedling for the synthesis of various metabolic products. Carbohydrate storage in the form of starch and oligosaccharides were hydrolyzed and increased the sugar levels due to metabolic changes in legume seeds during germination process (Urbano *et al.*, 2005). Pawar and Chavan (2007) reported the degradation of starch might be due to the enhanced action of α -amylase during the process of germination, which hydrolyzes the starch into simple carbohydrate. The entry of allelochemicals in plants may result in changes in growth with fluctuation in carbohydrate contents (Roushan Islam, 2016) and affect the various metabolic activities and growth components in plants (Mali and Kanade, 2004). Gulzar and Siddiqui, (2014) found that total carbohydrate content was increased in allelopathic treated plants. The result of present investigation showed that, the activity of amylase in guar after seed treatment of *Celosia argentea* L. plant parts leachates.

Activity of enzyme α -Amylase [E.C. 3.2.1.1]:-

The activity of amylase enzyme recorded after treatment of aqueous extracts of *C. argentea* L. on germinating seeds of guar in table. It was noticed the elevated activity of amylase after treatment of leachates of *C. argentea* L. The treatment of 40 to 80% inflorescence, leaf and root aqueous leachate recorded doubled amylase activity as compared to control in germinating seeds. The 5 to 80% inflorescence aqueous leachates treatment recorded 0.89, 1.07, 1.43, 1.64 and 1.97 μg amylose hydrolysed $\text{min}^{-1}\text{g}^{-1}$ respectively, Leaf leachates showed 1.12, 1.30, 1.81, 2.10 and 2.34 μg amylose hydrolysed $\text{min}^{-1}\text{g}^{-1}$ where as root leachates treatment reported elevated values of amylase as 0.95, 0.96, 1.12, 1.32 and 1.49 μg amylose hydrolysed $\text{min}^{-1}\text{g}^{-1}$ respective to treatment. The leaf leachates treatment influences more positively on activity of amylase as compared to treatment of inflorescence and root aqueous leachates. It indicated as allelochemicals are more in leaf than other plant parts of *C. argentea* L.

Amylase is an important hydrolytic enzyme synthesized during seed germination in plants. This enzyme is abundant in the germinating seeds and catalyses a random hydrolysis of α -1, 4 glucosidic linkage in the starch component (Kengar *et al.*, 2014). Seed development is closely associated with seed metabolism and transport processes (Weber *et al.*, 1998). It is involved in the mobilization of starch reserves which are transported as sugars and utilized by the growing embryo (Ernst David Floyd, 1971). The enzymes most commonly endorsed with the initial attack on starch granules are α -amylase and β -amylase, responsible for breakdown and initiating the mobilization of starch in germinating seeds (Trethewey & Smith, 2000).

Ramakrishnan *et. al.*, (2014) reported that the leaf leachates of *Gmelina arborea* on red gram, green gram, black gram, and chickpea they studied allelochemicals inhibited the expression and activity of the enzymes required for efficient germination. Allelopathic plants water extract application at low concentration improved the performance of maize which might be attributed to the presence of various secondary metabolic (Casimiro *et.al.*, 2001). Pawar and Chavan (2007) studied the effect of leaf leachates of *Eucalyptus globulus*, *Moringa oleifera*, *Parthenium hysterophorus* and *Glycine max* on seedlings of *Sorghum bicolor*, recorded decreased activity of α -amylase and invertase in *Sorghum bicolor*. The similar results were reported by Madane and Patil (2017), they observed increased α -amylase activity in *Cajanus cajan* and *Cicer arietinum* seeds during germination after treatment of *E. odoratum* at lower concentrations.

In present investigation, there is positively correlation between the aqueous leachate concentrations of *C. argentea* L. and activity of α -amylase in guar. The activity increased in germinating seeds of guar after treatment of aqueous leachates of all plant parts however the treatment of 40 to 80% inflorescence, leaf and root aqueous extract recorded doubled amylase activity as compared to control in germinating seeds. The activity of α -amylase was more pronounced in leaf leachate treated seedlings as compared to inflorescence and root leachate treatments. This elevation in activity of α -amylase is due to allelochemicals present in plant parts (Madane and Patil, 2017), indicated that allelochemicals are more in leaf than other plant parts. This study indicates that some allelochemicals are present in aqueous extract of *C. argentea* L. and it worked as enzyme amylase regulators.

CONCLUSION:

The present study indicated that the amylase activity in guar was stimulated in the all selected concentrations of aqueous leaf leachates of *C. argentea* with in germinating seeds of guar. This increased activity of α -amylase is due to allelochemicals present in *C. argentea* (Narwal, 1994). It needs further screening of allelochemicals and their characterization for detailed study. Therefore, present investigation recommended that, some eco-friendly preventing measures should be taken to minimize the deleterious effects of *C. argentea* L. at the time of growing crops.

Table 1: Effect of aqueous extract of *C. argentea* on Carbohydrate content and activity of alpha amylase in germinating seeds of Guar

All values are mean of three determinations

Values are expressed in μg amylose hydrolysed $\text{min}^{-1}\text{g}^{-1}$ fresh weight.

	Activity of Alpha amylase in germinating seeds of Guar # (μg amylose hydrolysed $\text{min}^{-1}\text{g}^{-1}$)		
Control	0.83		
Aqueous Leachates (%) of <i>C. argentea</i> L.	Inflorescence	Leaf	Root
5%	0.89	1.12	0.95
20%	1.07	1.30	0.96
40%	1.43	1.81	1.12
60%	1.64	2.10	1.32
80%	1.97	2.34	1.49

REFERENCE:

1. Batish, D.R., Lavanya, K., Singh, H. P. and Koohli, R.K., Phenolic allelochemicals released by *Chenopodium murale* affect the growth, nodulation and macromolecule content in chickpea and pea. *Plant Growth Regulation*, 2007, 51: 119-128.
2. Benyas, E., Hassanpouraghdam, M.B., Salmasi, S.Z. and Oskoei, O.S.K., Allelopathic effects of *Xanthium strumarium* L. shoot aqueous extract on germination, seedling growth and chlorophyll content of lentil (*Lens culinaris* Medic.). *Rom. Bio-tech. Lett.*, 2010, 15, 5223-5228.
3. Borisjuk L, Rolletschek H, Radchuk R, Weschke W, Wobus U, Weber H, Seed development and differentiation: a role for metabolic regulation. *Plant Biol.*, 2004 6: 375–386.
4. Casimiro I.A, Marchant R.P, Bhalerao T, Beckman S, Dhooge R, Swarup N, Graham D, Inze G, Sandberg P.J, Casero and Bennett M.J, Auxin transport promotes Arabidopsis lateral root initiation. *Plant Cell*, 2001, 13: 843–852.

5. Ernst, Floyd D, Amylase activity in dormant and germinating seeds of *Polygonum pensylvanicum*. Retrospective Thesis and Dissertations. 1971, paper 4536. <http://lib.dr.iastate.edu/cgi/viewcontent.cgi?article=5535&context=rtd>.
6. Gulzar A and Siddiqui MB, 2014. Evaluation of allelopathic effect of *Eclipta alba* (L.) Hassk on biochemical activity of *Amaranthus spinosus* L., *Cassia tora* L. and *Cassia sophera* L. *Afr. J. Env. Sci and Tech.*, 8: 1 - 5.
7. Inamdar, Archana and A. B. Kamble, Allelopathic Effects of the plant *Celosia argentea* L. on Seed Germination and Seedling Growth of *Vigna mungo* L. *Nature Environ. and Poll. Tech.*, 2009, 8 (1), 59-62.
8. Kadioglu, L., Yamhar, Y. and Asav, U. Allelopathic effects of weed leachates against seed germination of some plants. *J. of Env. Biol.*, 2005, 26: 169 – 173.
9. Katsumi, M. and Fukuhara, M., The activity of α -amylase in the shoot and its relation to gibberellin induced elongation. *Physiol. Plantarum*, 1969, 22, 68-75.
10. Kengar Y.D, Patil B.J and Sabale A.B, Effect of hexaconazole and triazophos on carbohydrate contents in germinating seeds of Spinach and Guar Cent. *Euro. J. Exp. Bio.*, 2014, 3 (3):16- 21.
11. Madane, Atul N. and Bhimrao J. Patil, Allelopathic effect of *Eupatorium odoratum* L. on amylase activity during seed germination of *Cicer arietinum* L. and *Cajanus cajan* (L) Millsp. *Bioscience Discovery*, 2017, 8(1):82-86.
12. Mali, A.A. and Kanade, M.B., Allelopathic effect of two common weed on seed germination, root-shoot length, biomass and protein content of jowar. *Ann. Biolog. Res.*, 2004, 5(3): 89-92.
13. Narwal, S.S. (1994): *In: Allelopathy in Crop Production*. Scientific Publishers, Jodhpur, India.
14. NRAA , Potential of Rainfed Guar (Cluster beans) Cultivation, Processing and Export in India. Policy paper No.3 National Rainfed Area Authority, NASC, Complex. DPS Marg, New Delhi-110012, India: 2014, pp109.
15. Pawar K.B and Chavan P. D, Influence of leaf leachates of soybean, Moringa, Parthenium and Eucalyptus on Carbohydrate metabolism in germinating seeds of *Sorghum bicolor* (L.) Moench, *Allelopathy Journal*, 2007, 19(2): 543-548.
16. Penfield S, Graham S, Graham I, Storage reserve mobilization in germinating oilseeds: Arabidopsis as a model system. *Biochem Soc Trans*, 2005, 33: 380–38
17. Ramakrishnan M. S., Shanmugham V., Abdul R. S., and Ramasamy R., 2014. Effect of allelochemicals from leaf leachates of *Gmelina arborea* on inhibition of some essential seed germination enzymes in green gram, red gram, black gram, and chickpea, *International Scholarly Research*, 2014, ID 108682, 7 pages
18. Roushan Islam, Allelopathy: Its Role, Recent Developments And Future Prospectus, *Int. J. of Institutional Pharmacy and Life Sci.*, 2016, 6(1):1-21

19. Saxena, S., K. Sharma, S. Kumar, N.K. Sand and Rao, P.B., Interference of three weed extracts on uptake of nutrient in three different varieties of paddy through radio tracer techniques. *J. Environ. Biol.*, 2004, 25(4): 387 – 393.
20. Trethewey, R.N And Smith A.M., Starch mobilization in leaves, *Advance in photosynthesis* (Eds): R.C.Leegod, T. D. Sharkry and VonCammerer. Vol. 9 photosynthesis: Physiology and metabolism. Dordrecht, the Netherlands, Kluwer Academic publishers, 2000, pp 205-231
21. Urbano, G., Lopez-Jurado, M., Frejnagel, S., GomezVillalva, E., Porres, J. M., Frias, J., Vidal-Valverde, C. and Aranda, P. Nutritional assesment of raw and germinated pea (*Pisum Sativum* L.) protein and carbohydrate by in vitro and in vivo techniques. *Nutrition*, 2005, 21(2): 230-239.
22. Weber, H., U. Heim, S. Golombek, L. Borisjuk and U. Wobus. 1998. Assimilate uptake and the regulation of seed development. *Seed Sci. Res.*, 8: 331-345.

