



DIVERSITY OF ZOOPLANKTON IN SHANIGARAM RESERVOIR, SIDDIPET DISTRICT, TELANGANA

Dr.T.Jagadeeshwara Chari¹B.Srishylam²,Dr.A.V.Rajashekhar³

Department of Fisheries(P.G)

¹Government Degree&PG College(A)Siddipet.Telangana State

ABSTRACT:

Zooplankton is an integral component of aquatic ecosystem and comprises of microscopic animal life that passively float or swim freely. In the present study the selected shanigaram reservoir for the purpose of the study. Zooplanktons are the smallest metazoans in all water bodies, ranging in size from about 0.05 to 10mm.They provide food for many species of fish and are therefore, vital role in the food web of ponds. A total of 16 Zooplankton taxa were observed in the lake and found dissimilarities in three seasons. During the study period the total numbers of 16 species are identified from the Selected reservoir during the study period. Rotifer, cladocera, copepod and Ostracoda. Seven number of Rotifer sps, Four number of cladocera sps, three number of copepod sps and Two number of ostracoda sps in the selected stations. In Rotifers were higher in pre-monsoon season cladocera during monsoon season and copepods in post-monsoon season were dominant taxa. This was the systematic survey on the fish diversity of this lake It is proposed that the scientific information on ichthyofaunal diversity and distribution status will surely help in serving the future purposes of sustainable exploration and concurrent conservation of fish resources. This was the systematic survey on the fish diversity of the lake.

Key words: Zooplanktons, Rotifers, Cladocera, Copepods and Ostracoda.

INTRODUCTION:

Zooplanktons form a major link in the energy transfer at secondary level in aquatic food webs between autotrophies and heterotrophy (Deivanai et al., 2004).In addition they act as indicator of water of pollution (Sharma,1983).The zooplankton occupies an intermediate position in the food web in the aquatic ecosystem. of the total rotifer count worldwide (2030) only 360 species have been reported from India. The number of cladocera species reported in India is 190(Ramachandran and Kumar,2002).The global diversity of cladocera is more than 600 species. The copepods have the longest and the strongest appendages which help them to swim faster than any other zooplankton. They are very sensitive to environmental changes and thus are of considerable potential value as water quality indicating (Gannon and Stemberger, 1978).Zooplanktons provides the main food for fishes and can be used as indicators of the tropic status of a water body. Zooplankton have long been used as indicators of the eutrophication (Weber etal 2005).

MATERIALS AND METHODS:

In the present study we carried out of zooplankton and their seasonal variation during the year May 2017-June 2018.Water samples of were collected in different stations of the lake during an early hours of the day 7.00 A.M-10.00 A.M. The plankton net is made by the bolting nylon silk (NO.25 Mesh size 50 μ) is used for collection of zooplankton and which is conical shape and reducing con with the bottle t its end. Collected samples were transected to labeled samples were containing 4% formalin. Collected samples were brought by using various authenticated monographs. After an accurate identification of ach genus, the density of zooplankton was calculated as per the Lackey Drop method.

The density of zooplankton was expressed as organisms per liter using formula:

$$N=n*v/V$$

N=Total no of organisms/Liter of water filtered.

n=No of organisms counted in 1 ml of sample.

v=Volume of concentrated sample (ml).

V=Volume of total water filtered/Liter (ml).

RESULTS AND DISCUSSION:

The present study report the zooplankton diversity community of Shanigaram reservoir. In the investigation period fifteen zooplankton were identified in this species of which Six of Rotifer four species of Cladocera Three species of Copepoda and two species of Ostracoda were identified. During the present investigation class Rotifer was dominated among all the zooplankton group in all the research showed Table-1.

Table-1: Seasonal distribution of zooplankton in Shaniagaram reservoir

Group	Species	Monsoon	Post monsoon	Pre monsoon
Rotifera	<i>Brachionus falcatus</i>	220	231	269
	<i>B. angularis</i>	198	215	264
	<i>Filinia longisepta</i>	116	125	139
	<i>Lecane monostyla</i>	123	136	144
	<i>Polyarthra remata</i>	110	105	126
	<i>Hexarthra</i> sps	132	158	160
	Total		1011	1104
Cladocera	<i>Daphniosoma</i>	118	120	142
	<i>Lactona</i>	95	96	107
	<i>Cereodaphnia</i>	383	322	373
	<i>Alonella</i> sps	134	130	128
	Total	730	668	750
Copepoda	<i>Mesocyclops</i> sp	385	371	352
	<i>Diaptonus</i> sp	83	105	118
	<i>Heliodiaptomus</i>	57	67	82
	Total	525	543	552
Ostracoda	<i>Cyprinotus</i> sp	285	384	402
	<i>Stenocypris</i> sp	184	176	145
	Total	469	560	547


Rotifers are regarded as bioindicators of water quality. Rotifers play a vital role in the trophic tiers of freshwater impoundments and serve as a living capsule of nutrition (Suresh Kumar et al. 1999). In the present study, they dominated with six species as compared to other groups of zooplankton. Taxonomic dominance has been reported in several water bodies (Kudari et al., 2005). This pattern is common in lakes, ponds, reservoirs and rivers (Naves et al., 2003). Since the rotifers have short reproductive stages, they increase in abundance rapidly under favorable environmental conditions (Dhanapathi, 2000). The rotifer group population shows the highest during the pre-monsoon season, lowest during the post-monsoon season. The similar observations were given by Kumar et al (1978).

Cladocerans are popularly known as water fleas as most of them move through the water with a series of hops and jumps. Cladoceros prefer to live deep water and constitute a major item of food for fish. The cladocera group population shows highest during the pre monsoon season, lowest during the post monsoon season. Zooplanktons are highly sensitive to environmental variation. Dutta et.al.,(2013).

Free living copepods are an essential link in the food chain occupying the intermediate tropic level between bacteria and algae n hand and small and large plankton predators on the other. Through they are well known as important intermediate hosts for helminthes parasites. The copepod group population shows highest during the pre monsoon season, lowest during monsoon season. Similar observations were given by Kambale et.al.,(2005).

Ostracoda commonly known as seed shrimps are small crustaceans having the bivalve carapace enclosing the laterally compressed body. They are found in a wide variety of aquatic habitats lakes, ponds, streams, swamps and especially shallow places; here weeds or algae are abundant. Majority of them are free living and a few are commensalisms on the gills of crayfishes and in the intestine of fishes and amphibians. The ostracoda group population shows highest during the post monsoon season, lowest during the monsoon season.

Seasonal average variation of zooplankton during the year 2017-2018



CONCLUSION:

In the present study, the zooplankton population of Shanigaram reservoir was observed that, the quantity of zooplankton found more during pre monsoon season. The rotifers were dominated among the population during pre monsoon. The ostracoda were comparatively in low profile in annual cycle.

REFERENCES:

- ❖ Deivannai, K. S. Arunprasath and Rajan, M.K. and Baskaran, S. 2004. Biodiversity of phyto and zooplankton in relation to water quality parameters in a sewage polluted pond at Ellayirampennai, Virudhanagar District. in: The proceedings of National Sustainable use, Organized by the center for biodiversity and Forest Studies, Madurai Kama raj University. Madurai.
- ❖ Dhanapathi M.V.S.S.S.(2000) Taxonomic notes on the Rotifers from India of IAAB.Publication,Hyderabad FAO,2000.
- ❖ Gannon,J.E.and Stremberger,R.S..”Zooplankton especially Rotifers and crustaceans as indicators of water quality”.Trans.Am.Micros.Soc.Vol.971.pp 16-35,1978.
- ❖ Kudari, V.A., Kadadevaru, G.G. and Kanamadu, “R.D. Zooplankton composition in some ponds of Haveri district, Karnataka” .Zoo^s Print Journal, Vol. 20(12),pp2094-2099,2005.
- ❖ Kamble, B. B. and Meshram, C.B.,A Preliminary study on zooplankton diversity at Khatijpur tank, near Achalpr, District Amravathi, Maharashtra.(2005).
- ❖ Raghunathan, M. B. and Kumar, S.R. 2002. Checklist of Indian Cladocera (crustacean).zoos; Ecology. 19(3):505-520.
- ❖ Sharma, B.K. 1983.The Indian Species of the genus Brachionus (Eurotarotia; Monogonota: Brachionida).Hydrobiol.104:34-39.
- ❖ Suresh Kumar,Altaff,R.K.and Raghunthan,M.B.” New record of a Chydorid Cladoceran,pleuroxuy Aucus Jurine,from Chennai,South India,with the description of the Development stages”.J.Aqua.Bio.,Vol.14(1&2),pp7-10,1999.
- ❖ Neves I.F., Recha O., Roche K.F. and Pinto A.A., “Zooplankton community structure of two marginal lakes of the river Cuiaba(Mato Grosso, Brazil)with analysis of Rotifer and cladocera diversity”,Braz.J.Biol.,Vol.63,pp1-20(2003).
- ❖ Weber Mona, Mayers Elecia Edwards, Cnbella and D. Wtebber (2005) phytoplankton and zooplankton as indicator of water quality in Discovery Bay Jamaica. Hydrobiologia 545:177-193.