



A Study on Electrophoresis Analysis of Alkaline Phosphatase Isozymes During Different Developmental Stages of Evolved Races R_1 and R_2 of *Bombyx mori* L

¹Manjula A.C and ^{2*}Keshamma E

¹Department of Sericulture, Maharani Cluster University, Palace Road,
Bengaluru, Karnataka, India

²Department of Biochemistry, Maharani Cluster University, Palace Road,
Bengaluru, Karnataka, India

Abstract: The molecular data, in particular gel electrophoresis of enzymes and numerical methods of analysis have proven useful in many groups of insects and will see much wider use in future. Therefore, present study was designed with the main purpose to analyze the activities of alkaline phosphatase isozymes by electrophoresis method during different developmental stages of newly evolved races R_1 and R_2 of *Bombyx mori* L. Standardized disc electrophoresis method was performed. Alkaline phosphatase isozymes form distinct enzymes zones in the photographs and in the zymogram and these have been numbered in cathodal to anodal sequence. These isozyme patterns have been established after repeated runs. The total isozymes of different developmental stages of R_1 and R_2 have been grouped into different zones. The nomenclature of enzyme banding pattern has been followed. Results demonstrated that changes in APH zymograms during development stages revealed a total of 18 bands R_1 , and 16 bands in R_2 . From the present results it is clear that APH the activity in eggs was low in R_1 and R_2 . In the larval stage, APH activity was higher in R_1 , and in R_2 moderate activity was exhibited in larval as well pupal stages. In pupal stage the activity was less in male pupae of R_1 & R_2 , and it was found to be high in female pupae R_1 & R_2 .

Keywords - *Bombyx mori* L, Alkaline phosphatase, Electrophoresis, Race 1 (R_1), Race 2 (R_2).

I. INTRODUCTION

A study on the taxonomy of closely related species is important though difficult. Earlier studies on the entomological taxonomy was based on morphologically defined species, the degree of morphological difference being the essential criterion. This topological approach still dominates date to day practices of most systematics. However, this concept has failed to deal adequately with sibling or cryptic species (Davidson et al., 1967). In such cases the use of biochemical study has been considered of some use in the separation of closely related species of insects.

Reliable methods for distinguishing members of the insect complex by chromatographic studies of pteridine species have been attempted which help to identify the members of the complex (Davidson et al., 1967, Micks et al., 1966 a,b Micks 1968). Immuno diffusing techniques have been used to separate insect species in particular mosquitoes (Cupp and Ibrahim, 1973, Cupp et al., 1970). Schumann (1973) analysed through gel diffusion techniques different strains of mosquitoes of different origin and identified them.

A science concerned with establishing durable classification has itself undergone tremendous changes in the last three decades. Traditional approach still is the basis of all taxonomical studies. The molecular data, in particular gel electrophoresis of enzymes (Yashitake 1963 1968, Eguchi 1965, 1968 Kai and Nishi 1976 Takeda et al., 1992) and numerical methods of analysis have proven useful in many groups of insects and will see much wider use in future (Berlacher 1984). With this background, current study was planned with the main purpose to analyze the activities of alkaline phosphatase isozymes by electrophoresis method during different developmental stages of new breeding lines and races of *Bombyx mori* L.

II. MATERIALS AND METHODS

Silkworm Varieties and Rearing

The pure races of bivoltine Kalimpong-A (KA) spinning oval white cocoons, New Bivoltine-18 (NB₁₈) spinning dumbbell white cocoons and multivoltine Pure Mysore (PM) spinning pointed yellow cocoons of mulberry silkworm *Bombyx mori* L. were selected for the present breeding programme. These races were obtained from their respective seed areas and are reared in cytogenetics laboratory, Jnana Bharathi, Bangalore University. The disease free layings were prepared as described by Krishnaswamy, and were incubated at 25°C and relative humidity of 60-70%. On 8th day composite layings were prepared (10-20 layings were prepared 100-200 eggs were collected from each laying). The hatched worms were reared according to the method described by Krishnaswamy. [Krishnaswamy, 1978] MS variety of mulberry leaves were used in rearing. The worms were reared in mass up to III instar, after III moult 300 worms were collected in three replicates in order to evaluate the rearing performance. Standard temperature and humidity were maintained in the rearing house.

Breeding

Single and three way crosses were made by using the above said three races. The first single cross involved KA females and PM males. The second single cross involved NB₁₈ females and PM males. During the course of breeding selection was made at the egg, larva, pupa and cocoon stages to fix the desirable traits. F₅ progenies of the respective crosses were back crossed to their respective bivoltine males to improve commercial characters.

Evolutions of New Lines R₁ and R₂

Females of KA and NB₁₈ were crossed with males of PM. The composite layings of F₁ hybrid were brushed and reared under standard laboratory conditions. The selection parameters explained earlier were applied to choose the seed cocoons for the preparation of F₂ layings. The replicates showing higher pupation rate were selected for intra family selection of cocoons. Further, segregation with respect to cocoon colour and built was noticed. Only white oval in case of KA x PM and dumbbell white in case of NB₁₈ x PM qualifying the parameter of selection were chosen for breeding in subsequent generations. The females of F₅ were backcrossed to the males of KA and NB₁₈ respectively in both the lines and reared up to 11 generations. At the end of the 11th generation the lines R₁ and R₂ were extracted with higher ERR than their respective better parents, with shorter larval period and with moderate cocoon productivity character in case of R₁ and R₂.

Breeding Plans I and II

I				II							
KA	O	O	x	PM	Cto	NB18	O	O	x	PM	Cfo
	+	i-					+	+			
			F1						F1		
			F2						F2		
			F3						F3		
			F4						F4		
F5	x	KA	O	er'		F5x	NB18	Cta	+		
			F1						F1		
			F2						F2		
			F3						F3		
			F4						F4		
			F5						F5		
			F6	(R1)					F6	(R2)	

Preparation of Enzyme Extract

The different developmental stages such as 1st day, 5th day and 9th day eggs, five larval instars (I, II, III, IV, and V instars), early, middle and late stages of male and female pupae, male moths before and after copulation. Female moths before and after egg laying of evolved races Race-1 (R₁) and Race-2 (R₂) were selected.

Electrophoresis

Disc electrophoresis was performed essentially according to Davis (1964) and Ornstein (1964). A discontinuous gel system consisting of 7.5% lower gel and 3.12 5% spacer gel was used. The lower gel consisted of one part of Tris-hydrochloric acid buffer (36.g Tris+ 48.0 ml of N HCl + 0.46 ml of TEMED, diluted to 100ml. pH 8.9), two parts of cyanogum 41 (3.08 g of cyanogum in 10ml of water), two parts of Ammonium persulphate (140mg of APS in 100ml of water) and three parts of distilled water. 1.2 ml of this solution was poured into clean, dry glass tubes (7 cm x 0.7 cm dia) held vertically. The solution was carefully over layered with distilled water and allowed to photopolymerise for 15 minutes under fluorescent lamp or day light. After polymerisation, the water layer was removed from the top and the spacer gel was added. The spacer gel consisted of 1 part of Tris phosphoretic acid buffer (5.7 g tris + 25.6 ml of 1M H₃PO₄ + 0.46 ml of TEMED diluted to 100 ml with distilled water pH 6.9) 2 parts of cyanogen 41 (1.25 g cyanogum 41 in 10ml of Distilled water) 1 part of APS (70 mg in 100 ml) and four parts of water. 0.2 ml of spacer gel was poured on the top of the lower gel each tube layered with a drop of water and allowed to Photopolymerise for 15 minutes. After polymerization the water was blotted off and the tubes with spacer gel were inserted into the rubber connectors of the upper electrode vessel. The electrode chambers were filled with electrode buffer (0.3 M boric acid and sodium hydroxide buffer pH 8.65). The

sample, suitably diluted with 20% sucrose containing bromophenol blue, was carefully layered on to each gel and subjected to electrophoresis in cold (4°C) imposing a current of 2mA per tube for 2 hours.

Staining Procedure

The stain of alkaline phosphatases (APH) constituted sodium 1-naphthyl phosphate 100 mg, polyvinylpyrrolidone 500 mg. Fast blue RR salt 100 mg. Manganese chloride 60 mg. Magnesium chloride 60 mg and sodium chloride 2 gms dissolved in 100 ml of 0.05 M Tris - HCl buffer at pH 8.5. The gels were incubated in the stain for 10-20 minutes until the bands appeared. The gels were then stored in 6% acetic acid. All pertinent gels were photographed and diagrammatic representation of the gels was presented in the form of zymograms as deeply stained, moderately stained and faintly stained bands.

Phosphatase isozymes form distinct enzymes zones in the photographs and in the zymogram and these have been numbered in cathodal to anodal sequence. These isozyme patterns have been established after repeated runs. The total isozymes of different developmental stages of R₁ and R₂ have been grouped into different zones. The nomenclature of enzyme banding pattern has been followed after Ayala et al. (1972). The relative front (Rf) of the esterase and phosphatase bands of all the developmental stages with reference to known indicator dye was calculated as follows:

$$R_f = \frac{\text{Length gel before staining / Length gel after staining}}{X}$$

(Distance moved by band / Distance moved by marker dye)

III. RESULTS

The zymograms of alkaline phosphatase isozymes of the race R₁ showed 8 APH zones. APH-1 zone consists of 1 band and is faintly stained in 144h female pupae, moderately stained in I instar larvae. APH-2 zone consists of 2 bands (2 and 3). Band 2 is faintly stained in 144h male pupae, 288h female pupae. Band 3 is faintly stained in 144h male pupae, 288h female pupae. APH-3 zone consists of 3 bands (4,5 and 6). Band 4 is faintly stained in 144h female pupae. Band 5 is moderately stained in 24h eggs. Band 6 is faintly stained in 216h eggs, darkly stained in 120h eggs, II and V instar larvae. APH-4 zone consists of 3 bands (7,8 and 9). Band 7 is faintly stained in male adult before copulation and female adult after oviposition, darkly stained in 120h, 216h eggs, III and V instar larvae. Band 8 is faintly stained in male adult after copulation, 24h female pupae, darkly stained in 120h, 216h eggs, III, IV, V instar larvae. Band 9 is moderately stained in 24h female pupae, darkly stained in 216h eggs, II, III, IV and V instar larvae, 144h, 288h male pupae, 288h female pupae. APH-5 zone consists of 3 bands (10, 11 and 12). Band 10 is moderately stained in male adult before copulation, darkly stained in II, III, IV and V instar larvae, 144h, 288h male pupae, 24h, 144h female pupae. Band 11 is darkly stained in II, III, IV instar larvae, 144h male pupae, 24h, 144h female pupae, male adult before and after copulation. Band 12 is moderately stained in 288h female pupae, darkly stained in II, III, IV instar larvae, 24h, 144h female pupae and male adult before copulation. APH-6 zone consists of 3 bands (13, 14 and 15). Band 13 is absent. Band 14 is lightly stained in 288h male pupae, darkly stained in 288h female pupae. Band 15 is absent. APH-7 zone consists of 3 bands (16, 17 and 18). Band 16 is darkly stained in 288h male pupae. Band 17 is moderately stained in 288h female pupae, and female adult after oviposition, darkly stained in I instar larvae, female adult before oviposition. Band 18 is darkly stained in II instar larvae. APH-8 zone consists of 3 bands (19, 20 and 21). Band 19 is absent. Band 20 is moderately stained in female adult before oviposition. Band 21 is commonly present in all the developmental stages (Figure 1, 2, and 3).




Figure 1: Alkaline phosphatase zymograms of R_1




Figure 2: Alkaline phosphatase zymograms of R_1




Figure 3: Alkaline phosphatase zymograms of R₁

The zymograms of the alkaline phosphatase isozymes of the race R₂ are grouped into 8 APH zones. APH-1 zone consists of one band and is moderately stained in female adult before copulation. APH-2 zone consists of 2 bands (2 and 3). Band 2 is faintly stained in 24h female pupae, moderately stained in 288h male pupae, female adult before oviposition. Band 3 is darkly stained in III instar larvae, 24h female pupae. APH-3 zone consists of 3 bands (4, 5 and 6). Band 4 is moderately stained in female adult before oviposition and darkly stained in I instar larvae. Band 5 is moderately stained in 144h male pupae, female adult after oviposition darkly stained in I instar larvae, 24, 288h male pupae. Band 6 is faintly stained in 120h eggs, moderately stained in 144h male pupae, female adult after oviposition darkly stained in 288h male pupae 24h, 288h female pupae. APH-4 zone consists of 3 bands (7, 8 and 9). Band 7 is darkly stained in male pupae. Band 8 is absent. Band 9 is faintly stained in 120h eggs, moderately stained in 144h male pupae, darkly stained in V instar larvae. APH-5 zone consists of 3 bands (10, 11 and 12). Band 10 is moderately stained in III instar larvae and 144h male pupae. Band 11 is moderately stained in 288h female pupae, darkly stained in 24h female pupae. Band 12 is moderately stained in III instar larvae, darkly stained in 144h female pupae, male adult before copulation. APH-6 zone consists of 3 bands (13, 14 and 15). Band 13 is darkly stained in 144h male pupae, 288h female pupae, male adult before copulation. Band 14 is moderately stained in III instar larvae, darkly stained in IV and V instar larvae. APH-7 zone consists of 3 bands (16, 17 and 18). Band 16 and 17 are absent. Band 18 is moderately stained in 120h eggs, 288h female pupae, female adult before oviposition and darkly stained in 216h eggs. APH-8 zone consists of 3 bands (19, 20 and 21). Band 19 is darkly stained in 216h eggs. Band 20 is darkly stained in 120h, 216h eggs. Band 21 is commonly present in all the developmental stages (Figure 4, 5, and 6).




Figure 4: Alkaline phosphatase zymograms of R_2




Figure 5: Alkaline phosphatase zymograms of R_2




Figure 6: Alkaline phosphatase zymograms of R₂

IV. DISCUSSION

The occurrence of phosphatases in silkworm *Bombyx mori* were first reported by Nakamura (1940) later many workers studied these enzymes (Day 1949, Bradfield 1951, Denuce 1952). The results obtained showed increased activity from eggs to V instar larvae and it was decreased in pupae and adults. This is supported by other workers also (Drilhon and Busnel 1945, Shridhara and Bhat 1963). The high activity of phosphatase during larval stage is due to the hydrolysis, histogenesis, cell differentiation and transformation. Low activity of the phosphatase in pupae and adult is because of the process of hydrolysis from V instar larval stage to pupae. This is also studied in Lepidopterous (Seang et al., 1983). On the basis of above studies, it is suggested that during metamorphosis midgut tissues undergo gradual degradation under the presence of hydrolases like acid and alkaline phosphatases which are originated from lysosomes in degenerated cells. This is also studied in *Antheraea mylitta*, tasar silkworm by Sinha et al. (1991) where ACPH activity was maximum in 3rd instar and decreased in spinning stage. APH activity was minimum and decrease again in spinning stage.

The changes in APH zymograms during development stages revealed a total of 18 bands R₁, and 16 bands in R₂. From the present results it is clear that APH the activity in eggs was low in R₁ and R₂. In the larval stage, APH activity was higher in R₁. Whereas, in R₂ moderate activity was exhibited in larval as well pupal stages. In pupal stage the activity was less in male pupae of R₁ & R₂, and it was found to be high in female pupae R₁ & R₂. Here also sexual dimorphism has been observed. The present results also agree with the findings of Hedge and Krishnamurthy (1984) where the activity of APH was low in eggs but increased gradually towards V instar larvae and it was still decreased in later stages.

The phosphatases are mainly concerned with the digestion of the ingested food and degradation of the cellular substances (de Duve, 1963). The fact that the silkworm egg shows neither digestive activity nor degradation of cellular substances accounts for low activity of APH in the eggs. Further, the larval stage is the only feeding stage in silkworm after which feeding stops. In the pupal and moth stages there is a lot of resorption of cells and tissues. Hence high phosphatase activity is seen in larval stage which gradually declines in later stages. The genes which control this mechanism are active in the larval stages and are less active in pupal and get inhibited in adult stage.

Phosphatases also show specificity. APH-4 and APH-5 in R₁ and APH-3, APH-5, APH-6 zones in R₂ are non-specific but are stage specific. However, the specificity varies from one race to another race. Here some zones are stage specific also. This specificity/differences in the electrophoretic mobilities of both the enzymes implies divergence in the molecular properties of the protein (Takeda et al., 1992). Genetic studies made in *Drosophila* (Beckman and Johnson, show electrophoretic variations in larval APH controlled by a pair of codominant alleles. Subsequently genetic and developmental relationship between larval and pupal APH were investigated by Willis and Fox (1968). Genetic inheritance of polymorphic m-APH in *Bombyx* midgut was studied by (Takeda et al., 1992). The functions of both APH have been well reported in the silkworm (Sridhar and Bhatt, 1963).

V. CONCLUSION

In conclusion, changes in APH zymograms during development stages revealed a total of 18 bands R_1 , and 16 bands in R_2 . From the present results it is clear that APH the activity in eggs was low in R_1 and R_2 . In the larval stage, APH activity was higher in R_1 , and in R_2 moderate activity was exhibited in larval as well pupal stages. In pupal stage the activity was less in male pupae of R_1 & R_2 , and it was found to be high in female pupae R_1 & R_2 .

REFERENCES

- Davidson, G., Patterson, H.E., Coluzzi, M., Mason, G.F. and Micks, D.W. 1967. The *Anopheles gambiae* complex. In "Genetics of insect vectors of disease" (J.W. Wright and R. Pal Editors) Elsevier Publishing Co., Amsterdam- 211.
- Micks, D.W., Rehmet, A. and Jennins, J. 1966a. Biochemical differentiation of morphologically strains of *Aedes aegypti* (Diptera Entomol. Soc. Amer., 59: 239.
- Micks, D.W., Rehmet, A., Jennins, J., Mason, G. and Davidson, G. 1966b. A chromatographic study of the systematic relationship with in the *Anopheles gambiae* Ball.
- Micks, D.W. 1968. The use of chromatography in the microtaxonomy of *Anopheles gambiae* complex. Proc. Int. Congr. Trop. Med. Malariol. 8th Teheran Abstr. Rev. 1299.
- Cupp, E.W. and Ibrahim, A. 1973. Identification of members of the *Culex pipiens* complex by immuno diffusion and immunoelectrophoresis. J. Med. Entomol., 10: 277.
- Cupp, E.W., Ibrahim, A.N., Gilotra, S.K. and Sweet, B.H. 1970. Application of immuno diffusion methods for identification of mosquitoes. J. Parasitol, 54(4) Sect. 2. Part. 1:64.
- Schumann, W. 1973. Immunogenetic and electrophoretic studies with extracts of different adult *Culex pipiens* strains. J. Insect. Physiol. 19: 1387.
- Yoshitake, N. 1963. On the esterase types in the midgut of the silkworm *Bombyx mori* L. J. Sericult. Sci., Tokyo, 32: 285-291.
- Yoshitake, N. 1968. Phylogenetic aspects on the Japanese race of silkworm *Bombyx mori* L. J. Sci. Japan, 37: 83-87.
- Eguchi, M. and Sugimoto, T. 1965. Changes in esterase zymograms in the silkworm, *Bombyx mori* L. during development J. Insect. Physiol. 11: 1145-1149.
- Eguchi, M. and Yoshitake, N. 1968. Electrophoretic variation of proteinase in the digestive juice of the silkworm, *Bombyx mori* L. Nature, 214: 843-844.
- Kai, H. and Nishi, K. 1976. Diapause development of eggs in relation to esterase A activity. J. Physiol. 22: 133-142. *Bombyx* Insect.
- Takeda, S., Azuma, M. and Eguchi, M. 1992. Alkaline phosphatase isozymes of the silkworm midgut polymorphism in the enzyme activity antigen and electrophoretogram. J. Serie. Sci. 59(2): 127-134.
- Berlocher, S.H. 1984. Insect molecular systematics. Ann.Rev. Entomol. 29: 403-428.
- Krishnaswamy, S. 1978. Improved rearing, CSB Publication.
- Davis, B.J. 1964. Disc electrophoresis. II. Methods applications to human serum protein. Ann. N.Y. Sci., 121: 404-412.
- Ornstein, L. 1964. Disc Electrophoresis-I method application to human serum proteins. Ann. N.Y. Sci. 121: 321-349.
- Ayala, F.J., Powell, J.R., Tracey, M.L., Mourao, C.A. and Sales, S.P. 1972. Enzyme variability in *Drosophila willistoni* group IV. Genic variation in natural population of *Drosophila wilsoni*. Genetics. 70: 113-121.
- Nakamura, I. 1940. The phosphorus metabolism during the growth of the animal. The behaviour of various phosphatases and compounds of *Bombyx mori* L. During growth. Mitt. Med. Acad. Kyoto, 28: 387-416.
- Day, M.F. 1949. The distribution of alkaline phosphatases in insects. Anst. J. Sci. Res. 2 31-41.
- Brandfield, J.R.J. 1951. Phosphatases and nucleic acids in silk glands. Cytochemical aspects of fibrillar in silk glands. Cytochemical aspects of fibrillar protein secretin. Quart. J. Mier. Sci. 92: 87-112.
- Denuce, J.M. 1952. Recherches sur le syseme phosphatasique des glandes sericigenes chez lever a soie (*B. mori* L.) Experimentia, 8: 64-65.
- Drilhon, A. and Bunsel, R.F. 1945. Recherches sur les d' insects. Bull. Soc. Chim. Biol. 27 phosphatases 415-418.
- Sridhara, S. and Bhat, J.V. 1963. Alkaline and acid phosphatases of the silk worm *Bombyx mori* L. J. Ins. Physiol. Vol. 9, 693-701.
- Seong, S.I., Kobayashi, M. and Yoshitake, N. 1983. Activities of acid phosphatase and nucleases during metamorphosis in the midgut of the silkworm *Bombyx mori*. J. Serie. Sci. Jpn. 52: 191-197.
- Sinha, A.K., Sinha, U.S.P., Shivaraju, K. and Sinha, S.S. 1991. Phosphatase activity in tasar silkworm *Antheraea mylitta* D. during its larval and pupal development. Indian J. Serie. 30, No. 1, 91-92.
- Krishnamurthy, N.B., Ramesh, S.R. and Rajasekara Setty, M.R. 1984. Developmental profiles in the isozymes of land B. Esterases in the embryogenesis of silkworm *Bombyx mori*. Current Science, Vol. 53, 281-284.
- deDuve, C. 1963. The lysosome concept In-ciba foundation symposium on lysosomes (Ed) De Reuck, A.V.S. and M.P.I. Cameron. Little, Brown, Boston.
- Willis, B.B. and Fox, A.S. 1968. Genetic and developmental relationship between two alkaline phosphatases in *Drosophila melanogaster*. Biochem. Genet. 2: 141-158.