IJCRT.ORG

ISSN: 2320-2882



INTERNATIONAL JOURNAL OF CREATIVE **RESEARCH THOUGHTS (IJCRT)**

An International Open Access, Peer-reviewed, Refereed Journal

To Standardization Of Yoghurt Fortification By **Sapota Powder And Honey**

¹Yogesh Kumar Saini Author, ²Todkar Ashish P.

¹M.Sc. Student, Department of Food Technology, Parul Institute of Applied Sciences, Parul University, Limda, India ²HOD, Department of Food Technology, Parul Institute of Applied Sciences, Parul University, Limda, India

Abstract: Milk preservation by fermentation is an old age technique. Yoghurt fortified by sapota powder and honey is a soft, smooth, sweet-sour, whole milk product, prepared from lactic fermented curd. It is nutritionally dense fermented milk and fortified product, its nutritional and therapeutic value can further be enhanced by incorporating filled milk, fruit pulp, fruits powder or probiotics. They are highly nutritious product because of increased vitamin content as selective cultures are being used in preparation of chkka. The basic aim of the study was "To standardization of yoghurt fortification by sapota powder and honey." Yoghurt was prepared using as per the standard method using activated culture of L.bulgaricus and S.thermophilus and sapota powder 25%(T1), honey 25%(T1) and 50% (T1) chakka while Control sample T0 was prepared without addition of sapota powder and honey and stored for 30 days at 4°c. Proximate composition of raw materials, Sensory attributes, Physico-chemical constituents and Microbiological analysis was analysed from prepared product and data collected from that were tabulated. The range of acidity for fortified yoghurt incorporated with sapota fruits powder and honey was found to be 2.09 and 2.34 for treatment T0 to T1, respectively and the pH range for sapota powder & honey fortified yoghurt was 4.82 to 5.12 per cent, respectively. The protein content for sapota powder & honey fortified yoghurt was 7.16 and 8.50 per cent for treatment T0, and T1 respectively and the fat content of sapota powder & honey fortified yoghurt was 7.28 and 8.73per centfor treatment T0 toT1 respectively. The moisture content of sapota powder & honey fortified yoghurt was found to be 56 and 54 percent for treatment T0 to T1 respectively and the range of Tss was found to be 9.7 nd 16 per cent for treatment T0, and T1 respectively. The ash content of prepapred sapota powder & honey fortified yoghurt 2.38 and 2.82 per cent for treatment T0 toT1 respectively and sucrose content was 0 and 14.25 percent for treatment T0 and T1 respectively, storage study of sapota powder & honey fortified yoghurt (absence of yeast and mold) had good storage stability during 30 days of storage at refrigeration temperature (4°c). Sensory analysis showed a significant difference in different sensory attributes of T1 sample with the rest of the treatments. T(25)honey and T1(25) supplementation of sapota powder to fortified yoghurt was much preferred. Hence, yoghurt fortification by sapota powder and honey can provide therapeutic benefits to the consumer with longer shelf life.

Index Terms - Chakka, Honey, Lactic Acid Bacteria, Milk, , Sapota Powder.

I. INTRODUCTION

Yogurt and fermented milk products are among the most famous dairy ingredients eaten all over the world. The acidic pH of yoghurt ionises calcium and, as a result, facilitates intestinal calcium absorption. Various researchers have demonstrated the fortification of yoghurt with a number of calcium salts. In the processing of flavoured yogurt, herbal fruits may be used. Yogurt is a fermented milk product that has many fitness advantages and is therefore rather viewed by dairy researchers. The blended impact of lactobacillus delbreuckeii bulgaricus and streptococcus thermophilus organisms is nicely described in yoghurt by its taste and texture. Recent buyers are interested in eating meals that provide more than just nutrition; they also need to provide fitness benefits such as cholesterol reduction, high blood pressure reduction, anti-diabetic benefits, and so on. Yoghurt is a cultured dairy product produced by fermenting milk with or without added nonfat dry milk with lactobacillus bulgaricus and streptococcus thermophilus bacteria. It typically incorporates 12-14 per cent total milk solids and has smooth friable custard like consistency and a clean wonderful flavor. The word "yoghurt" is derived from the turkish phrase "jugurt" and is a regular food beverage in the balkans and middle east. However, its popularity has additionally spread in india. Sapota is native to central and south america, especially from the yucatan peninsula of mexico to costa rica. Sapota is grown on commercial grounds in india, the philippines, sri lanka, malaysia, Mexico, Venezuela, Guatemala, and different Central American countries. The ripe fruits contain seventy-two to 78% moisture content (w.b.) and the Total Soluble Solids (TSS) range from 12 to 18 °Brix.

II. MATARIALS

Freeze-dried probiotic live culture (Streptococcus thermophilus and Lactobacillus delbrueckii subsp. Bulgaricus,), Buffalo milk, sapota fruits, honey, culture were purchased from local market of vadodara, gujrat.


III. MATHODE

A. Peprepration Of Freeze-Dried Culture

To Prepare a Lactobacillus MRS broth (100ml) - take 100ml distilled water in a clean and sterilized 500ml conical flask. Weigh 6.75 gms of MRS Powder and add to the taken distilled water in a conical flask. Close the mouth of the conical flask with cotton cork and autoclave for 15 min at 121°c. Select the freeze-dried culture of Lactobacillus delbrueckii and Streptococcus thermophilus and weigh 2gm. Inoculation of culture in MRS broth - Add the weighed freeze-dried culture to the sterilized broth. Incubation at 37°C for 68 - 72 hrs. Obtained growth in broth. Centrifugation to collect the bacteria pellets at 4500rpm for 30 mins and store the pellets in Eppendorf tubes at $\leq 4^{\circ}$ C. Addition of bacteria pellets to the pasteurized amul milk.

B. Prepration Of Sapota Powder

Take fresh and clean sapota fruit, cut into thin slices, and place them in your Excalibur dehydration tray. Dehydrate the fruit slices at 125°F for 24-48 hours, or until completely dry. Dry slices are added to the grinder to make a smooth powder.



D. Sensory Analysis

The sensory assessment was conducted out on prepared products with various ratios of sapota powder and honey taste, and each was evaluated to a control product using a 9-point headonic scale. Flavor, body and texture, colour and look, and general acceptance were all assessed using a standard scoring card.

E. Physico-Chemical Analysis

pH, acidity, moisture content, total solid content, ash content, fat content, protein content. All property analyses were carried out in accordance with AOAC standards.

F. Microbial Analysis

Microbial analysis of ice cream was done with three different tests.

- 1) yeast and mould count
- 2) coliform count
- 3) viability count

IV. MATHODE

A. Sensory Evaluation

Fig 1 sensory attributes color of the T0 was 7.83 whereas for the experimental sample T1, T2 the score were 8.73, 8.06 was observed. Appearance of the T0 yoghurt was 6.86, whereas for the experimental sample T1, T2 the score ware 7.87, 7.43 was observed. Taste of the T0 yoghurt was 7.68, whereas for the experimental sample T1, T2 the score ware 7.72, 7.12 was observed. Flavor of the T0 yoghurt was 7.19, whereas for the experimental sample T1, T2 the score ware 7.34, 7.13, was observed. Consistency of T0 yoghurt was 7.22, whereas for the experimental sample T1, T2 the score ware 7.43, 7.26 was observe. Including all overall acceptability of T0 yoghurt was 7.78, whereas the experimental sample T1, T2 the score ware 7.24, 8.42 was observed.




Fig. 1. Sensory analysis

B. Physico-Chemical Analysis

Fresh buffalo milk used for in making of sapota yoghurt was analyzed for proximate composition like pH, acidity, moisture, fat, protein. And then prepared product was analyzed for physio-chemical constituents such as pH, Acidity, Moisture, Total solid, Ash, Fat, Protein, calcium.

Table 1 Proximate analysis of sapota powder and honey voghurt.

pН	5.12
Moisture (gm%)	54.18
Total solid (gm%)	16.64
Titrable acidity (%)	0.86
Ash (gm%)	0.95
Protein (gm%)	8.50
Lactose (gm%)	5.28
Fat (gm%)	0.80
Calcium (gm%)	58.28
Carbohydrate (gm%)	21.18

Table 1 shows the proximate analysis of yoghurt which contains pH of 5.12, moisture (54.18%), content, total solid (16.12gm%), titrable acidity at 0.86 %, where titratable acidity varies from 0.86 to 0.95% in fresh buffalo milk and ash is 0.95, protein is 8.50, and percentage of fat present in yoghurt was 0.80 gm %, lactose is 58.28 gm% and calcium percent was 21.18%.

C. Physico-Chemical Properties

Stop the fortification of yoghurt with sapota yoghurt powder ", for proximate analysis, sensory, physio-chemical and microbiological analysis. Where it shows the amount of acidic pH from T2 treatment (4.52), the highest acidity percentage in T2 (0.99), the increase in the percentage of high moisture in T0 (58.64), the level of total solid content has dropped dramatically. T0 (29.76). In addition, the yoghurt ash content of the sap was high with T2 treatment (0.95), high fat content was observed in T2 (1.03), high protein content in T1 (8.50), These physio-chemical parameters were observed and sapota powder, stored in a cool place (-18 °c). From the current research it can be concluded that Sapota powder offers excellent use in the diary industry to enrich dairy products. A mixture of sapota powder for T1 treatment is more acceptable than other T2 test treatments due to high overall acceptance points. It was noted that as the increasing level of sap and honey flour increased and in Weight, Acidiy, protein, ash, fat and pH decreased, total yogurt intensification enhanced by sap powder and honey.

D. Microbial Analysis and Shelf Life of Product

1) Yeast And Mould Count

The yeast and mould growth in sapota yoghurt was analyzed at the regular intervals viz., 10th, 20th and 30th days of storage. Initially there was no yeast and mould growth in sapota yoghurt thus, yeast and mould were absent in fresh yoghurt. Then on 15th day of storage in one experimental (T1 -10⁴) sample there was no colony was observed on plate and after that at 30th day (T2-5th dilution, T1-10⁴ dilution) of storage no growth were seen on plate.

2) Coliform Count

In each treatment of prepared yoghurt, the coliform count was nil which indicates that hygienic conditions adapted during manufacturing as well as the sanitation was carried out which was adequate.

3) Viability

Only a cluster of colonies was observed in experimental samples T1 (108 dilution) which was uncountable due to the enormous cluster. After 15 and 30 days of storage it remained same uncountable in T2 (109 dilution) and skeptical to identify and to count, hence it was a fallible result.

G. Cost Of Production Of Sapota Powder And Honey Yoghurt

The cost of sapota yoghurt is made considering the existing cost of the ingredients only. Cost data is presented in Table 5.

Table 2 Organoleptic evaluation of sapota yoghurt

Sample	Color	Consistency	Taste	Flavor	Appearance	Overall acceptability
T0	7.83	7.22	7.68	7.19	6.86	7.78
T1	8.73	7.43	7.72	7.34	7.87	7.24
T2	8.06	7.26	7.12	7.13	7.43	8.42

Table 3 Physiochemical properties of sapota yoghurt

Sample	pН	Acidity	Moisture	Total solid	Ash	Fat	Carbohydrate	Protein
Т0	4.78	1.02	58.64	29.76	0.74	0.40	2.80	7.16
T1	5.12	0.86	54.18	16.64	0.82	0.80	21.18	8.50
T2	5.02	0.99	54.92	15.69	0.95	1.03	22.86	8.21

Table 4 Changes in microbial activity during storage of sapota yoghurt

	0 day			15 day			30 day		
Microbial quality	C	Е		C	E		C	Е	
	-	10^{8}	10^{9}	=	10^{8}	10 ⁹	-	10^{8}	10^{9}
Viability	-	- 1		UC	-	=	ı	UG	UG
Yeast and mold count	-		1 - 1	- 4		-	1	ı	ı
		7		- /	-		1	-	-
Coliformcount	-	10^{2}	10^{3}	- 1	10^{2}	10^{3}	_	10^{2}	10^{3}
	,	ŀ	-	1	-	-		-	-

^{*}C- control, *E-experimental *UC- uncountable, '-'= Nil

Table 5 Cost of production of sapota yoghurt

Ingredients	Sapota yoghurt treatments							
	T0		T1		T2			
	Qty	Cost	Qty	Cost	Qty	Cost		
Milk	480 ml	25.68	120 ml	6.42	120 ml	6.42		
Freeze-dried probiotics	50 mg	5	50 mg	5	50 mg	5		
Sapota powder	-	-	25 gm	7.14	30 gm	8.57		
Honey		-	25 gm	15.5	20 gm	12.4		

V. CONCLUSIONS

The present study was under taken for "To satandardization of yoghurt fortification by sapota powder and honey", for proximate, sensory, physio-chemical and microbiological analysis. The study was divided into six phases commencing with collection of raw materials, sample preparation and raw material preparation. After developing the product, physiochemical analysis of product was done and then finally in the last sixth step microbiology analysis of different experimental and control treatments was done. According to the sensory attributes all different treatments were accepted. Results obtained were satisfactory and the fortified yoghur showed good quality characteristics on pH, Acidity, Moisture, Total solid, Ash, Fat, Protein, Lactose and moisture. Sapota powder can be very well utilized by drying and grinding it in ratios with buffalo milk containing 8.5% fat, a starter culture of L.helveticus and S.thermophilus strain added to yogurt and then yogurt is added in 1:2 ratio. It can be recommended as health food for patients due to its lower fat content and being a probiotic such type of yoghurt will be beneficial to health conscious people. Yoghurt fortification by sapota powder and honey can be stored for 30 days due to absence of yeast and mold was also lower.

REFERENCES

- AbdelRaheem, Sherief, Sherief AbdAllah, and Khaled Hassanein. "The effects of prebiotic, probiotic and synbiotic supplementation on intestinal microbial ecology and histomorphology of broiler chickens", International Journal for Agro Veterinary and Medical Sciences, 2012.
- Afoakwa, . "Effects of Fermentation and Extended Pod Storage on Cocoa Bean Quality", Cocoa Production and Processing Technology, 2014.
- Alcazar, A.. "Enzymatic-spectrophotometric determination of sucrose in coffee beans", Talanta, 20051015
- A. L. Senghor, A. Ortega-Beltran, J. Atehnkeng, P. Jarju, P. J. Cotty, R. Bandyopadhyay. "Aasafe SN01 is the First Biocontrol Product Approved for Aatoxin Mitigation in Two Nations, Senegal and The Gambia", Plant Disease, 2021
- A.M. Abdel-Sala. "Functional Foods: Hopefulness to Good Health", American Journal of Food Technology, 2010
- Annel M Hernández-Alcántara, Alfonso Totosaus, M. Lourdes Pérez-Chabela. "Evaluation of AgroIndustrial Co-Products as Source of Bioactive Compounds: Fiber, Antioxidants and Prebiotic", Acta Universitatis Cibiniensis. Series E: Food Technology, 2016
- Adriana E. Manzi, Leland D. Powell, Ajit Varki. "Analysis of Monosaccharides", Current Protocols in Molecular Biology, 2001
- Ayman Suliman Ma, Omer Turki Mamdoh Ers. "The Benets of Lactic Acid Bacteria in Yogurt on the Gastrointestinal Function and Health", Pakistan Journal of Nutrition, 2009
- Balasundram, N.. "Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses", Food Chemistry, 2006
- 10. Beáta Koréneková, Jana Kottferová, Marián Korének. "Observation of the effects of nitrites and nitrates on yogurt culture", Food Research International, 1997
- 11. Benoit Foligné, Catherine Daniel, Bruno Pot. "Probiotics from research to market: the possibilities, risks and challenges", Current Opinion in Microbiology, 2013
- 12. Biannka Pompeo Cardoso. "Efeito do probiótico Lactobacillus acidophilus LA5 em modelo de periodontite induzida em camundongos diabéticos", Universidade de Sao Paulo, Agencia USP de Gestao da Informação Academica (AGUIA),
- 13. Buket Askin, Deniz Öztürk, Begüm Durusoy, Yalcin Kaya. "Properties of yogurt fortied with capsaicin extracts dissolved in high oleic sunower oil", Journal of Food Processing and Preservation, 2021
- 14. Cai, Yimin, Huili Pang, Zhongfang Tan, Yanping Wang, Jianguo Zhang, Chuncheng Xu, Jinsong Yang, and Yang Cao. "Application of Lactic Acid Bacteria for Animal Production", Lactic Acid Bacteria, 2014.
- 15. Carlo Pompei, Alessandro Spagnolello. "Inuence of Tumbling and Dextrose on Furosine Content in Cooked Ham", Journal of Agricultural and Food Chemistry, 1997
- 16. C Butler, Eleri Owen-Jones, Mandy Lau, David Gillespie et al. "Probiotics to reduce antibiotic administration in care home residents aged 65 years and older: the PRINCESS RCT", Ecacy and Mechanism Evaluation, 2021
- 17. Daniel J Merenstein, Tina P Tan, Aleksey Molokin, Keisha Herbin Smith et al. " Safety of subsp. () strain BB-12supplemented yogurt in healthy adults on antibiotics: a phase I safety study ", Gut Microbes, 2015
- 18. Daniel John Vogt, Joel P. Tilley, Robert L. Edmonds. "Soil and Plant Analysis for Forest Ecosystem Characterization", Walter de Gruyter GmbH, 2015
- 19. D. S. Mikkelsen, S. J. Toth. "Tissue-Testing Kit for Field Determination of the Magnesium Status of Plants", Agronomy Journal, 1949
- 20. Douglas B DiRienzo. "Effect of probiotics on biomarkers of cardiovascular disease: implications for heart-healthy diets", Nutrition Reviews, 2014
- 21. DURMUS SERT, MUSTAFA T. YILMAZ, MUSTAFA KARAKAYA, EBRU BAYRAK. "PEKMEZ-DAIRY DESSERTS FORTIFIED WITH SPRAY-DRIED DAIRY POWDERS: EFFECTS OF THE INTERACTION BETWEEN SKIM MILK POWDER, YOGHURT POWDER AND BUTTERMILK POWDER BY A MIXTURE DESIGN APPROACH", Journal of Texture Studies, 2009
- 22. Ekam I. Akpakpan, Eugene N. Onyeike, Charity U. Ogunka-Nnoka. "Phytochemical Composition and Nutritive Value of Unripe and Ripe Dennettia tripetala (DT) Fruits", Asian Food Science Journal, 2019
- 23. Ekakitie Lisa Ilobekemen, Okpoghono Joel, Osuvwe Clement Orororo, Ekakitie Andrew. "AMELIORATIVE PROWESS OF BEE HONEY IN THE TISSUES OF RATS ADMINISTERED ALUMINIUM NITRATE", Scientic African, 2021
- 24. Elisabeth Norin. "Interactions of Bacteria with the Host Alteration of Microora-Associated Characteristics of the Host; Non-Immune Functions", Microbial Ecology in Health and Disease, 2/20/2000
- 25. E. M. Tanvir, Md. Sakib Hossen, Ummay Mahfuza Shapla, Milon Mondal et al. "Antioxidant, brine shrimp lethality and analgesic properties of propolis from Bangladesh", Journal of Food Biochemistry, 2018
- 26. Fangfang Pu, Yue Guo, Ming Li, Hong Zhu, Shijie Wang, Xi Shen, Miao He, Chengyu Huang, Fang He. "Yogurt supplemented with probiotics can protect the healthy elderly from respiratory infections: A randomized controlled openlabel trial", Clinical Interventions in Aging, 2017
- 27. Fatima Drissi, Didier Raoult, Vicky Merhej. "Metabolic role of lactobacilli in weight modication in humans and animals", Microbial Pathogenesis, 2017
- 28. F Drissi, V Merhej, E Angelakis, A El Kaoutari, F Carrière, B Henrissat, D Raoult. "Comparative genomics analysis of Lactobacillus species associated with weight gain or weight protection", Nutrition & Diabetes, 2014
- 29. Gerber, P A, Spinas, G A, Berneis, K. "Small dense low-density lipoprotein particles: priority as a treatment target in type 2 diabetes?", 'Future Medicine Ltd', 2012
- 30. George M. Carter, Aryan Esmaeili, Hardikkumar Shah, Debbie Indyk, Matthew Johnson, Michael Andreae, Henry S. Sacks. "Probiotics in Human Immunodeciency Virus Infection: A Systematic Review and Evidence Synthesis of Benets and Risks", Open Forum Infectious Diseases, 2016

- 31. Giuseppina Sefora Rutella, Davide Tagliazucchi, Lisa Solieri. "Survival and bioactivities of selected probiotic lactobacilli in yogurt fermentation and cold storage: New insights for developing a bifunctional dairy food", Food Microbiology, 2016
- 32. Haribalakrishna Balasubramanian. "Early probiotics to prevent childhood metabolic syndrome: A systematic review", World Journal of Methodology, 2015
- 33. Isabelle Sodini, John Montella, Phillip S Tong. "Physical properties of yogurt fortied with various commercial whey protein concentrates", Journal of the Science of Food and Agriculture, 2005
- 34. James Heinis. "BOOK AND INTRNET REVIEWS", Journal of Agricultural & Food Information, 01/24/2008
- 35. Jayakanthan Kabeerdoss, R Shobana Devi, R Regina Mary, D Prabhavathi et al. "Effect of yoghurt containing Bidobacterium lactis Bb12®on faecal excretion of secretory immunoglobulin A and human beta-defensin 2 in healthy adult volunteers", Nutrition Journal, 2011
- 36. Jim M. Wild, F. William Hersman, Samuel Patz, Iga Muradian et al. "Chapter 4 MRI of Pulmonary Ventilation", Springer Science and Business Media LLC, 2009
- 37. Junhua Xie, Shaoping Nie, Qiang Yu, Junyi Yin, Tao Xiong, Deming Gong, Mingyong Xie. " NCU116 Attenuates Cyclophosphamide-Induced Immunosuppression and Regulates Th17/Treg Cell Immune Responses in Mice ", Journal of Agricultural and Food Chemistry, 2016
- 38. Jun Nishihira. "The Role of Probiotics in Sports", Elsevier BV, 2019
- 39. Kingsley C., Enya B., Emmanuel O.. "Chapter 9 Reducing Urogenital Infections Including HIV in SubSaharan Africa -Can Probiotics Be a Viable Paradigm?", IntechOpen, 2011
- 40. Laura Ailioaie, Gerhard Litscher. "Probiotics, Photobiomodulation, and Disease Management: Controversies and Challenges", International Journal of Molecular Sciences, 2021
- 41. Law, Chung, Wan Daud, and Arun Mujumdar. "Emerging Drying Technologies for Agricultural Products", Introduction to Advanced Food Process Engineering, 2014.
- 42. Liran C. Shan, Aoife De Brún, Maeve Henchion, Chenguang Li, Celine Murrin, Patrick G. Wall, Frank J. Monahan. "Consumer evaluations of processed meat products reformulated to be healthier – A conjoint analysis study", Meat Science, 2017
- 43. Marie Céleste Karam, Jeremy Petit, David Zimmer, Elie Baudelaire Djantou, Joël Scher. "Effects of drying and grinding in production of fruit and vegetable powders: A review", Journal of Food Engineering, 2016
- 44. Maria L Marco, Dustin Heeney, Sylvie Binda, Christopher J Cifelli et al. "Health benets of fermented foods: microbiota and beyond", Current Opinion in Biotechnology, 2017
- 45. Masato Yoneda, Takashi Uchiyama, Shingo Kato, Hiroki Endo et al. "Plasma Pentraxin3 is a Novel Marker for Nonalcoholic Steatohepatitis (NASH)", BMC Gastroenterology, 2008
- 46. Mohammad Asghari Jafarabadi, Azadeh Dehghani, Leila Khalili, Ali Barzegar, Mahdi Mesrizad, Tohid Hassanalilou. "A Meta-analysis of Randomized Controlled Trials of the Effect of Probiotic Food or Supplement on Glycemic Response and Body Mass Index in Patients with Type 2 Diabetes, Updating the Evidence", Current Diabetes Reviews, 2021
- 47. M. Remely, I. Tesar, B. Hippe, S. Gnauer, P. Rust, A.G. Haslberger. "Gut microbiota composition correlates with changes in body fat content due to weight loss", Benecial Microbes, 2015
- 48. Muhammad Norazan, Abu Hassan. "Fatty acid of moringa oleifera oil at different sample condition", 2013
- 49. Nazni Peerkhan, Sandra Nair. "Optimization of wheat dextrin yogurt formulation using response surface methodology", Journal of Food Science and Technology, 2020
- 50. Neelam Arya, Om Prakash, S. Tripathi, Vivekanand, A.K. Pant. "Curcuma longa L: Elemental and Nutritional Proling of Fifty Accessions from Uttarakhand Region in India", Asian Journal of Chemistry, 2015
- 51. Poonam Baniwal, Bahadur Singh Hathan, "Process Parameter Optimization for the Development of Ready-to-Eat Instant Sand Pear Candy Using Response Surface Methodology", Journal of Food Processing and Preservation, 2015
- 52. Qingqing Zhang, Yucheng Wu, Xiaoqiang Fei. "Effect of probiotics on glucose metabolism in patients with type 2 diabetes mellitus: A meta-analysis of randomized controlled trials", Medicina, 2016
- 53. Ramesh C. Chandan. "Role of Milk and Dairy Foods in Nutrition and Health", Wiley, 2015
- 54. Sadeq Hasan Al-Sheraji, Amin Ismail, Mohd Yazid Manap, Shuhaimi Mustafa, Rokiah Mohd Yusof, Fouad Abdulrahman Hassan. "Hypocholesterolaemic effect of yoghurt containing Bidobacterium pseudocatenulatum G4 or Bidobacterium longum BB536", Food Chemistry, 2012