



STUDY OF SOME FREDHOLM INTEGRAL EQUATIONS ARISING IN APPLICATIONS OF SCIENCE AND ENGINEERING

P.L.Suresh

Associate Professor,
B .V.Raju College,Bhimavarm
West Godawari Dist (AP)

Abstract:

We present the numerical solution of the Fredholm Integral Equations by using the analytic method (Adomian Decomposition Method). To illustrate the exactness and efficiency of the proposed method (ADM), some numerical examples have been performed. A Fredholm integral equations is solved by ADM which gives us the approximate solution of the problem that tends to the exact solution of the problem.

Keywords:

Adomian Decomposition Method, Equations,Fredholm Integral Equations

Adomian Decomposition Method

The Adomian Decomposition method (ADM) is very powerful method which considers the approximate solution of a nonlinear equation as an infinite series which actually converges to the exact solution in this paper, ADM is proposed to solve some first order, second order and third order differential equations and integral equations. The Adomian Decomposition method (ADM) was firstly introduced by George Adomain in 1981. This method has been applied to solve differential equations and integral equations of linear and nonlinear problem in Mathematics, Physics, Biology and Chemistry up to know a large number of research paper have been published to show the feasibility of the decomposition method.

Proposed method for solving the Fredholm integral equation.

The type of integral equation in which the limits of the integration are constant, in which a and b are constant are called the Fredholm Integral equations, and is given as

$$\phi(x) = f(x) + \lambda \int_a^b K(x,t)\phi(t)dt \quad (1)$$

Where the function and the kernel are given in the advance, and λ is a parameter. In this part, the process of the Adomian decomposition method is used. The Adomian decomposition method involving the decomposing of the unknown function $\phi(x)$ of any equation into a addition of an infinite number of constituents defined by the decomposition series

$$\phi(x) = \sum_{n=0}^{\infty} \phi_n(x) \quad (1.1)$$

Or equivalently $\phi(x) = \phi_0(x) + \phi_1(x) + \phi_2(x) + \dots$

When the constituents $\phi_n(x), \geq 0$ will be resolved. The Adomian decomposition method investigate itself with discover the components $\phi_0(x), \phi_1(x), \phi_3(x), \dots$

To organize the recurrence relation, we substitute (1.1) into the Fredholm integral equation (1) to get

$$\sum_{n=0}^{\infty} \phi_n(x) = f(x) + \lambda \int_a^b K(x,t) \sum_{n=0}^{\infty} \phi_n(t) dt \quad (1.2)$$

Or equivalently

$$\phi_0(x) + \phi_1(x) + \phi_2(x) + \dots = f(x) + \int_a^b K(x,t) [\phi_0(t) + \phi_1(t) + \phi_3(t) + \dots] dt$$

The zeroth component $\phi_0(x)$ is spotted by all terms that are not comprises under the integral sign. This signifies that the components $\phi(x), \geq 0$ of the unknown function $\phi(x)$ are totally resolved by the recurrence relation

$$\phi_0(x) = f(x), \phi_{n+1}(x) = \int_a^b K(x,t) \phi_n(t) dt, n \geq 0$$

Or equivally

$$\begin{aligned} \phi_0(x) &= f(x) \\ \phi_1(x) &= \int_a^b K(x,t) \phi_0(t) dt \\ \phi_2(x) &= \int_a^b K(x,t) \phi_1(t) dt \\ \phi_3(x) &= \int_a^b K(x,t) \phi_2(t) dt \\ \phi_4(x) &= \int_a^b K(x,t) \phi_3(t) dt \end{aligned}$$

And so on the other constituents

Thus the constituents $\phi_0(x), \phi_1(x), \phi_3(x), \dots$ are resolved totally.

Thus the solution of the Fredholm integral equation (1) is easily acquired in a series form by utilize the series as assumption in (1.1).

Applications Fredholm integral equations:

Example1. Consider the linear Fredholm integral equation given as

$$\phi(x) = \sin x - \frac{x}{4} + \frac{1}{4} \int_0^{\pi/2} xt \phi(t) dt \quad (1.3)$$

$$\text{Let } \phi(x) = \sum_{n=0}^{\infty} \phi_n(x)$$

Then by applying the Adomian decomposition method, equation (1.3) becomes

$$\sum_{n=0}^{\infty} \phi_n(x) = \sin x - \frac{x}{4} + \frac{1}{4} \int_0^{\pi/2} xt \sum_{n=0}^{\infty} \phi_n(t) dt \quad (1.4)$$

To determine the components of $\phi(x)$, we use the recurrence relation

$$\phi_0(x) = \sin x - \frac{x}{4}$$

$$\phi_{n+1}(x) = \frac{1}{4} \int_0^{\pi/2} xt \sum_{n=0}^{\infty} \phi_n(t) dt$$

This implies

$$\Phi_0(x) = \sin x - \frac{x}{4} \quad (1.6)$$

$$\Phi_1(x) = \frac{1}{4} \int_0^{\pi/2} xt[\Phi_0(t)]dt$$

$$\Phi_1(x) = \frac{1}{4} \int_0^{\pi/2} xt[\sin t - t]dt$$

$$\Phi_1(x) = \frac{x}{4} \int_0^{\pi/2} [t \cos t - t^2]dt$$

$$\Phi_1(x) = \frac{x}{4} \left[1 - \frac{\pi^3}{24}\right]$$

$$\Phi_2(x) = \frac{1}{4} \int_0^{\pi/2} xt[\Phi_1(t)]dt$$

$$\Phi_2(x) = \frac{1}{4} \int_0^{\pi/2} xt \left[\frac{t}{4} \left[1 - \frac{\pi^3}{24}\right] \right] dt$$

$$\Phi_2(x) = \frac{x}{4} \cdot \frac{1}{4} \left(1 - \frac{\pi^3}{24}\right) \int_0^{\pi/2} t^2 dt$$

$$\Phi_2(x) = \frac{x}{4} \left(\frac{\pi^3}{24} - \frac{\pi^5}{576} \right)$$

And so on

Now by using equation (1.1) we obtain

$$\Phi(x) = \sin x - \frac{x}{4} + \frac{x}{4} - \frac{x\pi^3}{96} + \frac{x\pi^3}{96} - \frac{x\pi^5}{2304} + \dots$$

thus the solution will be $\Phi(x) = \sin x$

which is converges to exact solution by the method of successive approximations $\Phi(x) = \sin x$

Example2.

Consider the Fredholm Integral equation given as

$$\Phi(x) = \frac{5x}{6} + \frac{1}{2} \int_0^1 xt\Phi(t)dt \quad (1.7)$$

$$\text{Let } \Phi(x) = \sum_{n=0}^{\infty} \Phi_n(x)$$

Then by applying the Adomian decomposition method, equation (1.7) becomes

$$\sum_{n=0}^{\infty} \Phi_n(x) = \frac{5x}{6} + \frac{1}{2} \int_0^1 xt \sum_{n=0}^{\infty} \Phi_n(t)dt \quad (1.8)$$

To determine the components of $\Phi(x)$, we use the recurrence relation

$$\Phi_0(x) = \frac{5x}{6}$$

$$\Phi_{n+1}(x) = \frac{1}{2} \int_0^1 xt \sum_{n=0}^{\infty} \Phi_n(t)dt$$

This implies

$$\phi_0(x) = \frac{5x}{6} \quad (1.9)$$

$$\phi_1(x) = \frac{1}{2} \int_0^1 xt[\phi_0(t)]dt$$

$$\phi_1(x) = \frac{1}{2} \int_0^1 xt \left[\frac{5t}{6} \right] dt$$

$$\phi_1(x) = \frac{5x}{12} \int_0^1 [t^2]dt$$

$$\phi_1(x) = \frac{5x}{12} \left[\frac{1}{3} \right]$$

$$\phi_1(x) = \frac{5x}{36}$$

$$\phi_2(x) = \frac{1}{2} \int_0^1 xt[\phi_1(t)]dt$$

$$\phi_2(x) = \frac{1}{2} \int_0^1 xt \left[\frac{5t}{36} \right] dt$$

$$\phi_2(x) = \frac{5x}{72} \int_0^1 [t^2]dt$$

$$\phi_2(x) = \frac{5x}{72} \left[\frac{1}{3} \right]$$

$$\phi_2(x) = \frac{5x}{216}$$

And so on

Now by using equation (1.1) we obtain

$$\phi(x) = \frac{5x}{6} + \frac{5x}{36} + \frac{5x}{216} + \dots$$

Which converges to exact solution by the method of successive approximations $\phi(x) = x$

Example 3: Consider the Fredholm Integral equation given as

$$\phi(x) = x + 2 \int_0^1 xt\phi(t)dt \quad (2.0)$$

$$\text{Let } \phi(x) = \sum_{n=0}^{\infty} \phi_n(x)$$

Then by applying the Adomian decomposition method, equation (2.0) becomes

$$\sum_{n=0}^{\infty} \phi_n(x) = x + 2 \int_0^1 xt \sum_{n=0}^{\infty} \phi_n(t)dt \quad (2.1)$$

To determine the components of $u(x)$, we use the recurrence relation

$$\phi_0(x) = x$$

$$\phi_{n+1}(x) = 2 \int_0^1 xt \sum_{n=0}^{\infty} \phi_n(t)dt$$

This implies

$$\phi_0(x) = x \quad (2.2)$$

$$\phi_1(x) = 2 \int_0^1 xt[\phi_0(t)]dt$$

$$\phi_1(x) = 2 \int_0^1 xt[t]dt$$

$$\phi_1(x) = 2x \int_0^1 [t^2]dt$$

$$\phi_1(x) = 2x \left[\frac{1}{3} \right]$$

$$\phi_1(x) = \frac{2x}{3}$$

$$\phi_2(x) = 2 \int_0^1 xt[\phi_1(t)]dt$$

$$\phi_2(x) = 2 \int_0^1 xt \left[\frac{2t}{3} \right] dt$$

$$\phi_2(x) = \frac{4x}{3} \int_0^1 [t^2]dt$$

$$\phi_2(x) = \frac{4x}{3} \left[\frac{1}{3} \right]$$

$$\phi_2(x) = \frac{4x}{9}$$

And so on

Now by using equation (1.1) we obtain

$$\phi(x) = x + \frac{2x}{3} + \frac{4x}{9} + \dots$$

Which converges to exact solution by the method of successive approximations $\phi(x) = 3x$

Conclusion:

The goal of this paper is to use the Adomian decomposition method for solving the Fredholm integral equations. It is can be visibly seen that the decomposition method for the Fredholm integral equation is equivalent to successive approximation method. Even though, the Adomian decomposition method is a very strong and useful appliance for solving the integral equations

.Following points have been identified while solving the numerical examples:

1. Linear Fredholm integral equation can be solved by this method.
2. It is clear that using the Adomian decomposition method when there is increasing in the 'n' order then there is the decreasing in the error.

References:

- [1] S. Balaji and V. Venkataraman, Solution of system of Nonlinear equations using integrated RADM and ADM, Pure and Appl.Math. (2017),367-373.
- [2] E. Babolian, J. Biazar, solution of a system of linear Volterra equations by Adomian decomposition method.Far East J. Math. Sci. 2 935-945.
- [3] .Shakeri,D. mehdi, application of the decomposition method of adomain for solving the pantograph equation of order m.Appl.Math.(2010),453-460.
- [4] E. Babolian, A.R.Vahidi and Gh. AsdiCordshooli, solving differential equation by adomain decomposition Appl. Math.Comput. 167 (2005) 1150-1155.
- [5] E.Babolian and J. Baizar, Solving Concrete Examples by Adomian Decomposition method, Appl. Math. Comput. 135 (2003) 161-167.
- [6]] G.Adomian ,a review of the decomposition method and some resent results for non linear equations , math compute model (1992),101-127.
- [7] G. Adomian, Solving Frontier Problem of Physics: The Decomposition method, Kluwer, Dordrecht, (1994).
- [8]G. Adomian, Nonlinear Stochastic Systems Theory and Applications to Physics, Kluwer, Dordrecht, (1989).
- [9] E.Babolian, and J.Baizar, Solving the problem of biological species living together by Adomain Decomposition method , Appl. Math. Comput. 129 (2002) 339-343.
- [10] F.S.Fawziah ,P.G.Kirtiwant, p.Priyanka , the approximate solution of fredholm integral equation by decomposition method and its modification, Appl.Math.(2018),327-336.
- [11] F.AL.Saar, K.P.Ghadle, P.Pathade, The approximate solution of Fredholm integral equation by Adomian Decomposition method and its modification,Appl.Math.(2018),327-336.
- [12] Z.Avazzedeh and G.B. Loghmani, Numerical solution of Fredholm integral equation of the second kind by using the integral mean value theorem, Appl.Math.(2013),432-442.

