



INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

Propolis: Chemical Constituent, Mechanism And Application In Topical Wound Healing

Prof. Yojana A. Kunjir,¹ Nilesh Manjare,² Pavin Bade,² Kiran Andhale,²

1. Professor, Department of Pharmaceutical chemistry, MKCOPR, Pune, Maharashtra, India.

2. Final Year Student, Department of Pharmaceutical chemistry, MKCOPR, Pune, Maharashtra, India.

Abstract

Background/Aims: Impaired wound healing is considered to be one of the most serious complications associated with diabetes as it significantly increases the susceptibility of patients to infection. Propolis is a natural bee product used extensively in foods and beverages that has significant benefits to human health. In particular, propolis has antioxidant, anti-inflammatory and analgesic effects that could be useful for improving wound healing. In this study, we investigated the effects of topical application of propolis on the healing and closure of diabetic wounds in a streptozotocin (STZ)-induced type I diabetic mouse model. **Methods:** Sixty male mice were distributed equally into 3 experimental groups: group 1, non-diabetic control mice; group 2, diabetic mice; and group 3, diabetic mice treated daily with a topical application of propolis. **Results:** We found that diabetic mice exhibited delayed wound closure characterized by a significant decrease in the levels of TGF- β 1 and a prolonged elevation of the levels of inflammatory cytokines (IL-1 β , IL-6 and TNF- α) and MMP9 in wound tissues compared with control non-diabetic mice. Moreover, the wound tissues of diabetic mice showed a marked reduction in the phosphorylation of Smad2 and Smad3 as well as a marked reduction in collagen production. Interestingly, compared with untreated diabetic mice, topical application of propolis significantly enhanced the closure of diabetic wounds and decreased the level of IL-1 β , IL-6, TNF- α and MMP9 to near normal levels. Most importantly, compared with untreated diabetic mice, the treatment of diabetic mice with propolis significantly enhanced the production of collagen via the TGF- β 1/Smad2,3 signaling axis in wounded tissues. **Conclusion:** Our findings reveal the molecular mechanisms underlying the improved healing and closure of diabetic wounds following topical propolis application.

Key Words

Cytokines • Diabetes mellitus • Inflammation • Propolis • Wound healing

1. Introduction

In a broad sense, trauma refers to body injuries caused by physical, chemical, and biological factors, including knife injuries, crush injuries, and frostbite. Trauma has a high incidence in the population. People have done a lot of research on drugs for wound repair from the perspectives of anti-inflammatory, antibacterial, antioxidant, and antiseptic [1]. Many drugs for treating local trauma have been developed, but most drugs have single pharmacological activity and may have certain side effects on the body [2, 3]. Silver sulfadiazine (AgSD), a silver compound, is often used to prevent or treat wound colonization, and also certain antibiotic-resistant bacteria. In vitro studies have shown that when acute rat wounds are used as a model, the topical antibacterial agents silver sulfadiazine and mafenyl acetate can destroy fibroblasts and have toxicity. This indicates that silver sulfadiazine and mafenyl acetate can significantly hinder wound contraction in the body [4–6]. Some researchers have found that when sulfadiazine is used to treat burns in experimental mice, it will also produce greater inflammatory reactions, such as redness and swelling [7]. Vaseline was used as the control group to treat the wounds of mice. Compared with the drug treatment in the experimental group, the wounds healed more slowly [8]. In a large number of studies, propolis, as a natural product that can promote tissue healing, has various biological activities, such as anti-inflammatory, antibacterial, and antioxidant. It has obvious advantages in promoting wound repair and has achieved ideal therapeutic effects. This kind of wound healing refers to the healing process after the body is broken or damaged by external forces, skin and other tissues, including the regeneration of various tissues and the complex combination of granulation tissue hyperplasia and scar formation, showing the synergy of various repair processes.

Propolis has always been considered as a folk medicine. Its research can be traced back to ancient times, and it was discovered about 300 years ago. Propolis is usually a sticky substance, which is formed by the resin collected from trees by Italian worker bees and the secretion of their maxillary glands. Propolis is extremely complex and contains a variety of compounds, such as flavonoids, terpenes, phenolic acids, aldehydes, and ketones, as well as a variety of hydrocarbons, minerals, trace elements, vitamins, and enzymes. Twelve different flavonoids, namely, pinocembrin, acacetin, chrysin, rutin, luteolin, kaempferol, apigenin, myricetin, catechin, naringenin, galangin, and quercetin; two phenolic acids, caffeic acid and cinnamic acid [9] (Figure 1). Propolis is a kind of mixture, which contains a variety of chemical components, but its most active chemical substances can also play a role alone. Phenethyl caffeic acid extracted from bee propolis is a receptor activator that regulates oxidation state and NF- κ B ligand (RANKL)/osteoprotegerin (OPG) signal, and has potential protective effect on glucocorticoid induced osteoporosis (GIO) [10]. It can also promote collagen deposition, re-epithelialization and wound healing in mice 12 days after pressure ulcer. In addition, it also promoted the inflammatory response, oxidative stress, and NRF2 expression, and made the skin wound of pressure ulcer in mice heal [11]. In order to determine the relationship between polyphenol derivatives in propolis and their antioxidant and antibacterial activities, some researchers studied the propolis extract from Bihor County, Romania. Different ethanol concentrations were used to extract propolis. The total polyphenols measured ranged from 1.5 to 91.2 mg/g. The final results showed that 50% ethanol extract

provided rich polyphenols and ensured good antioxidant capacity [12]. In addition to caffeic acid phenethyl ester and polyphenols, there is also an important single-substance flavonoids. chrysin (CR) is a flavone, which exists in propolis and many plants. Populin was used to treat LPS-induced sepsis in rats, which could reduce the levels of oxidative stress markers and cytokines in patients with sepsis [13]. With the deepening research on propolis by scholars at home and abroad, the biological and pharmacological activities of propolis have been further revealed. It plays an important role in antioxidation, scavenging free radicals, antibacterial and anti-inflammatory, protecting the liver, improving human immunity, antitumor, oral health, regulating blood lipids, gastrointestinal diseases, anti-vascular effects, reducing blood glucose, and so on [14] (Table 1).

Studies have shown that different forms of propolis have an effect on wound healing (Table 2). According to these results, it can be seen that propolis has a significant effect on various wound treatments. On the basis of previous studies, this paper summarizes the effect and related mechanism of propolis on wound healing and the synergistic effect of propolis and other compounds, which further clarified the medicinal value of propolis and provided a more powerful basis for propolis theory.






Figure 1: Chemical structural formula of some active ingredients in propolis.

Table 1: Treatment of other diseases with propolis.

SR.NO	Propolis type	Medical applications	Mian result	References
1.	Green, red, or brown propolis	Atherosclerosis	Reduce atherosclerotic lesion area regulating inflammatory and angiogenic factors	[15]
2.	Istanbul, Turkey propolis	Diabetes	Decrease of blood glucose significant improvement in pancreas, liver and kidney tissue	[16]
3.	Brazilian extract of propolis (EEP)	Oral health	Effectively remove dental improve marginal periodontal tissue	[17]
4.	Propolis	Radiation resistance	Inducing apoptosis increase the phosphorylation of Akt/mTOR and hinder cell migration	[18]
5.	Brazilian propolis	Anti-ulcer effect	Inhibition of diclofenac induced ulcer formation antagonism to histaminergic system	[19]
6.	Trigona sp. propolis	Dental pulp disease	Inhibition of IL-6 expression in dental pulp of inflammatory rats	[20]
7.	Brown propolis from Southern Brazil	Anti-angiogenesis	Inhibition of tubular structure formation of endothelial cells on matrix gel (tubulogenesis)	[21]
8.	Ethanol-soluble derivative of propolis	Protect liver activity	Elimination of hepatic collagen deposition, inflammatory signals and oxidative stress	[23]
9.	Propolis extract	Vaginal use	Reorganization of vaginal mucosa faster organizational recovery decreased inflammatory response	[24]
10.	Tekirdag-Turkey propolis	Antitumor activity	Activation of caspase cascade pathway induces apoptosis in C6 glioma cells	[25]
11.	Ethanolic extract of propolis of Chihuahua (EEPCh)	Hypoglycemic effect	Significantly inhibited the increase of blood glucose and weight loss in diabetes mice	[26]

TABLE 2: Effects of different forms of propolis on wound healing.

S. No	Propolis form	Experimental model	Main results	References
1		Square skin incision	Affect stimulating keratinocytes cell proliferation	[30]
2		Diabetic foot ulcer	Reduced ulcer area enhanced wound healing	[31]
3		Pulp wound	Maintain low inflammation and microbial cell population stimulating restorative dentin Stimulates the accumulation of glycosaminoglycans on the wound surface	[32]
4	Propolis ointment	Burn	required for granulation, tissue growth and wound closure to accelerate therepair of burn tissue	[33]
5		Chronic wound	Treatment of chronic wound infection caused by <i>Proteus mirabilis</i>	[34]
6		Cell CRL-7522	Enhance the proliferation, activation and growth of skin cells	[35]
7		Rat skin wound	Speed up wound healing high inflammatory cell infiltration rate highergranulation tissue	[36]
8		Tooth pulp wound	Dentin tubules are arranged more orderly as pulp capping agent	[37]
9		Skin wound	<i>Staphylococcus aureus</i> and <i>Staphylococcus epidermidis</i> have inhibitory effects improve skin antibacterial effect	[38]
10	Propolis extract	Damage caused by striking	Contains fibroblasts and collagen a large number of mitotic cells stimulate cell proliferation and tissue repair	[39]
11		Wound (about 11 mm)	Accelerate wound healing improve collagen deposition	[40]
12	Wound dressing	Third degree burn wound	The area of the burn was reduced no inflammation and edema	[41]
13	containing propolis	Back incision	Fibroblast proliferation contributes to collagen deposition increasedsynthesis of natural hyaluronic acid	[42]
14	Propolis contains	Back and neck trauma	Wound healing improved fibroblast and collagen production increased	[43]
15	biocellulose membrane	Wound (about 6 mm)	Tissue repair of contaminated wounds shorter time and better effect	[44]
16	Propolis and metal nanoparticles and biology medical thread combination	Surgically infected wound	Promote cell migration and proliferation gives effective antibacterial power to surgical sutures	[45]

1.2. The Possible Mechanism of Propolis on Wound Healing

The mechanism of wound healing by propolis mainly includes the following five aspects: antibacterial, anti-inflammatory, antioxidant, immune, and mast cell (Figure 2). characterized by redness, heat, pain, and dysfunction. It can be infectious inflammation caused by infection or non-infectious inflammation caused by infection. After experiencing burns and trauma, severe symptoms are manifested by a large number of degenerated and necrotic tissues, bacterial invasion, massive production of free radicals, and inflammation caused by stress response [65]. Scientists have found that propolis can alleviate inflammatory problems and promote wound healing. The anti-inflammatory biological activity of propolis is mainly related to the fact that propolis contains a large number of flavonoid anti-inflammatory substances, such as carnitine and galangin, as well as phenolic anti-inflammatory substances, such as caffeic acid, ferulic acid, phenethyl caffeic acid, and so on [66]. A large number of in vivo experiments on the repair activity of propolis have observed that propolis promotes wound repair, accompanied by inhibition of local inflammatory response of wound (Table 3).

Some researchers knocked out the smad3 gene in mice, which abnormally showed accelerated skin wound healing compared with wild-type mice. The results showed that the local mononuclear cell infiltration of the mouse wound was significantly reduced, and the re-epithelialization speed increased, reflecting the inhibitory effect of the inflammatory response on the re-epithelialization [71]. Dietary propolis has an effect on the metabolism of arachidonic acid in vitro and in vivo. It significantly inhibits the lipoxygenase pathway of arachidonic acid metabolism in the process of inflammation in the body [72]. Propolis ethanol extract may exert its anti-inflammatory effect by inhibiting the expression of iNOS gene, by acting on the iNOS promoter at the NF- κ B site and directly inhibiting the catalytic activity of iNOS [73]. There is another possibility that the anti-inflammatory effect of propolis may be caused by the carbon monoxide mechanism [74].

In general, inflammatory response can play a positive role in wound repair through anti-infection, tissue debridement and release some cytokines, but it will also inhibit some aspects of tissue repair.

Antibacterial Effect. Bacterial infection is an acute systemic infection caused by pathogenic bacteria or conditional pathogenic bacteria invading the blood circulation, growing and reproducing, producing toxins and other metabolites. Clinically, it is characterized by shivering, high fever, rash, joint pain, and hepatosplenomegaly, some patients may have septic shock and transitional lesions.

Since ancient times, humans have used the antibacterial properties of propolis, including as a wound healing promoter [75]. A large number of experiments have also shown the antibacterial properties of propolis (Table 4). At present, the research on the mechanism of antibacterial activity is not clear, and it may be combined with some other biological materials to play a role. Some researchers also pointed out the antibacterial activity of propolis ingredients, polyphenols and flavonoids against *Escherichia coli* and *Staphylococcus aureus* [80]. Studies have also shown that propolis extract drug-loaded preparations can inhibit *Staphylococcus aureus*.




Figure 2: The mechanism of propolis on wound healing. IL-6—interleukin-6, iNOS—inducible nitric oxide synthase, NO—nitric oxide, TGF-β—transforming growth factor-β, VEGF—vascular endothelial growth factor, Th1—helper T 1, Th2—helper T 2, SOD—superoxide dismutase, MDA—malondialdehyde.

Table 3: Study on anti-inflammatory mechanism of propolis promoting wound healing.

S. No	Propolis administration mode	Trauma model	Action effect	References
1	External application	Burn	Accelerate the tissue repair process reduced inflammation	[49]
2	Drip	Alkali burns of rabbit cornea	Decreased infiltration of inflammatory cells	[67]
3		Six traumas in diabetic mice	Reduce neutrophil infiltration normal macrophages inwound tissue	[68]
4	Apply	Superficial second-degree burn	Reduced inflammation rapid wound healing	[69]
5		Skin wounds in diabetic mice	Increase in damage shrinkage reduction of inflammatory symptoms	[70]

2. METHODOLOGY:- (Method And Analysis)

2.1. Propolis preparation:-

Honey Spring propolis (batch number 4A80) was collected from INDIA by , MKCOPER COLLAGE OF PHARMACY, DBATU University. Collection of the propolis and determination of its contents were identified in our laboratory using high- speed counter current chromatography and off-line atmospheric pressure chemical ionization mass- spectrometry injection as previously described [32]. Briefly, preparations of propolis extract consisted of three phases including drying, extracting, and evaporating. The drying process began by washing the sample, cutting it into small pieces, and putting them in the oven with a temperature of 40-60°C. Before the extraction process, samples were dried and then crushed by a blender. 200 grams of dry samples were weighed and put in 1 L Erlenmeyer glass, soaked with ethanol to the volume of 1 L (20%). Sample in ethanol was stirred for \pm 30 minutes and allowed to stand overnight to settle. Then, solution containing the active substance was filtered with filter paper. Soaking process was repeated three times and the last stage was evaporation. Extraction solvent was inserted into 1 L evaporation flask. Then, water bath was filled with water up to a full circuit and then installed according to an equipment protocol and set to a temperature of 90°C. Ethanol was allowed to drip in the flask (\pm 1.5-2 hours/flask containing \pm 900 ml). Extraction results obtained roughly one tenth of dried natural materials (20 grams extract/200 gram's sample). A final solution equivalent to 400 mg/ml was prepared by dissolving this extract (20 grams) in 50 ml of 70% ethanol and this final solution was stored in hermetically-sealed brown-glass bottles at room temperature. Previous studies have shown that the prepared propolis extract using this method is stable for 6 months, maintaining its antimicrobial and antioxidant activities over this period [33]. According to numerous data in our laboratory, using different animal models, a daily dose of ethanolic soluble derivative of propolis (5–20 mg topically applied to wounded area) does not show any toxic effects and subsequently this dose is categorized safe. We therefore, used an optimal dose of ethanolic soluble derivative of propolis (25 μ l equivalent to 10 mg) daily applied for the treatment of normal and diabetic mice.

2.2. Chemicals:-

Streptozotocin (STZ) was obtained from Sigma Chemicals Co. (St. Louis, MO, USA). STZ was dissolved in cold 0.01 M citrate buffer (pH 4.50), which was freshly prepared (within 5 min) as needed.

2.3. Animals and diabetes induction:-

A total of 60 sexually mature 12-week-old male BALB/c mice weighing 25-30 g each were obtained from the Central Animal House of the Faculty of Pharmacy at King Saud University. All animal procedures were conducted in accordance with the standards set forth in the Guidelines for the Care and Use of Experimental Animals by the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) and the National Institutes of Health (NIH). The Animal Ethics Committee of the Zoology Department, College of Science, King Saud University approved the protocol used in this study according to the Helsinki Principles. All animals were acclimated to metal cages in a well-ventilated room for 2 weeks prior to experimentation. The animals were maintained under standard laboratory conditions (23°C, 60- 70% relative humidity and a 12-h light/dark cycle), fed a standard diet of commercial pellets and given water *ad libitum*. All mice were fasted for 20 h prior to diabetes induction. The mice ($n = 40$) were rendered diabetic by intraperitoneal (i.p.) injection of STZ (60 mg/kg body weight in 0.01 M citrate buffer pH 4.5) daily for 5 consecutive days [34, 35]; mice in the control group ($n = 20$)

were injected with vehicle alone (0.01 M citrate buffer, pH 4.5). Mice were considered diabetic if glycemia was higher than 220 mg/dl. The animals were housed for 2 weeks prior to wound formation and propolis treatment.

2.4. Excisional wound preparation and macroscopic examination:-

Two weeks post-diabetes induction, the mice were wounded as previously described [34]. Briefly, the mice were anesthetized with a single i.p. injection of ketamine (80 mg/kg body weight) and xylazine (10 mg/kg body weight). The hair on the back of each mouse was shaved, and the back was cleaned with 70% ethanol. Two wounds (8 mm in diameter, 3-4 mm apart) were made on the back of each mouse by excising the skin and underlying panniculus carnosus. The animals were then divided into three experimental groups: group 1, control non-diabetic mice topically treated with 25 μ l 70% ethanol (vehicle)/wounded area/day for 15 days (n=20); group 2, diabetic mice topically treated with 25 μ l vehicle/wounded area/day for 15 days (n=20); and group 3, diabetic mice topically treated with 25 μ l of ethanolic soluble derivative of propolis (25 μ l equivalent to 10 mg) /wounded area/day for 15 days (n=20). The optimal dose of propolis was determined in our laboratory on the basis of the LD₅₀ value and several previously established parameters. Propolis was painted onto the entire wound surface with a sterile cotton bud. Skin biopsy specimens were obtained from the animals at 3, 6, 9, 12, and 15 days post-wounding. For each time point, a skin sample

– which included the scab, the complete epithelial and dermal compartments of the wound margins, the granulation tissue, and portions of the adjacent muscle and subcutaneous fat tissue – was excised from each individual wound. As a control, a similar sample of skin was collected from the backs of non-wounded wild-type mice. Each wound site was digitally photographed at the indicated time intervals to document the wound area. Changes in the wound area are expressed as a percentage of the initial wound area. At the indicated time intervals, the tissue of two wounds from ten animals (n = 20 wounds) was collected for RNA, western blot and ELISA analyses.

2.5. Measurement of hydroxyproline content in the wound sites:-

At the indicated time intervals after the injury, skin wound sites were removed using a sterile disposable biopsy punch and were dried for 24 h at 120°C, the levels of hydroxyproline, a major constituent of collagen in skin wound sites, was measured to quantify collagen accumulation at the wound site, as previously described [36]. Hydroxyproline content was calculated by comparison to standards and is expressed as the amount (μ g) per wounded tissue weight.

2.6. Blood analysis:-

Blood glucose levels were determined using an AccuTrend sensor (Roche Biochemicals; Mannheim, Germany). Luminex (Biotrend; Düsseldorf, Germany) was used to analyze serum insulin levels, according to the manufacturer's instructions.

2.7. Biochemical analysis of wounded tissues:-

Measuring cytokine levels. A 2.0-mm punch biopsy taken from the wound site was harvested and frozen in liquid nitrogen. Specimens were homogenized in cytoplasmic lysis buffer containing protease inhibitors (Roche Diagnostics), disrupted using Fast Prep (Q-Biogene; Solon, OH, USA), and centrifuged at 5000 \times g for 10 min. The protein concentration in each lysate was determined using the bicinchoninic acid (BCA) protein assay kit (Pierce; Rockford, IL, USA). The supernatants were then assayed to determine IL-1 β , IL-6, TNF- α , TGF- β 1, MMP2 and MMP9 levels using a commercial ELISA kit (R&D Systems; France), according to the manufacturer's instructions. The results are expressed as target molecule (picograms) per total protein (milligrams) for each sample.

2.8. Western blot analysis:-


The skin and wound tissue biopsies were homogenized in lysis buffer (1% Triton X-100, 137 mM NaCl, 10% glycerol, 1 mM dithiothreitol, 10 mM NaF, 2 mM Na_3VO_4 , 5 mM ethylenediaminetetra-acetic acid, 1 mM phenylmethylsulfonylfluoride, 5 ng/ml aprotinin, 5 ng/ml leupeptin and 20 mM Tris/HCl, pH 8.0), and the lysates were prepared as previously described [37]. Fifty micrograms of total protein from the skin lysates was analyzed using SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and western blot analysis. Antibodies (Abs) directed against anti-collagen type 1 antibody (1:5000) phospho-Smad2 (pSmad2) (1:1000), pSmad3 (1:1000), total Smad2 (1:2000), total Smad3 (1:2000) and β -actin (1:4000) (all from Abcam, Paris, France) were used in combination with horseradish peroxidase-conjugated secondary Abs, and the proteins were visualized using an enhanced chemiluminescence (ECL, Supersignal Westpico chemiluminescent substrate; Perbio, Bezons, France) detection system. The ECL signal was detected on Hyperfilm ECL. To quantify the band intensities, the films were scanned, saved as TIFF files, and then analyzed using the NIH Image J software program.

2.9. Statistical analysis:-

The data were tested for normality (using an Anderson-Darling test) and variance homogeneity prior to further statistical analysis. The data were normally distributed and are expressed as the mean \pm SEM (standard error of the mean). Significant differences between groups were analyzed using one-way analysis of variance (for more than two groups) followed by Tukey's post-test using SPSS software version

17. Data are expressed as the mean \pm SEM. Differences were considered statistically significant at $^*P < 0.05$ for diabetic vs. control; $^+P < 0.05$ for diabetic + propolis vs. control; or $^\#P < 0.05$ for diabetic + propolis vs. diabetic.

3. Modeling and Analsis



4. Results And Discussion

4.1. Results

4.1.1. Propolis enhances wound closure in diabetic mice

We evaluated macroscopic changes at the skin-excision wound sites in control mice, diabetic mice and diabetic mice treated topically with the ethanolic soluble derivative of propolis. Pictures were taken on day 0, immediately following injury. The wound sites exhibited a similar morphology in all 3 experimental groups on day 3 post-injury. The wounds in the control and diabetic mice treated with propolis showed nearly similar degree of closure at 15 days post-injury. By contrast, the diabetic mice exhibited delayed wound closure. A representative result is shown (Fig. 1A). The accumulated data showing the change in the percentage of wound closure at each time point (relative to the original wound area) from 10 mice per group are shown (Fig. 1B). These results showed that wound closure and healing were accelerated in diabetic mice treated topically with the ethanolic soluble derivative of propolis compared with diabetic mice treated with vehicle alone. We also measured the blood glucose and insulin levels in the 3 groups of mice before and throughout the indicated time points post-wounding. Accumulated data from 10 animals from each group revealed that the diabetic mice exhibited significant elevation in the glucose levels and significant decrease in the insulin levels compared to the control non-diabetic mice (Table 1). However, when the diabetic mice were treated with the ethanolic soluble derivative of propolis the blood glucose levels were not significantly decreased and the insulin levels were not significantly increased compared to the diabetic mice treated with vehicle (Table 1).

4.1.2. Topical application of propolis to diabetic wounds restores the levels of wound-tissue pro-inflammatory cytokines, TGF- β 1 and MMP9

ELISAs were used to measure the levels of pro-inflammatory cytokines, TGF- β 1 and MMP9, which play important roles in wound healing, in the excisional wound tissues collected from the 3 groups of mice on days 3, 6, 9, 12 and 15 post-wounding. Day 0 samples were collected one hour prior to wound formation (non-wounded skin tissue). The accumulated data from 10 individual mice from each group revealed elevated levels of pro-inflammatory cytokines (IL-1 β , IL-6, and TNF- α) and MMP9 in diabetic mice compared with controls, as shown in Fig. 2. In particular, the diabetic mice exhibited elevated levels of the pro-inflammatory cytokines IL-1 β (Fig. 2A), IL-6 (Fig. 2B) and TNF- α (Fig. 2C), as well as MMP9 (Fig. 2D), compared with control non-diabetic animals. Interestingly, the diabetic mice that were treated topically with the ethanolic soluble derivative of propolis showed a partial but significant restoration of wild-type expression for IL-1 β (Fig. 2A), IL-6 (Fig. 2B), TNF- α (Fig. 2C), and MMP9 (Fig. 2D) compared with vehicle-treated diabetic animals. Nevertheless, in contrast to the levels of pro-inflammatory cytokines and MMP9, diabetic mice exhibited a marked and significant reduction in the levels of TGF- β 1 compared with control non-diabetic mice (Fig. 2E). Most importantly, the diabetic mice that were treated topically with the ethanolic soluble derivative of propolis showed a partial but significant restoration of TGF- β 1 levels compared with vehicle-treated diabetic animals.

4.1.3. Propolis-treated diabetic mice exhibit clear changes in TGF- β 1-mediated phosphorylation of Smad2 and Smad3 in the wound tissue.

TGF- β 1 mediates activation of the transcription factors Smad2 and Smad3, leading to the production of collagen, which in turn acts as a regulator of the wound healing processes. Therefore, the expression and activation of Smad2 and Smad3 were investigated in the excisional wound tissues from the three groups of mice at 6, 9 and 12 days post-wounding. A representative immunoblot showing one of the five experiments is shown (Fig. 3A & B). In particular, immunoblots of phosphorylated Smad2, total Smad2 and total β -actin (loading control) (Fig. 3A), phosphorylated Smad3, total Smad3 and total β -actin (loading control) (Fig. 3B) are shown in control non-diabetic mouse (CM), vehicle-treated diabetic mouse (DM) and diabetic mouse treated with propolis (DM+P) in wounded tissues at 6, 9 and 12 days post-wounding. We found that the phosphorylation of Smad2 and Smad3 were markedly reduced in vehicle-

treated diabetic mice, as the immunoblots showed very low intensity bands over the entire wound healing process compared with control non-diabetic mice. By contrast, the propolis-treated diabetic mice displayed enhanced phosphorylation of Smad2 and Smad3, which peaked at 9 and 12 days post-injury, similar to the levels observed in the control non-diabetic mice. Nevertheless, treatment of diabetic mice with propolis had no effect on the expression of Smad2 and Smad3. Accumulated data from five individual mice per group are shown for the normalized phosphorylation of Smad2 to the total relevant Smad2 (Fig. 3C) and normalized phosphorylation of Smad3 to the total relevant Smad3 (Fig. 3D). We found that vehicle-treated diabetic mice (closed black bars) exhibited a significant reduction in the normalized Smad2 and Smad3 phosphorylation from days 6 to 12 post-wounding compared with control (open bars) and propolis-treated diabetic mice (gray dotted bars).

4.1.4. Propolis enhances the production of collagen and accelerates diabetic wound healing

Increased collagen content in the extracellular matrix is a characteristic change observed during the proliferative phase of the wound healing process. As hydroxyproline is found almost exclusively in collagen, we used hydroxyproline content as an indicator of collagen type I levels at the wound sites. The accumulated data from 10 individual mice per group demonstrated that hydroxyproline content was significantly decreased in diabetic mice with decreased wound closure compared with control mice (Fig. 4A). Compared with control mice, there was less collagen accumulation at wound sites in diabetic mice, consistent with the delays in wound healing. However, diabetic mice that were treated topically with the ethanolic soluble derivative of propolis exhibited a significant restoration of hydroxyproline content compared with vehicle-treated diabetic mice. To confirm the alteration in the expression of collagen type I in the diabetic animals when treated topically with the ethanolic soluble derivative of propolis, Western blot analysis was assessed to monitor the expression of collagen type I at its protein level and one representative immunoblot showing one of the five experiments is shown (Fig. 4B) in control non-diabetic mouse (CM), vehicle-treated diabetic mouse (DM) and diabetic mouse treated with propolis (DM+P) in wounded tissues at 6, 9 and 12 days post-wounding. Our data demonstrated that the expression of collagen type I was markedly reduced in vehicle-treated diabetic mice, as the immunoblots showed very low intensity bands compared with control non-diabetic mice. By contrast, the propolis-treated diabetic mice displayed an obvious increase in the expression of collagen type I at 9 and 12 days post-injury, similar to the levels observed in the control non-diabetic mice. Accumulated data from five individual mice per group are shown (Fig. 4C) for the normalized values of collagen type I expression to the relative intensities of β -actin values. Our data revealed that vehicle-treated diabetic mice (closed black bars) exhibited a significant reduction in the collagen type I expression (from days 6 to 12 post-wounding compared with control (open bars) and propolis-treated diabetic mice (gray dotted bars). Interestingly, when diabetic mice were treated topically with the ethanolic soluble derivative of propolis, they exhibited a significant restoration of collagen type I expression.




Fig.4.A. Macroscopic changes in skin excisional wounds during wound closure. (A) The wound sites were photographed at the indicated intervals. The pictures at day 0 were taken immediately after injury. (B) Re- presentative data from 10 individual mice per group indicating changes in the percentage of wound closure. The data are presented as mean ± SEM values. *P < 0.05, diabetic vs. control; +P < 0.05, diabetic + propolis vs. control; #P < 0.05, diabetic + propolis vs. diabetic.



Table 4.B. Blood glucose and insulin level

	Blood glucose level (mg/dL)		
	Control mice	Diabetic mice	Diabetic + propolis
Onset of diabetic induction	120±11	260±15.8 *	244±16.4 +
Two weeks post-diabetes induction	127±12	344±18.2 *	315±18.4 +
0 day post-wounding	101±10	360±16 *	311±17.2 +
3 days post-wounding	90±9.8	368±19 *	339±18 +
6 days post-wounding	131±13.8	370±22 *	353±21 +
9 days post-wounding	128±11.6	345±21 *	326±19 +
12 days post-wounding	162±12.4	392±21.6 *	362±20.8 +
15 days post-wounding	139±14.4	386±19.4 *	359±21 +
	Blood insulin level (ng/ml)		
	Control mice	Diabetic mice	Diabetic + propolis
Onset of diabetic induction	5.6±0.5	3.1±0.29 *	3.7±0.28 +
Two weeks post-diabetes induction	5.7±0.36	2.2±0.22 *	2.6±0.26 +
0 day post-wounding	4.9±0.32	1.6±0.18*	1.7±0.21 +
3 days post-wounding	5.1±0.4	1.8±0.19 *	1.9±0.18 +
6 days post-wounding	5.2±0.4	1.9±0.22 *	1.95±0.19 +
9 days post-wounding	4.9±0.33	2.1±0.2 *	2.3±0.2 +
12 days post-wounding	5.4±0.45	2.1±0.16 *	2.2±0.25 +
15 days post-wounding	4.98±0.35	2.05±0.18 *	2.4±0.16 +

*P<0.05 Diabetic vs control.
+P<0.05 Diabetic + propolis vs control.

the experiments

Fig. 4.C. Changes in the levels of pro-inflammatory cytokines and MMP9 during wound healing. The levels of pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α and TGF- β) and MMP9 were measured in the indicated days post-wounding. The results are presented as cy values ($n = 10$). $P < 0.05$, diabetic vs. control; * $P < 0.05$, diabetic






Fig. 4.D. Topical application of propolis affects the phosphorylation of Smad2 and Smad3 in wounded tissues. Immunoblots showing one representative experiment is shown for phosphorylated Smad2 and total Smad2 (A), phosphorylated Smad3 and total Smad3 (B) in control non-diabetic mouse (CM), diabetic mouse (DM) and diabetic mouse treated topically with propolis (DM+P). Accumulated data from five independent experiments are shown for normalized Smad2 (C) and Smad3 (D) phosphorylation in control non-diabetic (open bars), diabetic (closed black bars) and diabetic animals treated topically with propolis (gray dotted bars) in wounded skin samples (at 6, 9 and 12 days post-wounding). Values represent the mean \pm SEM. * $P < 0.05$, diabetic vs. control; # $P < 0.05$, diabetic + WP vs. control; + $P < 0.05$, diabetic + WP vs. diabetic (ANOVA with Tukey's post-test).

Fig. 4.E. Propolis enhances collagen production in diabetic wounds.


(A) Hydroxyproline content, an index of collagen accumulation at wound sites, was determined. Values represent the mean \pm SEM. (B) Western blot analysis was performed to measure the expression of collagen type I as a protein and β -actin in wounded skin samples (at 6, 9 and 12 days post-wounding). (C) Accumulated data from five independent experiments are shown for normalized collagen type I expression in control non-diabetic (open bars), diabetic (closed black bars) and diabetic animals treated topically with propolis (gray dotted bars) in wounded skin samples (at 6,

9 and 12 days post-wounding).

Values represent the mean \pm SEM.

* $P < 0.05$, diabetic vs. control;

* $P < 0.05$, diabetic + WP vs. control; # $P < 0.05$, diabetic + WP vs. diabetic (ANOVA with Tukey's post-test).



4.2. Discussion

Natural antioxidants play central roles in enhancing the immune system through mechanisms dependent on the oxidative stress which, in turn, seems to play significant roles in many human diseases. In this context, we previously demonstrated the beneficial effects of thymoquinone in the treatment of multiple myeloma and improving the diabetic complications by restoring the T cell immune response in diabetic offspring [37–40]. Most interesting, we shown that natural antioxidants isolated from ants venoms were able to enhance the normal lymphocyte functions and exerts antitumor effects on the breast cancer cells [41]. Moreover, we provide clear evidences for the effects of camel whey protein for accelerating the healing process of diabetic wounds in experimental animal models [34, 42, 43]. Propolis is a natural antioxidant product found in plant materials and is processed by worker bees.

The present study showed that the topical application of propolis to diabetic wounds in mice accelerated wound closure to a rate that was similar to that in non-diabetic control mice and significantly faster than that in untreated diabetic mice. These improvements were evident during the earliest stages post-incision and continued over the entire two-week study period, indicating that propolis application impacts all stages of the healing process. Our data are consistent with previous studies in humans and

animals, as well as with older reports describing the use of propolis to treat ulcers [28-30]. Topical application of propolis had no effect on the glucose levels as shown in Table 1 suggesting that treatment of diabetic wounds with propolis accelerates the healing process by reversing the diabetic complication rather than lowering the glucose level.

The wound repair-enhancing effects of propolis are partly due to its anti-inflammatory properties. It is thought that prolonged inflammation impairs the healing process in diabetic patients, and it is recognized that elevated IL-1 β , IL-6, and TNF- α levels are found in diabetic wounds [34, 44]. Therefore, it has been suggested that targeting inflammatory mediators could be an effective strategy for improving healing dynamics in diabetes. In the present study, we show that propolis application abrogated the inflammatory process associated with diabetic wounds and restored the expression of IL-1 β , IL-6 and TNF- α to near wild-type levels. A direct inhibitory effect of propolis on cytokine production by immune cells has been documented [45], supporting the anti-inflammatory profile shown in the current study. Restoration of the proper expression levels of these cytokines is likely associated with inhibition of the inflammation feedback loop [45], as well as with decreased degradation of the ECM through the inhibition of MMP expression [46, 47]. Indeed, MMP9 levels were correlated with pro-inflammatory cytokine levels in this study, consistent with the hypothesis that propolis reduces proteolytic activity during cutaneous inflammation. It was previously reported that propolis and its extracts reduce MMP9 expression in diabetic wounds, particularly TNF-induced MMP9 [30]. In particular, flavonoids inhibited TNF- α -induced MMP9 expression in keratinocytes at both the gene and protein levels [48].

Propolis increased the levels of both collagen type I and its major constituent hydroxyproline in the wounds of diabetic mice. This finding was consistent with the accelerated healing observed in the treatment group compared with the diabetic PBS-treated group, which showed reduced expression of collagen type I and hydroxyproline. Similar healing profiles in the ECM (i.e., increased levels of collagen and its degradation products) were reported in rat excisional [28] and burn wounds [49] following propolis application. Additionally, enhanced healing and early stage replacement of collagen type III with collagen type I in burn wounds have been reported using propolis extract-coated collagen dressings [50]. Diminished collagen deposition is a mechanism of delayed wound healing, and propolis can restore the composition and quality of the ECM. These improvements may be due to reduced MMP9 levels, as described above, and indeed, the restoration of proper MMP9 levels using laser therapy improved ECM quality in diabetic mice [50]. However, in our study, increased collagen type I was observed at the level of gene expression, as well as at the protein level, as demonstrated by higher hydroxyproline values.

To further define the role of TGF- β in enhancing wound healing following propolis application, we characterized TGF- β 1 gene expression at mRNA and protein levels, as well as phosphorylation of the downstream Smad transcription factors. Downregulation of the Smad2/3 signaling pathway is correlated with decreased collagen type I transcription in fibroblast cell lines [51]. Additionally, multiple studies using a variety of pharmacological and physical agents have demonstrated the induction of collagen expression and synthesis in dermal fibroblasts in a TGF- β /Smad-dependent manner [52-54]. Therefore, a means to positively target the TGF- β pathway would certainly be useful for the treatment of diabetic wounds. Strikingly, propolis application to diabetic wounds resulted in the upregulation of TGF- β gene expression and enhanced Smad2/3 phosphorylation compared with Vehicle-treated diabetic wounds, and these effects were observed from the earliest stages, peaking around day 9 of the study. In addition to the transcriptional activation of collagen expression, the downregulation of both TNF- α and MMP9 in the cutaneous wounds of diabetic mice is consistent with increased TGF- β /Smad signaling, given that the upregulation of TGF- β /Smad is accompanied by the downregulation of TNF- α and MMP9 as well as decreased inflammation. Indeed, the promoters of both factors harbor TGF- β /Smad response elements, and TGF- β has been shown to downregulate TNF- α -induced MMP9 expression in monocytes [55]. Other potential mechanisms through which propolis application could promote wound healing via TGF- β /Smad upregulation are the stimulation of keratinocyte migration and increased integrin expression [56], which will require further investigation.

It has been shown that HoxD3, a homeobox transcription factor that promotes angiogenesis and collagen synthesis, is up-regulated during normal wound repair whereas its expression is diminished in impaired healing wounds of the genetically diabetic (db/db) mouse as compared to wild-type mice with normal healing wounds [57]. In this context, it was concluded that decreased expression of HoxD3 was due to marked reduction in the expression of cytokines and growth factors including TGF- β , which are all reduced in diabetic models. [57]. Therefore, it is possible that treatment of diabetic wounds with propolis restored TGF- β expression and signaling-mediated HoxD3 and Smad and subsequently collagen expression.

It has been established that TGF- β induced proliferation of keratinocytes and fibroblasts, led to new formation of capillaries in the granulation tissue and modulated extracellular matrix deposition and reconstitution of the injured area. Additionally, topical application of growth factors was successful to accelerate healing of full thickness wound in normal mice and normalizes a delayed healing response of diabetic rats. Miyazono and Heldin reported that TGF- β is a family of multifunctional 25KDa protein (TGF- β 1, 2, 3) which stimulates collagen and fibronectin formation in variety of fibroblast cell lines [58]. Moreover, TGF- β is known to regulate the differentiation of cells, induce chemotaxis of inflammatory cells and induce the accumulation of extracellular matrix protein [59]. Sporn et al. stated that TGF- β is a human DNA-derived polypeptide growth factor that induces normal soft tissue repair mechanism and reverses deficient repair rates. This growth factor is released by platelets, monocytes/macrophages, endothelial cells and fibroblasts, cells that are essential to the repair process [60]. In this work by Sporn et al. they found that TGF- β played a central role in wound healing. It influenced the inflammatory response, angiogenesis, granulation tissue formation, reepithelization, extracellular matrix deposition and remodeling. Therefore, it is possible the effects of propolis in accelerating healing process of diabetic wounds was due to its direct effect on the expression of TGF- β and its downstream signaling. The present study indicates that the topical application of propolis enhances the wound repair process in the context of diabetes by promoting TGF- β /Smad signaling, leading to increased expression and deposition of collagen type I, reduced MMP expression, and decreased inflammation.

5. Abbreviations

Diabetes mellitus (DM); Extracellular matrix (ECM); Interleukin (IL); Matrix metalloproteinase (MMP); Streptozotocin (STZ); Transforming growth factor-beta (TGF- β); Tumor necrosis factor-alpha (TNF-a).

6. Conclusions

When people realize the benefits of propolis to human beings, scientists at home and abroad have done a lot of research on propolis. The chemical components of propolis are very complex, and different chemical components play different roles. At present, propolis has been widely used in food, health products, cosmetics, and beauty products and has a broad market and application value.

Propolis is rich in flavonoids, polyphenols, terpenoids, aromatic acids, and other pharmacological active ingredients. Flavonoids can promote the synthesis of collagen, and flavonoids and other ingredients also have antibacterial and anti-inflammatory functions. In skin wound healing, propolis can reduce scar formation, shorten healing time, increase wound contraction, accelerate tissue repair, and ultimately improve the quality of life of patients. It can be seen that the importance of propolis to the human body is extreme. In order for my country's propolis products to win more shares in the international market, it is necessary to conduct more in-depth discussions on some issues in the research, development and application of propolis. For example, first, there is little research on the combination of propolis and other substances. In this regard, it can be increased to use propolis with low toxicity with other drugs to play a greater medicinal value. Second, propolis can be made into different dosage forms for clinical use. At present, there is little research on intelligent materials and nano-materials using propolis, which will be a very important research direction in the future research. Third, propolis can be further purified and optimized, and the role of each effective substance can be brought into full play.

At present, some of the mechanism of propolis is not perfect, and there are many directions worthy of research and discussion. This article only reviews the effects and mechanisms of propolis on wound healing and the effects of propolis and other compounds in order to provide more effective and comprehensive information and provide some ideas for the development and utilization of propolis in the future.

7. References

- [1] E. Farstvedt, T. S. Stashak, and A. Othic, "Update on topical wound medications," *Clinical Techniques in Equine Practice*, vol. 3, no. 2, pp. 164–172, 2004.
- [2] J. Chen, C. M. Han, X. W. Lin, Z. J. Tang, and S. J. Su, "Effect of silver nanoparticle dressing on second degree burn wound," *Zhonghua Wai Ke Za Zhi*, vol. 44, no. 1, pp. 50–52, 2006.
- [3] P. Olczyk, G. Wisowski, K. Komosinska-Vassev et al., "Propolis modifies collagen types I and III accumulation in the matrix of burnt tissue," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 423809, 10 pages, 2013.
- [4] G. Sandri, M. C. Bonferoni, F. D'Autilia et al., "Wound dressings based on silver sulfadiazine solid lipid nanoparticles for tissue repairing," *European Journal of Pharmaceutical and Biopharmaceutics*, vol. 84, no. 1, pp. 84–90, 2013.
- [5] I. O. W. Leitch, A. Kucukcelebi, and M. C. Robson, "Inhibition of wound contraction by topical antimicrobials," *ANZ Journal of Surgery*, vol. 63, no. 4, pp. 289–293, 1993.
- [6] M. J. Muller, M. A. Hollyoak, Z. Moaveni, T. L. H. Brown, D. Herndon, and J. Heggors, "Retardation of wound healing by silver sulfadiazine is reversed by aloe vera and nystatin," *Burns*, vol. 29, no. 8, pp. 834–836, 2003.
- [7] L. G. Wasef, H. M. Shaheen, Y. S. El-Sayed et al., "Effects of silver nanoparticles on burn wound healing in a mouse model," *Biological Trace Element Research*, vol. 193, no. 2, pp. 456–465, 2020.
- [8] S. Iyyam Pillai, P. Palsamy, S. Subramanian, and M. Kandaswamy, "Wound healing properties of Indian propolis studied on excision wound-induced rats," *Pharmaceutical Biology*, vol. 48, no. 11, pp. 1198–1206, 2010.
- [9] M. L. Khalil, "Biological activity of bee propolis in health and disease," *Asian Pacific Journal of Cancer Prevention*, vol. 7, no. 1, pp. 22–31, 2006.
- [10] M. F. Tolba, A. T. El-Serafi, and H. A. Omar, "Caffeic acid phenethyl ester protects against glucocorticoid-induced osteoporosis in vivo: impact on oxidative stress and RANKL/OPG signals," *Toxicology and Applied Pharmacology*, vol. 324, pp. 26–35, 2017.
- [11] B. Romana-Souza, J. S. Dos Santos, and A. Monte-Alto-Costa, "Caffeic acid phenethyl ester promotes wound healing of mice pressure ulcers affecting NF- κ B, NOS2 and NRF2 expression," *Life Sciences*, vol. 207, pp. 158–165, 2018.
- [12] M. M. Nichitai, A. M. Josceanu, R. D. Isopescu et al., "Polyphenolics profile effects upon the antioxidant and antimicrobial activity of propolis extracts," *Scientific Reports*, vol. 11, no. 1, Article ID 20113, 2021.
- [13] F. Koc, M. Y. Tekeli, M. Kanbur, M. Ö Karayigit, and B. C. Liman, "The effects of chrysin on lipopolysaccharide-induced sepsis in rats," *Journal of Food Biochemistry*, vol. 44, no. 9, Article ID e13359, 2020.
- [14] H. Arjun, "Banskota and Yasuhiro Tezuka and Shigetoshi Kadota recent progress in pharmacological research of propolis," *Phytotherapy Research*, vol. 15, no. 7, pp. 561–571, 2001.
- [15] J. B. Daleprane, V. da Silva Freitas, A. Pacheco et al., "Anti-atherogenic and anti-angiogenic activities of polyphenols from propolis," *The Journal of Nutritional Biochemistry*, vol. 23, no. 6, pp. 557–566, 2012.
- [16] R. El Adaouia Taleb, N. Djebli, H. Chenini, H. Sahin, and

- S. Kolayli, "In vivo and in vitro anti-diabetic activity of ethanolic propolis extract," *Journal of Food Biochemistry*, vol. 44, no. 7, Article ID e13267, 2020.
- [17] D. Skaba, T. Morawiec, M. Tanasiewicz et al., "Influence of the toothpaste with Brazilian ethanol extract propolis on the oral cavity health," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 215391, 12 pages, 2013.
- [18] K. Anjaly and A. B. Tiku, "Caffeic acid phenethyl ester induces radiosensitization via inhibition of DNA damage repair in androgen-independent prostate cancer cells," *Environmental Toxicology*, vol. 37, no. 5, pp. 995–1006, 2022.
- [19] N. Paulino, L. A. Coutinho, J. R. Coutinho, G. C. Vilela, V. P. D. Silva Leandro, and A. S. Paulino, "Antitumorogenic effect of Brazilian propolis formulation in mice," *Pharmacology & Pharmacy*, vol. 6, no. 12, pp. 580–588, 2015.
- [20] A. Sabir and A. Sumidarti, "Interleukin-6 expression on inflamed rat dental pulp tissue after capped with *Trigona* sp. propolis from south Sulawesi, Indonesia," *Saudi Journal of Biological Sciences*, vol. 24, no. 5, pp. 1034–1037, 2017.
- [21] C. Meneghelli, L. S. D. Joaquim, G. L. Q. Félix et al., "Southern Brazilian autumnal propolis shows anti-angiogenic activity: an in vitro and in vivo study," *Microvascular Research*, vol. 88, pp. 1–11, 2013.
- [22] M. A. Alsherbiny, D. J. Bhuyan, I. Radwan, D. Chang, and C. G. Li, "Metabolomic identification of anticancer metabolites of Australian propolis and proteomic elucidation of its synergistic mechanisms with doxorubicin in the MCF7 cells," *International Journal of Molecular Sciences*, vol. 22, no. 15, p. 7840, 2021.
- [23] G. Badr, E. A. Sayed, H. Waly, K. A. H. Hassan, M. H. Mahmoud, and Z. Selamoglu, "The therapeutic mechanisms of propolis against CCl₄-mediated liver injury by mediating apoptosis of activated hepatic stellate cells and improving the hepatic architecture through PI3K/AKT/mTOR, TGF- β /Smad2, Bcl2/BAX/P53 and iNOS signaling pathways," *Cellular Physiology and Biochemistry*, vol. 53, no. 2, pp. 301–322, 2019.
- [24] A. P. Bonfim, K. M. Sakita, D. R. Faria et al., "Preclinical approaches in vulvovaginal candidiasis treatment with mucoadhesive thermoresponsive systems containing propolis," *PLoS One*, vol. 15, no. 12, Article ID e0243197, 2020.
- [25] Z. M. Coskun, M. Ersoz, M. Gecili, A. Ozden, and A. Acar, "Cytotoxic and apoptotic effects of ethanolic propolis extract on C6 glioma cells," *Environmental Toxicology*, vol. 35, no. 7, pp. 768–773, 2020.
- [26] N. Rivera-Yañez, M. Rodriguez-Canales, O. Nieto-Yañez et al., "Hypoglycaemic and antioxidant effects of propolis of chihuahua in a model of experimental diabetes," *Evidence-Based Complementary and Alternative Medicine*, vol. 2018, Article ID 4360356, 10 pages, 2018.
- [27] C. Ji, Y. Pan, S. Xu et al., "Propolis ameliorates restenosis in hypercholesterolemia rabbits with carotid balloon injury by inhibiting lipid accumulation, oxidative stress, and TLR4/NF- κ B pathway," *Journal of Food Biochemistry*, vol. 45, no. 4, Article ID e13577, 2021.
- [28] M. Barary, R. Hosseinzadeh, S. Kazemi et al., "The effect of propolis on 5-fluorouracil-induced cardiac toxicity in rats," *Scientific Reports*, vol. 12, no. 1, p. 8661, 2022.
- [29] M. Y. Song, D. Y. Lee, and E. H. Kim, "Anti-inflammatory and anti-oxidative effect of Korean propolis on helicobacter pylori-induced gastric damage in vitro," *Journal of Microbiology*, vol. 58, no. 10, pp. 878–885, 2020.
- [30] E. Sehn, L. Hernandez, S. L. Franco, C. Goncalves, and M. Baesso, "Dynamics of reepithelialisation and penetration rate of a bee propolis formulation during cutaneous wounds healing," *Analytica Chimica Acta*, vol. 635, no. 1, pp. 115–120, 2009.
- [31] M. Afkhamizadeh, R. Aboutorabi, H. Ravari et al., "Topical propolis improves wound healing in patients with diabetic foot ulcer: a randomized controlled trial," *Natural Product Research*, vol. 32, no. 17, pp. 2096–2099, 2018.
- [32] W. A. Bretz, D. J. Chiego Jr., M. C. Marcucci, I. Cunha, A. Custodio, and L. G. Schneider, "Preliminary report on the effects of propolis on wound healing in the dental pulp," *Zeitschrift Für Naturforschung C*, vol. 53, no. 11-12, pp. 1045–1048, 1998.
- [33] P. Olczyk, K. Komosinska-Vassev, K. Winsz-Szczotka, J. Stojko, K. Klimek, and E. M. Kozma, "Propolis induces chondroitin/dermatan sulphate and hyaluronic acid accumulation in the skin of burned wound," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 290675, 8 pages, 2013.
- [34] J. Kwiecińska-Piróg, K. Skowron, A. Śniegowska et al., "The impact of ethanol extract of propolis on biofilm forming by proteus mirabilis strains isolated from chronic wound infections," *Natural Product Research*, vol. 33, no. 22, pp. 3293–3297, 2019.
- [35] A. Jacob, A. Parolia, A. Pau, and F. Davamani Amalraj, "The effects of Malaysian propolis and Brazilian red propolis on connective tissue fibroblasts in the wound healing process," *BMC Complementary and Alternative Medicine*, vol. 15, no. 1, p. 294, 2015.
- [36] O. D. Eyarefe, C. A. Ozota, and T. A. Jarikre, "Pathological and immunohistochemical evaluation of wound healing potential of Nigerian bee propolis in albino rats," *Comparative Clinical Pathology*, vol. 28, 2018.
- [37] N. Likitpongpipat, S. Sangmaneeadet, P. Klanrit, R. Noisombut, S. Krisanaprakornkit, and P. Chailertvanitkul, "Promotion of dental pulp wound healing in New Zealand white rabbits' teeth by Thai propolis product," *Journal of Veterinary Dentistry*, vol. 36, no. 1, pp. 17–24, 2019.
- [38] K. Huanbutta, W. Sittikijyothin, and T. Sangnim, "Development of topical natural based film forming system loaded propolis from stingless bees for wound healing application," *Journal of Pharmaceutical Investigation*, vol. 50, 2020.
- [39] A. A. Berretta, A. P. Nascimento, P. C. P. Bueno, M. M. de Oliveira Lima Leite Vaz, and J. M. Marchetti, "Propolis standardized extract (EPP-AF), an innovative chemically and biologically reproducible pharmaceutical compound for treating wounds," *International Journal of Biological Sciences*, vol. 8, no. 4, pp. 512–521, 2012.
- [40] A. Eskandarinia, A. Kefayat, M. Gharakhloo et al., "A propolis enriched polyurethane-hyaluronic acid nanofibrous wound dressing with

- remarkable antibacterial and wound healing activities,” *International Journal of Biological Macromolecules*, vol. 149, pp. 467–476, 2020.
- [41] M. Stojko, J. Włodarczyk, M. Sobota et al., “Biodegradable electrospun nonwovens releasing propolis as a promising dressing material for burn wound treatment,” *Pharmaceutics*, vol. 12, no. 9, p. 883, 2020.
- [42] A. Eskandarinia, A. Kefayat, M. Rafenia, M. Agheb, S. Navid, and K. Ebrahimpour, “Cornstarch-based wound dressing incorporated with hyaluronic acid and propolis: in vitro and in vivo studies,” *Carbohydrate Polymers*, vol. 216, pp. 25–35, 2019.
- [43] F. Marquele-Oliveira, H. da Silva Barud, E. C. Torres et al., “Development, characterization and pre-clinical trials of an innovative wound healing dressing based on propolis (EPP- AF)-containing self-microemulsifying formulation incorporated in biocellulose membranes,” *International Journal of Biological Macromolecules*, vol. 136, pp. 570–578, 2019.
- [44] H. D. S. Barud, A. M. de Araújo Júnior, S. Saska et al., “Antimicrobial Brazilian propolis (EPP-AF) containing biocellulose membranes as promising biomaterial for skin wound healing,” *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 703024, 10 pages, 2013.
- [45] T. Baygar, “Characterization of silk sutures coated with propolis and biogenic silver nanoparticles (AgNPs); an eco-friendly solution with wound healing potential against surgical site infections (SSIs),” *Turkish Journal of Medical Sciences*, vol. 50, no. 1, pp. 258–266, 2020.
- [46] J. M. Reinke and H. Sorg, “Wound repair and regeneration,” *European Surgical Research*, vol. 49, no. 1, pp. 35–43, 2012.
- [47] S. Bruno, C. C. Grange, F. Collino et al., “Microvesicles derived from mesenchymal stem cells enhance survival in a lethal model of acute kidney injury,” *PLoS One*, vol. 7, no. 3, Article ID e33115, 2012.
- [48] S. Gatti, S. Bruno, M. C. Deregis et al., “Microvesicles derived from human adult mesenchymal stem cells protect against ischaemia-reperfusion-induced acute and chronic kidney injury,” *Nephrology Dialysis Transplantation*, vol. 26, no. 5, pp. 1474–1483, 2011.
- [49] A. G. T. Pessolato, D. D. S. Martins, C. E. Ambrósio, C. A. F. Mancanares, and A. F. de Carvalho, “Propolis and amnion reepithelialise second-degree burns in rats,” *Burns*, vol. 37, no. 7, pp. 1192–1201, 2011.
- [50] E. B. de Almeida, J. Cordeiro Cardoso, A. Karla de Lima et al., “The incorporation of Brazilian propolis into collagen-based dressing films improves dermal burn healing,” *Journal of Ethnopharmacology*, vol. 147, no. 2, pp. 419–425, 2013.
- [51] P. Olczyk, K. Komosińska-Vassev, K. Winsz-Szczotka et al., “Propolis modulates vitronectin, laminin, and heparan sulfate/heparin expression during experimental burn healing,” *Journal of Zhejiang University-Science B*, vol. 13, no. 11, pp. 932–941, 2012.
- [52] M. Bayrami, A. Bayrami, A. Habibi-Yangjeh et al., “Biologically-synthesised ZnO/CuO/Ag nanocomposite using propolis extract and coated on the gauze for wound healing applications,” *IET Nanobiotechnology*, vol. 14, no. 7, pp. 548–554, 2020.
- [53] O. D. Eyarefe, C. A. Ozota, T. A. Jarikre, and B. O. Emikpe, “Pathological and immunohistochemical evaluation of wound healing potential of Nigerian bee propolis in albino rats,” *Comparative Clinical Pathology*, vol. 28, no. 2, pp. 455–466, 2019.
- [54] M. Nani, A. Leone, V. P. Bom et al., “Evaluation and comparison of wound healing properties of an ointment (alpaWash) containing Brazilian micronized propolis and peucedanum ostruthium leaf extract in skin ulcer in rats,” *International Journal of Pharmaceutical Compounding*, vol. 22, no. 2, pp. 154–163, 2018.
- [55] N. Gheib, A. Farzand, Z. Habibi, and F. Samiee-Rad, “The effect of oral consumption of propolis alone and in combination with silver nanoparticles on wound healing in male wistar rats,” *Wound Management & Prevention*, vol. 66, no. 4, pp. 38–46, 2020.
- [56] D. S. Ernawati and A. Puspa, “Expression of vascular endothelial growth factor and matrix metalloproteinase-9 in *Apis mellifera* Lawang propolis extract gel-treated traumatic ulcers in diabetic rats,” *Veterinary World*, vol. 11, no. 3, pp. 304–309, 2018.
- [57] A. Kiderman, R. Torten, A. L. Furst, and K. Reinus, “Bilateral eosinophilic ulcers in an infant treated with propolis,” *Journal of Dermatological Treatment*, vol. 12, no. 1, pp. 29–31, 2001.
- [58] N. Samet, C. Laurent, S. M. Susarla, and N. Samet-Rubinsteen, “The effect of bee propolis on recurrent aphthous stomatitis: a pilot study,” *Clinical Oral Investigations*, vol. 11, no. 2, pp. 143–147, 2007.
- [59] V. R. Santos, R. T. Gomes, R. A. D. Mesquita et al., “Efficacy of Brazilian propolis gel for the management of denture stomatitis: a pilot study,” *Phytotherapy Research*, vol. 22, no. 11, pp. 1544–1547, 2008.
- [60] M. Kucharzewski, M. Kozka, and T. Urbanek, “Topical treatment of nonhealing venous leg ulcer with propolis ointment,” *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 254017, 5 pages, 2013.
- [61] S. M. Bergin and P. Wraight, “Silver based wound dressings and topical agents for treating diabetic foot ulcers,” *Cochrane Database of Systematic Reviews*, vol. 2006, no. 1, Article ID CD005082, 2006.
- [62] F. R. Henshaw, T. Bolton, V. Nube et al., “Topical application of the bee hive protectant propolis is well tolerated and improves human diabetic foot ulcer healing in a prospective feasibility study,” *Journal of Diabetes and Its Complications*, vol. 28, no. 6, pp. 850–857, 2014.
- [63] W. N. Hozzein, G. Badr, A. A. Al Ghamdi, A. Sayed, N. S. Al-Waili, and O. Garraud, “Topical application of propolis enhances cutaneous wound healing by promoting TGF-beta/Smad-mediated collagen production in a streptozotocin-induced type I diabetic mouse model,” *Cellular Physiology and Biochemistry*, vol. 37, no. 3, pp. 940–954, 2015.
- [64] V. Mujica, R. Orrego, R. Fuentealba, E. Leiva, and J. Zuniga-Hernandez, “Propolis as an adjuvant in the healing of human diabetic foot wounds receiving care in the diagnostic and treatment centre from the regional hospital of Talca,” *Journal of Diabetes Research*, vol. 2019, Article ID 2507578, 10 pages, 2019.
- [65] D. Z. Peng, “The pathogenetic factors, molecular mechanism and the management strategies of postburn inflammatory reaction,” *Chinese Journal of Burns*, vol. 21, no. 6, pp. 405–409, 2005.
- [66] N. S. Guimarães, J. C. Mello, J. S. Paiva et al., “Baccharis dracunculifolia, the main source of green propolis, exhibits potent antioxidant activity and prevents oxidative mitochondrial damage,” *Food and Chemical Toxicology*, vol. 50, no. 3–4, pp. 1091–1097, 2012.
- [67] F. Öztürk, E. Kurt, M. Cerçi et al., “The effect of propolis extract in experimental chemical corneal injury,” *Ophthalmic Research*, vol. 32, no. 1, pp. 13–18, 2000.
- [68] S. V. McLennan, J. Bonner, S. Milne et al., “The anti-inflammatory agent propolis improves wound healing in a rodent model of experimental diabetes,” *Wound Repair and Regeneration*, vol. 16, no. 5, pp. 706–713, 2008.

- [69] S. R. Gregory, N. Piccolo, M. T. Piccolo, M. S. Piccolo, and J. P. Heggors, "Comparison of propolis skin cream to silver sulfadiazine: a naturopathic alternative to antibiotics in treatment of minor burns," *Journal of Alternative & Complementary Medicine*, vol. 8, no. 1, pp. 77–83, 2002.
- [70] A. Picolotto, D. Pergher, G. P. Pereira et al., "Bacterial cellulose membrane associated with red propolis as phyto-modulator: improved healing effects in experimental models of diabetes mellitus," *Biomedicine & Pharmacotherapy*, vol. 112, Article ID 108640, 2019.
- [71] G. S. Ashcroft, X. Yang, A. B. Glick et al., "Mice lacking smad3 show accelerated wound healing and an impaired local inflammatory response," *Nature Cell Biology*, vol. 1, no. 5, pp. 260–266, 1999.
- [72] O. K. Mirzoeva and P. C. Calder, "The effect of propolis and its components on eicosanoid production during the inflammatory response," *Prostaglandins, Leukotrienes and Essential Fatty Acids*, vol. 55, no. 6, pp. 441–449, 1996.
- [73] Y. S. Song, E. H. Park, G. M. Hur, Y. S. Ryu, Y. M. Kim, and C. Jin, "Ethanol extract of propolis inhibits nitric oxide synthase gene expression and enzyme activity," *Journal of Ethnopharmacology*, vol. 80, no. 2-3, pp. 155–161, 2002.
- [74] K. Tan-No, T. Nakajima, T. Shoji et al., "Anti-inflammatory effect of propolis through inhibition of nitric oxide production on carrageenin-induced mouse paw edema," *Biological and Pharmaceutical Bulletin*, vol. 29, no. 1, pp. 96–99, 2006.
- [75] X. P. Cao, Y. F. Chen, J. L. Zhang, M. M. You, K. Wang, and F. L. Hu, "Mechanisms underlying the wound healing potential of propolis based on its in vitro antioxidant activity," *Phytomedicine*, vol. 34, pp. 76–84, 2017.
- [76] J. J. Veloz, M. Alvear, and L. A. Salazar, "Antimicrobial and antibiofilm activity against streptococcus mutants of individual and mixtures of the main polyphenolic compounds found in Chilean propolis," *BioMed Research International*, vol. 2019, Article ID 7602343, 7 pages, 2019.
- [77] N. R. J. Lupatini, P. Danopoulos, R. Swikidisa, and P. V. Alves, "Evaluation of the antibacterial activity of green propolis extract and meadowsweet extract against *Staphylococcus aureus* bacteria: importance in wound care compounding preparations," *International Journal of Pharmaceutical Compounding*, vol. 20, no. 4, pp. 333–337, 2016.
- [78] D. Khodabakhshi, A. Eskandarinia, A. Kefayat et al., "In vitro and in vivo performance of a propolis-coated polyurethane wound dressing with high porosity and antibacterial efficacy," *Colloids and Surfaces B: Biointerfaces*, vol. 178, pp. 177–184, 2019.
- [79] D. C. Zancanela, C. S. Funari, R. D. Herculano et al., "Natural rubber latex membranes incorporated with three different types of propolis: physical-chemistry and antimicrobial behaviours," *Materials Science and Engineering: C*, vol. 97, pp. 576–582, 2019.
- [80] E. A. Tosi, E. Ré, M. E. Ortega, and A. F. Cazzoli, "Food preservative based on propolis: bacteriostatic activity of propolis polyphenols and flavonoids upon *Escherichia coli*," *Food Chemistry*, vol. 104, no. 3, pp. 1025–1029, 2007.
- [81] F. B. Guo, "The antibacterial activity of propolis," *Journal of Biochemistry*, vol. 36, no. 3, pp. 10–12, 2004.
- [82] D. M. Qi, "The pharmacological action and clinical application of propolis," *Medical Times*, no. 4, pp. 44–45, 2008.
- [83] A. Parihar, M. S. Parihar, S. Milner, and S. Bhat, "Oxidative stress and anti-oxidative mobilization in burn injury," *Burns, gastric healing activity promoted by Brazilian green propolis and the healing efficacy of artemisinin C*, *Inflammopharmacology*, vol. 28, no. 4, pp. 1009–1025, 2020.
- [84] A. Ocakci, M. Kanter, M. Cabuk, and S. Buyukbas, "Role of caffeic acid phenethyl ester, an active component of propolis, against NaOH-induced esophageal burns in rats," *International Journal of Pediatric Otorhinolaryngology*, vol. 70, no. 10, pp. 1731–1739, 2006.
- [85] Y. Y. Wang, Y. Y. Huang, and D. U. Juan, "Effects of honey and propolis on wound healing in rats with second-degree scald," *Journal of Tianjin University of Traditional Chinese Medicine*, vol. 31, no. 03, pp. 154–156, 2006.
- [86] J. L. Zhang, K. Wang, and F. L. Hu, "Advance in studies on antioxidant activity of propolis and its molecular mechanism," *China Journal of Chinese Materia Medica*, vol. 38, no. 16, pp. 2645–2652, 2013.
- [87] P. Olczyk, P. Ramos, K. Komosinska-Vassev, J. Stojko, and B. Pilawa, "Positive effect of propolis on free radicals in burnwounds," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 356737, 12 pages, 2013.
- [88] A. A. Aytekin, S. Tuncay Tanriverdi, F. Aydın Köse, D. Kart, İ Eroğlu, and Ö Özer, "Propolis loaded liposomes: evaluation of antimicrobial and antioxidant activities," *Journal of Liposome Research*, vol. 30, no. 2, pp. 107–116, 2020.
- [89] S. Touzani, H. Imtara, S. Katekhaye et al., "Determination of phenolic compounds in various propolis samples collected from an African and an Asian region and their impact on antioxidant and antibacterial activities," *Molecules*, vol. 26, no. 15, p. 4589, 2021.
- [90] M. Y. Boufadi, J. Soubhye, and P. Van Antwerpen, "Anti-inflammatory, antioxidant effects, and bioaccessibility of Tizirt propolis," *Journal of Food Biochemistry*, vol. 45, no. 4, Article ID e13663, 2021.
- [91] S. Martinotti, G. Pellavio, U. Laforenza, and E. Ranzato, "Propolis induces AQP3 expression: a possible way of action in wound healing," *Molecules*, vol. 24, no. 8, p. 1544, 2019.
- [92] J. M. Sforcin, "Propolis and the immune system: a review," *Journal of Ethnopharmacology*, vol. 113, no. 1, pp. 1–14, 2007.
- [93] V. Dimov, N. Ivanovska, V. Bankova, and S. Popov, "Immunomodulatory action of propolis: IV prophylactic activity against gram-negative infections and adjuvant effect of the water-soluble derivative," *Vaccine*, vol. 10, no. 12, pp. 817–823, 1992.

- [94] T. Tatefuji, N. Izumi, T. Ohta, S. Arai, M. Ikeda, and M. Kurimoto, "Isolation and identification of compounds from Brazilian propolis which enhance macrophage spreading and mobility," *Biological & Pharmaceutical Bulletin*, vol. 19, no. 7, pp. 966–970, 1996.
- [95] S. Ansorge, D. Reinhold, and U. Lendeckel, "Propolis and some of its constituents down-regulate DNA synthesis and inflammatory cytokine production but induce TGF- β 1 production of human immune cells," *Zeitschrift Für Naturforschung C*, vol. 58, no. 7-8, pp. 580–589, 2003.
- [96] C. L. Baum and C. J. Arpey, "Normal cutaneous wound healing: clinical correlation with cellular and molecular events," *Dermatologic Surgery*, vol. 31, no. 6, pp. 674–686, 2006.
- [97] Y. Hiromatsu and S. Toda, "Mast cells and angiogenesis," *Microscopy Research and Technique*, vol. 60, no. 1, pp. 64–69, 2003.
- [98] L. J. Walsh, "Mast cells and oral inflammation," *Critical Reviews in Oral Biology & Medicine*, vol. 14, no. 3, pp. 188–198, 2003.
- [99] T. A. Wilgus and B. C. Wulff, "The importance of mast cells in dermal scarring," *Advances in Wound Care*, vol. 3, no. 4, pp. 356–365, 2014.
- [100] K. Nishida, A. Hasegawa, S. Yamasaki et al., "Mast cells play role in wound healing through the ZnT2/GPR39/IL-6 axis," *Scientific Reports*, vol. 9, no. 1, Article ID 10842, 2019.
- [101] D. E. A. Komi, K. Khomtchouk, and P. L. Santa Maria, "A review of the contribution of mast cells in wound healing: involved molecular and cellular mechanisms," *Clinical Reviews in Allergy and Immunology*, vol. 58, no. 3, pp. 298–312, 2020.
- [102] M. S. Cho, W. S. Park, W. K. Jung et al., "Caffeic acid phenethyl ester promotes anti-inflammatory effects by inhibiting MAPK and NF- κ B signaling in activated HMC-1 human mast cells," *Pharmaceutical Biology*, vol. 52, no. 7, pp. 926–932, 2014.
- [103] M. A. Nader, "Caffeic acid phenethyl ester attenuates IgE-induced immediate allergic reaction," *Inflammopharmacology*, vol. 21, no. 2, pp. 169–176, 2013.
- [104] T. Kanda, H. Akiyama, A. Yanagida et al., "Inhibitory effects of apple polyphenol on induced histamine release from RBL-2H3 cells and rat mast cells," *Bioscience, Biotechnology, and Biochemistry*, vol. 62, no. 7, pp. 1284–1289, 1998.
- [105] M. Kawai, T. Hirano, S. Higa et al., "Flavonoids and related compounds as anti-allergic substances," *Allergology International*, vol. 56, no. 2, pp. 113–123, 2007.
- [106] S. Yano, D. Umeda, T. Yamashita et al., "Dietary flavones suppresses IgE and Th2 cytokines in OVA-immunized BALB/c mice," *European Journal of Nutrition*, vol. 46, no. 5, pp. 257–263, 2007.
- [107] R. Nakamura, R. Nakamura, K. Watanabe et al., "Effects of propolis from different areas on mast cell degranulation and identification of the effective components in propolis," *International Immunopharmacology*, vol. 10, no. 9, pp. 1107–1112, 2010.
- [108] Y. Bae, S. Lee, and S. H. Kim, "Chrysin suppresses mast cell-mediated allergic inflammation: involvement of calcium, caspase-1 and nuclear factor- κ B," *Toxicology and Applied Pharmacology*, vol. 254, no. 1, pp. 56–64, 2011.
- [109] P. R. Barroso, R. Lopes-Rocha, E. M. F. Pereira et al., "Effect of propolis on mast cells in wound healing," *Inflammopharmacology*, vol. 20, no. 5, pp. 289–294, 2012.
- [110] B. Pippi, A. J. Lana, R. C. Moraes et al., "In vitro evaluation of the acquisition of resistance, antifungal activity and synergism of Brazilian red propolis with antifungal drugs on *Candida spp.*," *Journal of Applied Microbiology*, vol. 118, no. 4, pp. 839–850, 2015.
- [111] S. Stepanović, N. Antić, I. Dakić, and M. Svabić-Vlahović, "In vitro antimicrobial activity of propolis and synergism between propolis and antimicrobial drugs," *Microbiological Research*, vol. 158, no. 4, pp. 353–357, 2003.
- [112] W. Krol, S. Scheller, J. Shani, G. Pietsch, and Z. Czuba, "Synergistic effect of ethanolic extract of propolis and antibiotics on the growth of staphylococcus aureus," *Arznei-mittelforschung*, vol. 43, no. 5, pp. 607–609, 1993.
- [113] J. C. Ahn, R. Biswas, and P. S. Chung, "Synergistic effect of radachlorin mediated photodynamic therapy on propolis induced apoptosis in AMC-HN-4 cell lines via caspase dependent pathway," *Photodiagnosis and Photodynamic Therapy*, vol. 10, no. 3, pp. 236–243, 2013.