



EFFECT OF FUNGICIDAL SEED TREATMENT ON SEED MYCOFLORA AND SEED GERMINATION DURING STORAGE OF SORGHUM

Dr. P.V. Pawar

Madhavrao Patil ACS College,
Palam Dist. Parbhani

ABSTRACT

The Sorghum Cv. Nirmal - 21 seed lots were treated with fungicides Thiram, Carbendazim, Mancozeb, Copper oxychloride, Captan, Captafol @2.5g/kg seed and untreated lot was kept as control. These seed lots were tested for seed mycoflora and seed germination for 36 months at 2 months interval. There was significant increase in seed mycoflora and reduction in seed germination after 36 months storage period. The germination was significantly superior in Thiram treated (83%) than Carbendazim (80%), Mancozeb, (76%) Copper oxychloride (75%), Captan (74%), Captafol (59%) and untreated control (54%). y words-Sorghum vulgare (Pers.) Cv. Nirmal-21), Fungicides

KEYWORDS: Seeds, Fungi, Plant, Mycoflora

INTRODUCTION

Sorghum seeds, treated with fungicides, are reported to harbour reduce seed mycoflora and maintain seed germination for longer period than untreated ones. The reports of Vyas and Nene (1975); Nagrajan and Karivaratharaju (1976); Vidhyasekaran et al. (1980); Mughate and Raut (1982) and Patil and Jarhad (1988) confine to the storability of either hybrids or varieties of Sorghum. This study was undertaken to assess the longevity of fungicide treated seeds of Sorghum Cv Nirmal-21 during storage under ambient condition

MATERIALS AND METHODS

The experiment was conducted during 2007-08, 08-09 and 09-10. The seeds obtained after harvest in October 2010 were adjusted to 9-10% moisture by drying in sunlight. The seeds were treated with fungicides thiram, Carbendazim, Mancozeb, Copper oxychloride, Captan and Captafol seed @ 2.5g/kg seeds and untreated seed lot was considered as control. Both treated and untreated seeds were stored in cloth bags. The seeds were tested at two month intervals for 36 months for germination and seed mycoflora by blotter paper method. The main objectives of seed health testing are related to the actual policy towards seed improvement, seed trade and plant protection (De Tempe 1953, ISTA 1966 and Anonymous 1976).

RESULTS AND DISCUSSION

The results (Table 1) indicated that there was a significant increase in mycoflora from 5 to 31% and reduction in seed germination from 89 to 57% after 36 months storage in control. The minimum germination

Table : 2 Seed mycoflora and seed germination as influenced by Fungicides in sorghum Cv. Nirmal – 21

Fungicides	Seed Mycoflora (%)	Seed germination (%)
Thiram	07	83
Carbendazim	08	80
Mancozeb	08	76
Copper oxychloride	10	75
Captan	14	74
Captafol	17	59
Control	29	54
S.E.	2.7	3.8
C.D. at 5%	6.7	9.3

for maintaining standards (75%) was maintained up to a period of 14 months after storage under normal conditions at udgir.

The germination percentage was significantly in thiram treated seed than superior carbandazim, mancozeb, copper oxychloride, captan captafol and untreated control. Seed mycoflora was also less in the thiram treated seed (Table 2) .Similar beneficial effects of seed treatment were obtained in sorghum by Vyas and Nane (1975) Nagrajan and Karivaratharaju (1976), Muggbate and Raut (1982) and patil and Jarhad (1988).

It was thus evident from this study the sorghum Cv. Nirmal-21 could maintain its germination above certification standard (75%) for about 14 months from harvest at udgir station. There is no need of retreatment or revalidation in up to 14 months in storage for sorghum Cv. Nirmal-21.

ACKNOWLEDGEMENT

I do appreciate the encouragement given and wish to thank Principal Dr. H. R. Aglave Lokmanya Mahavidyalaya Sonkhed and Principal, Dr. D. R. More N. W. College Akhada Balapur.

Table 1 : Seed mycoflora and seed germination influenced by period in sorghum Cv. Nirmal - 21

Period (Months)	Seed Mycoflora (%)	Seed germination (%)
0	5	89
2	6	84
4	7	82
6	7	80
8	8	79
10	9	77
12	9	75
14	11	75
S.E.+	1.49	1.5
C.D. at 5%	3.5	04
16	13	73
18	14	72
20	16	72
22	16	69
24	17	68
26	18	67
S.E.+	0.6	22
C.D. at 5%	1.5	56
28	12	64
30	27	62
32	28	60
34	30	58
36	31	57
S.E.+	1.2	1.14
C.D. at 5%	3.5	3.16

References :-

- Anonymous. 1976. International rules for seed testing. Proc. Int. Seed. Test. Assoc. 32: 562-598.
- De Temp J. 1953. The blotter method of seed health testing programme. ISTA 28: 133-151.
- Garud, TB Syed Ismail and Shinde BM. 2000. Effect of two mould-causing fungi on seed germination of sorghum seed. International Sorghum News Letter 41:54.
- ISTA. 1966. International rules for seed testing. Proc. Int. Seed. Test. Assoc. 32:565-589.
- Jadhav AS. 2009. Seed health management of Jawar. Ph.D thesis S.R.T.M.U; Nanded.
- Mughate AG and Raut JG. 1982. Efficacy of nine different fungicides against fungi frequently associated with sorghum seed. Pesticides. 16: 16-18.
- Nagrajan K and Karivaratharaju TV. 1976. Storage studies in sorghum, Bajraand Maize seed viability in relation to moisture content. Seed Res. 4: 167-173.
- Patil RB and Jarhad AL. 1988. Storability of seeds of sorghum inbreds under ambient conditions. Seed Res. 16:88091
- Vidhyasekaran P Thulasidas G Ramassamy KR and Kandaswamy TK. 1980. Preservation of viability of sorghum seeds by controlling seed borne fungi; Indian phytopath. 33: 225-230.
- Pardeshi VF Reddy VG and Nalwandikar PK. 1989. Effect of different fungicides on seedling vigour and seed viability in Soyabean. J. Maharashtra, Agric. Uni. 14 (1)10-15.
- Vyas SC and Nene YL. 1975 Influence of storing thiram treated sorghum on germination JNKVV. Res J. 9: 114-119.

