

A RETROSPECTIVE STUDY ON EPIDEMIOLOGY OF ASTHMA AND CHRONIC OBSTRUCTIVE PULMONARY DISEASE

Akhila Janga, Bhavana Pamulapati, Divya Bharathi Jekkula, Dr. M.V. Sree Keerthi*, Rajeev Gollapalli*

1. Akhila Janga - Pharm.D, Department of Pharmacy Practice, St. Peter's Institute of Pharmaceutical Sciences, Hanamkonda, Warangal - 506001, Telangana, India.
2. Bhavana Pamulapati - Pharm.D, Department of Pharmacy Practice, St. Peter's Institute of Pharmaceutical Sciences, Hanamkonda, Warangal - 506001, Telangana, India.
3. Divya Bharathi. Jekkula - Pharm.D, Department of Pharmacy Practice, St. Peter's Institute of Pharmaceutical Sciences, Hanamkonda, Warangal - 506001, Telangana, India.
4. Dr. Sree Keerthi MV – MBBS, Chest Physician, DTRD, Vennela Chest Clinic, Jagityal District, Telangana , India,
5. Rajeev Gollapalli – Associate Professor, St. Peter's Institute of Pharmaceutical Sciences, Hanamkonda, Warangal - 506001 , Telangana, India.

Abstract:

Chronic respiratory disease which majorly includes Asthma and Chronic Obstructive pulmonary disease. There were only few studies from India on the field of epidemiology of Asthma and COPD. The prevalence rates for both disorders reported in different studies have widely varied. In phase-I of the observational study on epidemiology of Asthma and COPD was conducted at a chest clinic. Population prevalence of Asthma and COPD in all age groups was reported and analyzed. In phase-II of observational study a standardized and validated questionnaire was used employing a uniform methodology in all patients. The main objective of this project is to find differences in the epidemiology of asthma and chronic obstructive pulmonary disease, including in the prevalence rates and other casual variables (age, gender, residence, occupation), if any, in different places in a Chest Clinic. And also to assess the influence of exposure to tobacco smoking, ETS (Environmental tobacco smoke) and other risk factors, on the prevalence of both Asthma and COPD. The results were reported and analyzed. Based on the results, it was found that, the prevalence rate of Asthma was found to be 0.35% and COPD of 0.27% (approximately).

Keywords: Asthma, Chronic Obstructive pulmonary disease, Risk factors, Epidemiology

INTRODUCTION

India is a country with vast economical, racial & religious differences. Chronic respiratory disease includes majorly Asthma and COPD. These diseases account for 100 million individuals in India. Both are the inflammatory disorders of airways.

ASTHMA:

Asthma is a chronic inflammatory disorder characterized by airway hyper-responsiveness to a variety of stimuli. It results from a complex interaction among inflammatory cells, mediators and airways. For practical purposes, asthma may be defined as a disorder of the airways characterized by the following:

- 1) Paroxysmal and/or persistent symptoms such as dyspnea, chest tightness, wheezing and cough, with or without mucus production.
- 2) Variable airflow limitation demonstrated by chest auscultation and/or repeated measurements of peak expiratory flow (PEF) or other spirometric indices.
- 3) Airway hyper-responsiveness to a variety of specific and nonspecific inhalational stimuli. Severity of obstruction is variable and believed to be as a result of bronchoconstriction, airway wall edema, mucus plug formation. It reduces the

ventilation to some lung regions which causes a perfusion imbalance that leads to hypoxemia. According to 2001 census, in India prevalence of Asthma is 2.03% -men,1.87%-women total 1.95% [1]

COPD:

Chronic obstructive pulmonary disease is also an inflammatory disorder of the airways which is a major cause of disability and death all over the world, including in India. Unlike asthma, COPD is primarily a disease of adults and runs a progressive downhill course. It includes the clinical conditions of chronic bronchitis and emphysema where chronic bronchitis is defined as the presence of cough and mucoid expectoration for at least 3months in a year continuously for two or more years and Emphysema is a clinical condition causing permanent damage to the alveoli. In this respiratory tissue inflammation occurs which results in vasodilation, congestion, mucosal edema, goblet cell hypertrophy. These events trigger goblet cells to produce excess amount of mucous. In severe condition, clusters of alveoli merge further leading to diminishing the number of alveoli finally resulting in increased space available for air trapping. According to 2001 census in India, prevalence of COPD observed as men-4.18%, women-2.74% and total-3.48% [1].

NEED OF THE STUDY:


Due to no particular differentiation of Asthma and COPD symptoms, the prevalence rates greatly vary in different conditions and different studies.

This variation may also due to place of residence, socio-economic status. Smoking and tobacco chewing also shows significant difference. Thus epidemiological studies are required to assess the respiratory burden of particular area with respective to the exposure history and risk factors. This study was carried out in a district named Jagityal, with a geographical area of 3,043.23 km² and a population of 1,03,930 (2001 census).It is located in Telangana state, India. The population of this district consists predominantly of backward classes and people belonging to the lower rung of the social ladder. Agriculture and beedi workers are the two main occupations of this region which are found to be main reasons for the progression of diseases asthma and COPD.

PREVIOUS STUDIES:

There were only few studies from India on the field of epidemiology of Asthma and COPD. The prevalence rates for both disorders reported in different studies have widely varied. According to 2001 census, in India prevalence of Asthma is 2.03% - men, 1.87% - women, total 1.95%, whereas for COPD men-4.18%,women-2.74%,total-3.48%.

In a multicenter study during 2006-2009, "Indian study on epidemiology of Asthma, Respiratory symptoms and Chronic bronchitis (INSEARCH)", the prevalence rates of Asthma and COPD in various regions were as follows



AIMS AND OBJECTIVES OF THE STUDY

Aims:

To study the epidemiology of both the diseases that includes Asthma and Chronic Obstructive Pulmonary Disease.

Objectives:

- 1) To find out the differences in the epidemiological aspects of asthma and chronic obstructive pulmonary disease including the prevalence rates and other casual variables (age, gender, residence, occupation) in different places of study site .
- 2) To assess the influence of exposure to tobacco smoking, ETS (Environmental tobacco smoke) and other risk factors on the prevalence of both Asthma and COPD.

METHODOLOGY

STUDY SITE: The proposed study was carried out at Vennela Chest Clinic, Jagityal. During the study period, a total number of 175 patients were enrolled into the study. Out of them, 60 asthma patients and 50 COPD patients were considered.

Vennela Hospital is a secondary health care hospital setup in Jagityal dist. This was the private center with the aim to provide effective service to the people of Jagityal and surrounding areas. Pulmonology is the major department. It provides all the facilities along with the adequate laboratory needs (it is equipped with facilities for spirometry, bronchoscope, Respiratory critical care unit and emergency requirements)

STUDY DESIGN:

A Retrospective Observational Study was done during May 2016

A Retrospective Observational study looks backwards and examines exposures to suspected risk or protection factors in relation to an outcome that is established at the start of the study, to determine how these factors affects the rates of a certain outcome.

STUDY CRITERIA:

Inclusion criteria:

- Patients who are diagnosed with asthma and chronic obstructive pulmonary disease.
- Symptoms and family history showing evidence of both the diseases.
- Patients of either sex.
- All age groups (mostly from 0yrs-85yrs).

Exclusion criteria:

- Patients with serious life threatening conditions
- Patients with Tuberculosis
- Patients with uncontrolled infections.

Study materials:

All the relevant and necessary data was collected in the data collection form which includes Interviewing patients or patient care takers, health care professionals, Prescriptions, Patient case sheet and any other relevant sources.

The data collected include: Demographic details, Chief complaints, Past medical history, Past medication history, Family history, Social history, General examination, information about sensitivity towards occupational and environmental triggering factors, social habits like smoking and tobacco usage, diagnosis and Medication chart.

Study procedure:

Patients visiting Vennela chest clinic were reviewed on a daily basis. The patients who met the study criteria were enrolled into the study and the following information was collected: patient demographics, chief complaints, PMHx, FHx, SHx, sensitivity towards occupational and environmental factors, smoking and tobacco usage and current medication. The patients with positive diagnosis towards asthma and chronic obstructive pulmonary disease were included. The data collected manually was reviewed and entered into Microsoft Excel database and then analyzed statistically.

RESULTS

The roles of several factors were assessed in the study to help clinicians in decision making.

1. **Gender:** The percentage of allergy is greater in women than in men.

Table 1: Gender wise distribution

Gender	Total	asthma	COPD
Male	42%	31%	54%
Female	58%	68.33%	46%

In this study, the significant occurrence of Asthma and COPD was strongly associated with female population with (P=0.58) and the percentage of female population are higher when compared to males with frequencies of 64(58%) and 46(42%) respectively. The ratio of females- males (F/M) is 1.38:1. The p values of males was found to be P=0.316 in and P=0.68 in females in Asthma, whereas in COPD P=0.540 in males and P= 0.460 in females

2. **Age:** Asthma is more among the individuals of young age 10-25years, whereas COPD was more between the age group of 60-70years.

ASTHMA

AGE (IN YEARS)	PERCENTAGE
35-40	2%
40-50	20%
51-60	24%
61-70	38%
71-80	12%
81 and above	4%

COPD

AGE IN YEARS	PERCENTAGE
0-25	42%
26-40	32%
41-60	22%
61 and above	4%

In this Study population, the percentage of asthma is higher in the age group of 0-25years (40%), followed by 26-40years (33.3%), 41-60 years (21.66%), 61 and above (5%), and among COPD higher number of patients were in age group of 61-70 years (38%), 51- 60years(24%), then 40 to 50 years (20%),71-80 years (12%) 81 and above (4%),35-40years(2%) age groups patients.

3. Other risk factors (smoking and tobacco chewing)

	ASTHMA			COPD		
	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
Smoking history	5%	5%	-	38%	38%	-
Tobacco chewing	11.66%	3.33%	8.33%	44%	10%	34%

Significant occurrence of Asthma and COPD was found in relation with the risk factor tobacco chewing, predominantly in females.

Male patients (5%) diagnosed with asthma who having habit of smoking, were as male patients (38%) with COPD were smokers. In tobacco chewing in asthma (n=7) patients(11.66%) were observed, among them (n=2) male patients 3.33%) female patients (n=5) 8.33% and in COPD(n= 22) patients 44% were observed, among them male patients (n=5) 10% and female patients (n=17) 34% were observed. The p value of COPD for smoking was found to be 0.380 in males and p=0.440 in females in case of tobacco chewing and it is the predominant.

4. Family History:

The family history of the Asthma patients was taken during the patient interview.

Table 4: Assessment of Family History

Total	25%
Male	10%
Female	15%

In this study population, family history of asthma was found to be significant with current asthmatic patients and it was found to be 25% (n=15 & P=0.240). There is no evidence of genetic predisposition in COPD patients.

5. Occupation:

Table 5: Assessment of Occupation

OCCUPATION	ASTHMA	COPD
Beedi worker	20%	8%
Farmer	21.66%	68%
Unskilled worker	16.66%	22%
House wife	10%	2%
Student	26.66%	-
Others	5%	-

In both the diseases asthma and COPD, farmers occupy a major number of total populations followed by beedi workers. More number of students also observed in case of asthma.

DISCUSSION

Epidemiology is the study of the occurrence, distribution and determinants of health conditions in populations, and the application of this knowledge to control health problems. It is fundamental to improving public health. Chronic respiratory disease which includes Asthma and Chronic Obstructive pulmonary disease, may account for an estimated burden India. There were only few studies from India on the field of epidemiology of Asthma and COPD. The prevalence rates for both disorders reported in different studies have widely varied. According to 2001 census, in India prevalence of Asthma is 2.03% -men, 1.87% - women, total 1.95%, whereas for COPD men-4.18%,women-2.74%,total-3.48%.Based on the study ,it was found that, the prevalence rate of Asthma was found to be0.35% and COPD of 0.27%(approximately).

In this study, the percentage of female population was higher when compared to males. The percentage of asthma is higher in the age group of 10-25years which indicates the genetic predisposition, and among COPD higher number of patients were in age group of 61-70 years, as COPD is a chronic disease and is progressed as age increases. Among Asthma patients75% were allergic to occupational and environmental triggers, were as in COPD 74% were allergic as most of the population are beedi workers and farmers which are the main risk factors. Among COPD 38% were smokers.

The tobacco chewing was found to be 44% in COPD. Family history of asthma was found to be more, and it is observed to have 25% of total asthma patients. There is no evidence of genetic predisposition in COPD patients. In the study population in both the diseases asthma and COPD farmers occupy a major number of total populations followed by beedi workers. More number of students is also observed in case of asthma.

LIMITATIONS OF THE STUDY:

1. Short duration of study period and less number of samples may not give the exact prevalence in this particular geographical area.
2. Seasonal variation in the prevalence could not be determined.

CONCLUSION:

The strong environmental and occupational triggering factor that contributes for the development of COPD was observed with air pollution.

Genetic predisposition plays a major risk factor causing Asthma in addition to air pollution. Apart from these risk factors, occupation contributes to the progression of disease. Its alarming time to bring out significant monitoring in the respiratory disease prone areas considering their risk factors and exposure history. Different patient counselling strategies should be adopted in such areas and awareness among the population considering the risk factors and triggers that could have a strong significant relation with Asthma and COPD.

This plays a vital role in decreasing the prevalence and incidence rates. Further understanding of risk factors in relation to pathophysiology could improve management in this vulnerable population.

ACKNOWLEDGMENT:

I would like to pay special thankfulness, warmth and appreciation to persons below who made my research successful and assisted me at every point to cherish my goal

My expert advisor, Dr. Sree Keerthi whose help and sympathetic attitude at every point during my research helped me to work in time

My guide, Mr. Rajeev Gollapalli for his vital support and assistance. His encouragement made it possible to achieve the goal.

Our college principal, Dr. Rajshekar, Chairman Mr. Jaypal Reddy and all the faculty members, my family members and friends who supported throughout my project.

BIBLIOGRAPHY:

- **Epidemiology of chronic obstructive pulmonary disease: a literature review**
Catherine E Rycroft¹, Anne Heyes¹, Lee Lanza² and Karin Becker³
- **Chronic Obstructive Pulmonary Disease Among Adults — United States, 2011**(Morbidity and Mortality Weekly Report (*MMWR*)
Weekly, November 23, 2012 / 61(46);938-943
- **Chronic airway obstruction in a population-based adult asthma cohort: Prevalence, incidence and prognostic factors.**
Backman H¹, Jansson SA¹, Stridsman C², Muellerova H³, Wurst K³, Hedman L⁴, Lindberg A⁵, Rönmark E⁶
- **Prevalence and trend of COPD from 1995-1997 to 2006-2008: The HUNT study, Norway.**
Bhatta L¹, Leivseth L², Mai XM³, Chen Y⁴, Henriksen AH⁵, Langhammer A⁶, Brumpton BM⁷
- **A cross-sectional study on prevalence of chronic obstructive pulmonary disease (COPD) in India: rationale and methods.**
Rajkumar P¹, Pattabi K¹, Vadivoo S¹, Bhome A², Brashier B³, Bhattacharya P⁴, Mehendale SM⁵.
- **Assessing the Prevalence and Incidence of Asthma and Chronic Obstructive Pulmonary Disease in the Eastern Mediterranean Region.**
Masjedi M¹, Ainy E², Zayeri F³, Paydar R⁴.
- **Fall episodes in elderly patients with asthma and COPD - a pilot study.**
Bozek A¹, Jarzab J¹, Hadas E¹, Jakalski M², Canonica GW³.
- **Risk factors for return to the emergency department for asthma: a population-based study.**
To T¹, Zhu J², Ryckman K³, Gershon A⁴.
- **[Study on the current status of smoking, intention of tobacco cessation and related risk factors among 18-65-year olds patients with chronic diseases in Beijing].**
[Article in Chinese; Abstract available in Chinese from the publisher]
Jiang B¹, Ma AJ, Li H, Fang K, Dong J, Xie J, Qi K, Xie C, Zhou Y, Zhao Y, Dong Z.
- **Diagnosis and Management of Asthma - The Swiss Guidelines.**
Rothe T¹, Spagnolo P^{2,3}, Bridevaux PO^{4,5}, Clarenbach C⁶, Eich-Wanger C⁷, Meyer F⁸, Miedinger D^{2,3}, Möller A⁹, Nicod LP¹⁰, Nicolet-Chatelain G¹¹, Sauty A¹², Steurer-Stey C^{13,14}, Leuppi JD^{2,3}.

- **Exacerbations of COPD.**

Viniol C¹, Vogelmeier CF²

- **Diagnosis of asthma-COPD overlap: Is it possible a global definition?**

Miravittles M¹

- **Observational Studies of Inhaled Corticosteroid Effectiveness in COPD: Lessons Learned.**

Suissa S¹, Ernst P².

- **Maternal Smoking during Pregnancy and Early Childhood and Development of Asthma and Rhinoconjunctivitis - a MeDALL Project.**

Thacher JD¹, Gehring U², Gruzieva O¹, Standl M³, Pershagen G^{1,4}, Bauer CP⁵, BerdelD⁶, Keller T⁷, Koletzko S⁸, Koppelman GH^{9,10}, Kull I^{1,11,12}, Lau S¹³, Lehmann I¹⁴, Maier D¹⁵, Schikowski T¹⁶, Wahn U¹³, Wijga AH¹⁷, Heinrich J^{3,18}, Bousquet J^{19,20}, Anto JM^{21,22,23,24}, von Berg A⁶, Melén E^{1,4,11}, Smit HA²⁵, Keil T⁷, Bergström A^{1,4}

- **Rates of asthma exacerbations and mortality and associated factors in Uganda: a 2-year prospective cohort study.**

Kirenga BJ^{1,2}, de Jong C^{3,4}, Mugenyi L^{1,5}, Katagira W², Muhofa A², Kamya MR¹, BoezenHM⁶, van der Molen T^{3,4}

- The overlap syndrome of asthma and COPD: what are its features and how important is it? Professor P Gibson, Department of Respiratory and Sleep Medicine, Level 3 HMRI, John Hunter Hospital, Locked Bag 1, Hunter Mail Centre, Newcastle, NSW Australia 2010; Volume 64, Issue 8.

- . Passive smoking and risk of adult asthma and COPD: an update by David B Coultas, 1998, Volume 53, Issue 5

- Overlap Syndrome of Asthma and COPD Predicts Low Quality of Life, Paula Kauppi, Henna Kupiainen, Ari Lindqvist et al. 2011 Volume 48, Issue 3: 279-285

- Risk factors for new-onset asthma and early COPD in a cohort of young adults. Lisbet Krogh Traulsen, Jesper Baelum, Anders Halling et al. European Respiratory Journal 2015 Volume 46:1118-1122.

- www.Census2011.co.in(INDIA)

ABBREVIATIONS

COPD	Chronic Obstructive Pulmonary Disease
ETS	Environmental Tobacco Smoke
PMHx	Past medical history
FHx	Family history
SHx	Social history