



A DESCRIPTIVE STUDY TO ASSESS THE KNOWLEDGE AND ATTITUDE REGARDING EXCLUSIVE BREAST FEEDING AMONG PRIMIGRAVIDA MOTHERS IN SELECTED HOSPITALS OF KASHMIR WITH A VIEW TO DEVELOP INFORMATION BOOKLET.

Mr. Hilal Ahmad Bhat¹, Ms. Sukhmanpreet Kaur²

¹Mr. Hilal Ahmad Bhat M.sc Nursing Final Year, DBUSON, Mandi Gobindgarh, Punjab.

²Ms. Sukhmanpreet Kaur, Assistant Professor, DBUSON, Mandi Gobindgarh, Punjab.

Abstract

Breastfeeding is the feeding of babies and young children with milk from a woman's breast. The world Health Organization (WHO) recommends that breastfeeding begins within the first hour of a baby's life and continues as often and as much as the baby wants. Mothers may pump milk so that it can be used later when breastfeeding is not possible. For this study Conceptual Frame Work Based on Ludwig Von Bertalanffy in 1968. The approach used for this study was Non experimental descriptive research design. The research variable of the study is Knowledge of Primigravida mothers regarding Exclusive breast feeding and extraneous variable are age, religion, mother education, mother occupation, type of family, family income, trimester, history of minor ailments previous knowledge on breast feeding and source of information etc. The pilot study was conducted to find the feasibility of the study 3rd of March 2021 to 10th of March 2021 among Primigravida mothers. The main study was conducted from 12-05-2021 to 15-05-2021 among 60 subjects; the subject was selected by purposive sampling technique and data was analyzed and interpreted using descriptive and inferential statistics. The overall knowledge on exclusive breast feeding. The mean score is 28.88 with a mean percentage of 72.2% with a standard deviation of 2.558. The overall attitude scores of primi gravid mothers on breast feeding were found to be 57.53% with standard deviation 2.3.

KEYWORDS: Breast feeding, knowledge and attitude Prime Gravida mothers.

Introduction

Breastfeeding is the feeding of babies and young children with milk from a woman's breast. The World Health Organization (WHO) recommends that breastfeeding begins within the first hour of a baby's life and continues as often and as much as the baby wants. During the first few weeks of life babies may nurse roughly every two to three hours, and the duration of a feeding is usually ten to fifteen minutes on each breast. Older children feed less often. Mothers may pump milk so that it can be used later when breastfeeding is not possible.

Breastfeeding is when you feed your baby breast milk, usually directly from your breast. It's also called nursing. Making the decision to breastfeed is a personal matter. It's also one that's likely to draw opinions from friends and family.

Objectives of the study

- ❖ To assess the level of knowledge regarding breast feeding among Primi gravida mothers.
- ❖ To assess the level of attitude regarding breast feeding among Primi Gravida mothers.
- ❖ To find out the association between knowledge and attitude score on breastfeeding with demographic variables.

Methodology

In the present study Non-experimental descriptive research design was used. The present study was conducted at Lal Ded Hospital Srinagar. The population in the present study included 60 Primigravida mothers who are attending OPD at Lal Ded Hospital Srinagar were drawn by Non Probable Convenient sampling technique. The data was collected by a structured demographic sheet and self-structured questionnaire. Data has been collected through forms within 10 days' time period of the month of July 2021.

Findings of the Study: Major Findings are:

Assessment of the level of knowledge regarding breast feeding among Primi gravidmothers.

Sr. No.	Knowledge regarding breast feeding among Primigravida mothers	Knowledge levels					
		Inadequate Below 50%		Moderate 51 – 75%		Adequate Above 75%	
		No	%	No	%	No	%
1	OVER ALL	35	58.3	25	41.7	00	00

Table -1 represents that knowledge level of Primigravida mothers regarding breastfeeding, 35(58.3%) had inadequate knowledge, 25(41.7%) had moderate level of knowledge and none of them adequate knowledge.

Assessment of the level of attitude regarding breast feeding among Primi gravidmothers.

Sr. No.	Attitude regarding breast feeding Among Primi Gravida mothers	Attitude levels					
		Unfavorable attitude Below 50%		Moderate Attitude 51 – 75%		Favorable attitude 75%	
1	Over all	No	%	No	%	No	%
		8	16.0	37	74.0	5	10.0

Table 2 shows that the represents that attitude level of Primigravida mothers regarding breast feeding, 8(16.0%) had Unfavorable attitude, 37(74.0%) had moderate attitude and 5(10%) had favorable attitude.

The Association of knowledge scores of Primigravida mothers with selected Socio-demographic variables

Demographic variables			Level of Knowledge		CHI-SQUARE TEST
			Moderately adequate	Inadequate	
Age	18-20 Years	23	3	20	chi square value=4.149 p=0.126
	21-24 Years	23	6	17	
	25-30 Years	14	6	8	
Religion	Muslim	44	10	34	chi square value= 3.143 p=0.208
	Sikh	1	1	0	
	Hindu	15	4	11	
Mother's education	No formal education	6	3	3	chi square value=4.807 p=0.186
	Primary school	20	2	18	
	Middle school	16	5	11	
	Graduation and above	18	5	13	
Mother's occupation	House wife	24	4	20	chi square value=5.126 p=0.163
	Daily wages	15	7	8	
	Employed	12	2	10	
	Self Employed	9	2	7	
	Self Employed	4	5	9	
Family	Nuclear family	48	13	35	chi square

Type	Joint family	12	2	10	value=0.556 p=0.456
Family Income	Rs. 5001-8000	13	1	12	chi square value=2.679 p=0.262
	Rs. 8001-10000	31	9	22	
	More than 10001	16	5	11	
Trimester	Irregular	26	5	21	chi square value=3.377 p=0.185
	First	12	5	7	
	Second	31	8	23	
History of minor ailments	Third	17	2	15	chi square value=0.247 p=0.619
	Yes	6	2	4	
	No	54	13	41	
Previous information breast feeding	Yes	6	0	6	chi square value=2.222 p=0.136
	No	54	15	39	
Source of information	Health personnel	11	2	9	chi square value=5.910 p=0.206
	Friends	13	5	8	
	Teachers	8	4	4	
	Mass media	7	1	6	
	No information	21	3	18	

The association between socio-demographic variables and the knowledge scores regarding breast feeding. All the selected demographic variables are not significantly associated with the knowledge scores. The association was determined by using Pearson chi square test

The Association of attitude scores of Primigravida mothers with selected demographic variables.

Demographic variables			Level of attitude		CHI-SQUARE TEST
			Below median	Median and above	
Age	18-20 Years	23	12	11	chi square value=0.824 p=0.662
	21-24 Years	23	9	14	
	25-30 Years	14	6	8	
Religion	Hindu	44	24	20	chi square value= 6.226 p=0.044
	Sikh	1	0	1	
	Muslim	15	3	12	
Mother's education	No formal education	6	5	1	chi square value=15.224 p=0.002
	Primary school	20	6	14	
	Middle school	16	3	13	
	Graduation and above	18	13	5	
Mother's occupation	House wife	24	12	12	chi square value=6.914 p=0.075
	Daily wages	15	10	5	
	Employed	12	3	9	
	Self Employed	9	2	7	
	Self Employed	14	5	9	
Family Type	Nuclear family	48	24	24	chi square value=2.424 p=0.119
	Joint family	12	3	9	
Family	Rs. 5001-8000	13	2	11	chi square

The association between socio-demographic variables and the attitude scores regarding breast feeding. Association with religion (chi square 6.226, P=0.044), Mother's education (chi square=15.224, P=0.002), Family Income (Chi Square= 17.507, P=0.000), and source of information (Chi square=10.546, P=0.032) are significantly associated with the attitude scores. The association was determined by using Pearson chi square test.

LIMITATION OF STUDY

- ❖ The findings could be generalized only to the population which fulfilled the criteria in the study.
- ❖ The study limited to assessment of knowledge and attitude.
- ❖ The sample was limited to 60 only.
- ❖ Long-term follow-up could not be carried out due to time constraints.

CONCLUSION

The present study was to assess the knowledge and attitude of Primigravida mothers regarding breast feeding, 35(58.3%) had inadequate knowledge, 25(41.7%) had moderate level of knowledge and none of them adequate knowledge.

REFERENCE

1. Parul Datta, Pediatric Nursing second edition Jaypee brothers, Publication, New Delhi, 2009 Page No.150.
2. Swarna Rekha Bhat Achar's Text book of paediatrics 4th edition, Universities Press Private Limited, India, 2009 Page No.200.
3. Horii N, Allman J, Martin-Prével Y, Waltisperger D. Determinants of early initiation of breastfeeding in rural Niger: cross-sectional study of community-based child healthcare promotion. International breastfeeding journal. 2017 Dec;12(1):41.
4. Tema T. Knowledge, Attitude and Practice Towards Exclusive Breast-Feeding at Jimma, Ethiopia. Ethiopian Journal of Health Sciences. 2000;10(1).
5. Layla E. Arafat (Master student KAU, Head Nurse OB/GYN King Faisal Specialist Hospital & Research Center) Knowledge and Attitude of Exclusive Breast-Feeding Among Saudi Women in Primary Health Care Centers in Jeddah City, Saudi Arabia IOSR.
6. Journal of Nursing and Health Science Volume 6, Issue 6 Ver. VIII. (Nov.- Dec.2017), PP 01-08
7. Deithra Leonard Lowdermilk Maternity and women's health care 9th edition published by mosby elseier Page no 709.
8. Joshi et al. Prevalence of exclusive breastfeeding and associated factors among mothers in rural Bangladesh: a cross-sectional study International Breastfeeding Journal 2014, 9:7.
9. UNICEF Child survival, neonatal mortality current status and progress.
10. WHO Global health observatory data of neonatal mortality, 2018.
11. Ruby L. Wesley, Nursing theories and Model's 2nd edition Page No 84 - 85.
12. Denise F. Polit Essentials of Nursing Research 7th published by Wolters KLUWER Lippincott Williams & Wilkins .
13. Ikonen R, Paavilainen E, Helminen M, Kaunonen M. Preterm infants' mothers' initiation and frequency of breast milk expression and exclusive use of mother's breast milk in neonatal intensive care units. Journal of clinical nursing. 2018 Feb; 27(3 4):e551-8.
14. Horie S, Nomura K, Takenoshita S, Nakagawa J, Kido M, Sugimoto M. A relationship between a level of hemoglobin after delivery and exclusive breastfeeding initiation at a baby friendly hospital in Japan. Environmental health and preventive medicine. 2017 Dec; 22(1):40.