



# A REVIEW ON: SPINAL CORD INFECTIONS

## *WALKING DISORDERS, SPINAL INFECTIONS AND IMAGING TECHNIQUES*

<sup>1</sup>Mr. Sagar Sapkal, <sup>2</sup>Dr. Gajanan Sanap, <sup>3</sup>Mr. Aniket Bankar, <sup>4</sup>Mr. Sudarshan Chavan, <sup>5</sup>Mr. Paresh Patil

<sup>1</sup>Student, <sup>2</sup>Principal, <sup>3</sup>Student, <sup>4</sup>Student, <sup>5</sup>Student

<sup>1</sup>Department of Pharmacy,

<sup>1</sup>Late Bhagirathi Yashwantrao Pathrikar College of Pharmacy, Pathri, Aurangabad, 431001. Maharashtra, India.

**Abstract:** This article consists of infections of the vertebral body, intervertebral disc, ligaments and para-vertebral gentle tissues, epidural space, meninges and subarachnoid space, and the spinal cord. A huge variety of infective organisms can be implicated and the incidence of some, drastically tuberculosis, is growing because of multiplied immune compromised and different factors. Imaging performs a key position in early prognosis of those diseases, which can be extreme and probably life-threatening Infection can also additionally be received by the hematogenous route by infection from contiguous systems or from direct inoculation. Of to be had imaging techniques, CT and MRI provide the clinically most valuable strategies of comparing all the systems probably involved in infection. Although many signs are non-specific, indication is given of in which appearances increase strong suspicion of contamination. The quantity of the inflammatory procedure is well evaluated by imaging, particular lying phrases of severity and morbidity of medical sequelae, which can also additionally be severe. Early research is obligatory and as MRI currently has the key function in research, it must be employed at an early degree of medical suspicion.

**Keywords - Spine Cord Infection, Spondylitis, Spondylodiscitis, Intramedullary infection, Epidural Abscess, Imaging Techniques used in Spinal Cord Infection, Walking Disorders due to Spinal Cord Injury.**

### I. INTRODUCTION

Infections of the axial skeleton can also additionally have an effect on the vertebral body (spondylitis), the intervertebral disk (spondylodiscitis), the ligaments and paravertebral gentle tissues, the epidural space (epidural abscesses), the meninges and subarachnoid space, and finally, very rarely, the spinal wire (myelitis, spinal wire abscesses). Age, the route of contamination, the causative organism, and any underlying disorder impact the morphologic look and the medical route of spinal infection. The maximum common infecting organism is Staphylococcus aureus determined in about 55–80% of cases.<sup>[1, 2]</sup> Other organisms consist of Streptococcus, Pneumococcus, Enterococcus, E.coli (gram positive), Salmonella, Pseudomonas aeruginosa, and Klebsiella (gram negative). Granulomatous infections originate from Mycobacterium tuberculosis, Brucellosis, fungi, and parasites. The upward push of HIV infection, the growth in age of the population, the development of resistance to anti biotherapy, and the rise of immune-compromised sufferers connected to extra competitive therapies have reversed the consistent decline of infections and specially the decline of tuberculous spondylitis. In the U.S., 40,000 new instances of tuberculosis have passed off between 1985 and 1991 as a right away end result of the AIDS epidemic.<sup>[3]</sup> In Europe and North America, tuberculous spondylitis impacts adults with predisposing elements including alcoholism, drug addiction, and immunosuppression.<sup>[1, 4]</sup> Imaging performs a key position in early analysis of those extreme and probably life-threatening lesions. Unfortunately, specificity is frequently missing at the even maximum touchy imaging procedures, including MRI, and the causative agent may be recognized way to laboratory checks or biopsy.




Figure 1, A & B: - L5-S1 Spondylodiscitis.

A) Sagittal & B) Anteroposterior plain film of the lumbo-sacral region: reduction of the height of the L5-S1 disk. Especially to the right arrow, the inferior vertebral plate of L5 is slightly ill.

## II. WALKING DISORDERS DUE TO SPINAL CORD INJURY

A first hanging distinction located in maximum SCI-IMFL topics and in maximum instances of spastic paresis from different neurologic problems is a discounted walking pace, as compared with everyday topics.<sup>[5-9]</sup> However, the walking pace in SCI-IMFL topics stages from “now no longer capable of walk” in very excessive instances to “everyday strolling pace” withinside the least excessive instances. A second critical distinction discovered in SCI-IMFL topics is an extended cycle duration and shorter stride duration. The increase in cycle length is from an increase in each stance and swing intervals. This augmentation of the length parameters ought to be seen, in part, on account of the low on foot velocity of SCI-IMFL topics as opposed to a function of on foot on this population.<sup>[10]</sup> Indeed, in ordinary topics on foot at low velocity, the cycle, stance, and swing intervals were accelerated and the stride duration reduced compared with the values acquired at ordinary comfortable speed.<sup>[11-12]</sup> Some of the characteristics in SCI-IMFL on foot styles as in comparison with ordinary topics on foot on the identical velocity (0.3m/set) are summarized here. Conrad and colleagues<sup>[5-10]</sup> found that subjects with spastic paresis from numerous reasons had shorter cycle length than ordinary topics at 55 m/set. However, opposite consequences have additionally been published.<sup>[10]</sup>

Angular displacements of decrease limb joints withinside the sagittal aircraft in spastic paretic topics display critical modifications in comparison with normal topics.<sup>[5-10]</sup> Hip joint angular tour become more withinside the spastic paretic topics than in the normal subject. All spastic paretic topics made foot touch with the knee in a flexed role, and for a few the knee remained flexed during the stance. At the a joint, the spastic paretic topics tended to have a greater dorsiflexed role at foot touch and the ankle remained greater dorsiflexed than regular even at push-off in overdue stance. During swing, a more variability of the ankle joint role was noted. It may want to continue to be greater dorsiflexed or, in contrast, it may want to continue to be plantarflexed due to a foot drag;<sup>[5-10]</sup> comparable results have additionally been suggested in SCI-IMFL.<sup>[11]</sup> Electromyographic (EMG) recordings of decrease limb muscle mass at some point of on foot screen changes withinside the timing and the amplitude of the interest in SCI-IMFL subjects<sup>3</sup> Coactivation of antagonist muscle mass at proximal and distal joints is frequently suggested in spastic paretic and SCI-IMFL gait.<sup>[13,14]</sup> Abnormal activation of triceps surae muscle in overdue swing or early stance followed with clonus is normally visible, in particular in topics with marked spasticity.<sup>[5,10-14]</sup>

Hyperactive reflexes and hypertonia are frequently associated with joint stiffness encountered in spastic paresis ensuing from SCI.<sup>[25]</sup> The contribution of the stretch reflex and the nonreflex factor to the stiffness of the ankle extensors in regular and spastic topics has been studied.<sup>[16-18]</sup> Spastic topics confirmed a growth of the reflex benefit and a scarcity of inhibition at some point of the swing section compared with regular topics. Further, spastic topics confirmed a small growth of the nonreflex torque at some point of gait. These findings endorse that will increase withinside the reflex benefit and withinside the nonreflex torque may want to each make a contribution to the stiffness of the ankle joint visible in spastic topics at some point of gait. Hence, changes in imperative mechanisms and adjustments in intrinsic homes of the muscle fibers may want to be accountable for the extended stiffness. Other research additionally suggested better benefit of soleus H-reflex in spastic SCI-IMFL topics at some point of on foot compared with regular topics.<sup>[19]</sup> Reduction of presynaptic inhibition of Ia terminals<sup>[20,21]</sup> and faulty reciprocal inhibition<sup>[22]</sup> were cautioned as possible mechanisms contributing to hyperactive reflexes in SCI subjects These hyperactive spinal reflexes had been emphasized in early definitions of spasticity alongside the locating of increased resistance to passive motion.<sup>[23-25]</sup> Recent discussions of spasticity have emphasized the relation among hyperactive reflexes and different motion problems consisting of a number of the additives supplied in parent 1.<sup>[26,27]</sup> Most remedies for spasticity, however, had been designed to deal with handiest the hyperactive spinal reflexes or resistance to passive stretch, even though it has been stated that a few medicinal drugs can also additionally also alternate dystonia.<sup>[28-29]</sup> Their outcomes on easy voluntary movements<sup>[30]</sup> and complicated motor behaviors had been evaluated greater rarely. The relation among remedies for spasticity and for movement, along with walking, is more and more essential because of the excessive occurrence of antispastic remedy withinside the SCI population.




Figure 2: - Factors involved in walking disorders and recovery after SCI.

An epidemiological take a look at of spasticity amongst new SCI sufferers pronounced that one zone had obtained clinical or surgical antispastic remedy earlier than being discharged from rehabilitation. Furthermore, the ones sufferers with Frankel B or C severity of damage have been nearly two times as probable to have obtained antispastic remedy than the others.<sup>[31]</sup> Data from the United States Spinal Cord Injury Model Systems (US-SCIMS) database confirmed that nearly 1- 0.33 of sufferers obtained some shape of antispastic remedy earlier than discharge and nearly half had obtained remedy for spasticity in the first yr after damage. Among individuals with tetraplegia withinside the US-SCIMS database, thirds of these with Frankel B or C severity and a couple of 0.33 of these with Frankel D severity had obtained antispastic remedy inside an yr of damage<sup>[32]</sup> It has Little JW, Goldstein BA, Hammond MC, Dillingham TR, Belandres PV. Spinal cord injury rehabilitation. In Rehabilitation of the Injured Combatant 1998 (Vol. 1, pp. 162-205). Office of the Surgeon General at TMM Publications Institute. There were many pharmacologic techniques developed to deal with the trouble of spasticity in diverse neurologic disorders. Some, e.g., topical anesthetics,<sup>[33,34]</sup> dantrolene sodium,<sup>[35]</sup> and botulinum toxin injections,<sup>[36,37]</sup> were directed at interrupting peripheral factors of the reflex.

Most of the pharmacologic remedies for spasticity, however, are directed at the imperative anxious system (CNS). Their foundation of antispastic impact is their movement at numerous distinct receptor sites in the spinal cord. Many additionally have results at different imperative receptor websites in addition to at peripheral sites, probably contributing to the antispastic impact and really giving upward thrust to a number of the facet results. The records of antispastic remedy consists of the look at of numerous gamma-aminobutyric acid (GABA)-ergic and glycinergic drugs, in addition to noradrenergic, serotonergic, and different classes of drugs.<sup>[38,39]</sup> GABAergic drugs have been widely used for numerous decades to deal with spasticity. They include diazepam, a GABA\* agonist<sup>[40]</sup>; progabide, a GABA\* and GABA\* receptor<sup>[41]</sup> agonist; and baclofen. Baclofen is a GABA agonist that has been found to be exceptionally beneficial for treating spasticity and been called the drug of preference for spasticity from spinal cord disease, at least in part due to the fact that its side-impact profile is relatively benign. For people whose spasticity cannot be controlled with oral baclofen, intrathecal way of drug delivery were explored.<sup>[42-43]</sup> Several glycinergic tablets have also been studied, which include glycine itself and one of its precursors, threonine.<sup>[44]</sup> The use of intrathecal morphine is based on the not unusual affiliation between ache and spasticity.<sup>[45]</sup>




Figure 3: - Pharmacological strategies to address the spasticity after SCI.

### III. INFECTIONS IN SPINAL CORD

Following are some infections seen in spinal cord.

#### A. Spondylitis and Spondylodiscitis

Pyogenic spondylitis is unusual and represents simplest 2–4% of pyogenic osteomyelitis. Mainly adults withinside the 6th to 7th decades is affected. The rule of 50% need to be remembered: 50% of the sufferers are older than 50 years, fever is found in most effective 50% of the cases; genitourinary tract is the primary supply of contamination in 50% of cases, *Staphylococcus aureus* is the causative agent in nearly 50%; the lumbar spine is affected in 50%; sufferers present with a record of greater than three months of signs and symptoms in 50% of cases.<sup>[49,50,51]</sup> Disc hypo-intensity on CT has been stated as being an early sign<sup>[52]</sup>, even though MRI is currently the modality of choice. The exceedingly vascularized vertebral body in adults is affected first and contamination spreads toward the disc area only secondarily.<sup>[53,54]</sup>

#### B. Intramedullary infection

Intramedullary spinal cord abscesses are extraordinarily uncommon lesions. Since the authentic description in 1830, fewer than a hundred instances had been stated withinside the literature.<sup>[55,56]</sup> Their rarity, in comparison with brain abscess, has been attributed to the smaller quantity of the spinal cord and its specific blood supply. Mechanisms of infection consist of hematogenous spread from an extraspinal cognizance of infection, contiguous spread from an adjoining cognizance of infection, direct inoculation (penetrating trauma, post-neurosurgery), and cryptogenic mechanisms (no documented extraspinal cognizance of infection). In a 1998 overview of 25 instances, hematogenous spread accounted for 8%, contiguous spread accounted for 24%, direct inoculation accounted for 4%, and 64% have been cryptogenic.<sup>[57]</sup> Bacteria, consisting of staphylococci and streptococci, are the maximum not unusualplace organisms accountable for those infections; however, infections additionally can be viral, fungal, or resulting from *Cysticercosis*, *Mycobacterium tuberculosis*, *Brucellosis*, *Listeria monocytogenes*, *Toxoplasma gondii*, or different parasites. The maximum usually affected region is the dorsal thoracic spinal cord. The mortality rate withinside the absence of surgical intervention is nearly 100%. The contemporary mortality rate (instances stated withinside the past 2 years) is 8–12%, while 70% of survivors have chronic neurologic deficit.<sup>[58]</sup>

#### C. Epidural Abscess

A current estimate is 2.8 instances per 10,000 admissions to a tertiary care centre. Most instances rise up from hematogenous seeding of the epidural space from a far-off source of infection. When the hematogenous spread influences in simple terms the epidural area the adjoining bony structures can be normal: for those motives especially, MRI need to straight away be used, that is the simplest imaging process to illustrate the lesion in a non-traumatic and simple manner. The thoracic and cervical levels are the maximum generally affected. Another not unusual place etiology is extension of infection from adjoining vertebral osteomyelitis or from a paraspinal or retro-spinal recognition of suppuration. *Staphylococcus aureus* stays the maximum common bacteriologic reason in 62% of infections.<sup>[59]</sup>

### IV. IMAGING TECHNIQUES USED IN SPINAL INFECTION

Following are the imaging techniques used check the spinal cord infection

### A. Skeletal Scintigraphy

The sensitivity of technetium-labelled 99 m bone scintigraphy for the detection of osteomyelitis, discitis, and aseptic spinal illnesses could be very high (>90%), however bone scans are non-specific; fractures or post-surgical adjustments can be accountable for identical modifications. Moreover, specific evaluation of extent, localization, and differentiation among bony factors and gentle tissues are not possible.

### B. Radiography

Plain films can be beneficial for assessing vertebral destruction, however on the early stage of the disorder they will be normal: it is essential to consider that on the time of onset of clinical symptoms, most effective 25% of sufferers show abnormalities. The earliest radiographic sign of infectious spondylitis is an ill-described vertebral end plate, seen possibly as early as 2–3 weeks following infection (Figure 1). Associated with bony impairment, expansion of peri-vertebral gentle tissues can also be detected. Conventional tomography used to reveal those anomalies better, however currently CT is preferred by far. Plain films are nevertheless used for follow-as much as locate vertebral misalignment, secondary bone fusion, or impairment of the spinal dynamics.

### C. Positron Emission Tomography

FDG PET can also additionally show beneficial for differentiation of degenerative and infectious end plate anomalies detected on MR imaging. Even in active (Modic type I) degenerative end plate abnormalities, PET did now no longer show elevated FDG uptake as stated by Stumpe et al. recently.<sup>[60]</sup>

### D. Magnetic Resonance Imaging

Magnetic resonance imaging (MRI) is currently the imaging technique of desire to locate early infection and to assess absolutely the volume of ailment affecting the spine.<sup>[61]</sup> Direct evaluation of the disc space on sagittal images permits early prognosis of end-plate destruction. The earliest reaction to vertebral osteomyelitis is the accumulation of water in extracellular bone marrow, accountable for bone marrow edema. The preferred MR technique for evaluating spinal infection includes consequently a preliminary sagittal spin echo collection using short TR and TE (T1-weighted) observed with the aid of using a sagittal T2-weighted collection and a sagittal SPIR or STIR collection. Additional sagittal and axial T1-weighted sequences after intravenous management of gadolinium-primarily based totally contrast medium are essential. The para spinal gentle tissues are great assessed on multiplanar (axial, coronal) T1-weighted sequences after enhancement and with fats suppression.<sup>[62]</sup>

### V. CONCLUSION

Spinal and spinal cord infections constitute intense pathologic situations that need to be recognized in an early stage, averting extreme musculoskeletal and neurologic complications. Magnetic resonance imaging has the key function to play and need to be employed as quickly as infection is clinically suspected. Also, the comprehensive assessment of walking behavior requires obligations regarding the distinctive control variables. This comprehensive assessment may be used to represent the procedure of restoration of walking in addition to the effectiveness of numerous treatments.

### VI. REFERENCES

- [1] Vandavelde AG, Jenkins SG, Hardy PR. Sclerosing osteomyelitis and *Actinomyces naeslundii* infection of surrounding tissues. *Clinical infectious diseases*. 1995 Apr 1;20(4):1037-9.
- [2] Tyrrell PN, Cassar-Pullicino VN, McCall IW. Spinal infection. *European radiology*. 1999 Jul;9(6):1066-77.
- [3] American Thoracic Society. Control of tuberculosis in the United States. *Am Rev Respir Dis*. 1992;146:1623-33.
- [4] MacGregor RR. Tuberculosis: from history to current management. In *Seminars in roentgenology* 1993 Apr 1 (Vol. 28, No. 2, pp. 101-108). WB Saunders.
- [5] Knutsson E, Richards C. Different types of disturbed motor control in gait of hemiparetic patients. *Brain: a journal of neurology*. 1979 Jun 1;102(2):405-30.
- [6] Dietz V, Quintern J, Berger W. Electrophysiological studies of gait in spasticity and rigidity. Evidence that altered mechanical properties of muscle contribute to hypertonia. *Brain: a journal of neurology*. 1981 Sep 1;104(3):431-49.
- [7] Conrad B. Pathophysiological aspects of human locomotion. Motor control mechanisms in health and disease.. 1983:717-27.
- [8] Wainberg M, Barbeau H, Gauthier S. Quantitative assessment of the effect of cyproheptadine on spastic paretic gait: a preliminary study. *Journal of neurology*. 1986 Oct;233(5):311-4.
- [9] Wagenaar RC, Beek WJ. Hemiplegic gait: a kinematic analysis using walking speed as a basis. *Journal of biomechanics*. 1992 Sep 1;25(9):1007-15.
- [10] Pepin A, Barbeau H. Effects of treadmill speed and incline on the walking pattern of normal and spastic paretic subjects. In *Soc Neurosci Abstr* 1992 (Vol. 18, p. 860).
- [11] Nilsson J, Thorstensson A, Halbertsma JN. Changes in leg movements and muscle activity with speed of locomotion and mode of progression in humans. *Acta Physiologica Scandinavica*. 1985 Apr;123(4):457-75.
- [12] Conrad B. Gait disturbances in paraspastic patients. *Clinical neurophysiology in spasticity*. 1985:155-74.
- [13] Fung J, Barbeau H. A dynamic EMG profile index to quantify muscular activation disorder in spastic paretic gait. *Electroencephalography and clinical neurophysiology*. 1989 Sep 1;73(3):233-44.
- [14] Fung JO, Barbeau HU. Effects of conditioning cutaneous stimulation on the soleus H-reflex in normal and spastic paretic subjects during walking and standing. *Journal of neurophysiology*. 1994 Nov 1;72(5):2090-104.
- [15] Mirbagheri MM, Kearney RE, Barbeau H, Ladouceur M. Increased reflex contributions to dynamic ankle stiffness in spastic spinal cord injured patients. In *Proceedings of 18th Annual International Conference of the IEEE Engineering in Medicine and Biology Society* 1996 (Vol. 2, pp. 607-608). IEEE.
- [16] Sinkjær T, Magnussen I. Passive, intrinsic and reflex-mediated stiffness in the ankle extensors of hemiparetic patients. *Brain*. 1994 Apr 1;117(2):355-63.
- [17] Dietz V, Berger W. Normal and impaired regulation of muscle stiffness in gait: a new hypothesis about muscle hypertonia. *Exp Neurol* 1983;79:680-7

- [18] Given JD, Dewald JP, Rymer WZ. Joint dependent passive stiffness in paretic and contralateral limbs of spastic patients with hemiparetic stroke. *Journal of Neurology, Neurosurgery & Psychiatry*. 1995 Sep 1;59(3):271-9.
- [19] Yang JF, Fung J, Edamura M, Blunt R, Stein RB, Barbeau H. H-reflex modulation during walking in spastic paretic subjects. *Canadian Journal of Neurological Sciences*. 1991 Nov;18(4):443-52.
- [20] Delwaide PJ. Human monosynaptic reflexes and presynaptic inhibition. In *Human reflexes, pathophysiology of motor systems, methodology of human reflexes 1973* (Vol. 3, pp. 508-522). Karger Publishers.
- [21] Faist M, Mazevet D, Dietz V, Pierrot-Deseilligny E. A quantitative assessment of presynaptic inhibition of Ia afferents in spastics: Differences in hemiplegics and paraplegics. *Brain*. 1994 Dec 1;117(6):1449-55.
- [22] Crone C, Nielsen J, Petersen N, Ballegaard M, Hultborn H. Disynaptic reciprocal inhibition of ankle extensors in spastic patients. *Brain*. 1994 Oct 1;117(5):1161-8.
- [23] ASHWORTH B. Preliminary trial of carisoprodol in multiple sclerosis. *Practitioner*. 1964;192:540-2.
- [24] Lance JW. Symposium synopsis. Spasticity: disordered motor control. 1980:487-9.
- [25] Young RR. Drug therapy: spasticity (first of two parts). *N Engl J Med*. 1981;304:28-33.
- [26] Wiesendanger M. Neurophysiological bases of spasticity. In *Neurosurgery for spasticity 1991* (pp. 15-19). Springer, Vienna.
- [27] Young RR. Spasticity: a review. *Neurology*. 1994 Nov 1;44(11 Suppl 9):S12-20.
- [28] Meythaler JM, McCary A, Hadley MN. Prospective assessment of continuous intrathecal infusion of baclofen for spasticity caused by acquired brain injury: a preliminary report. *Journal of neurosurgery*. 1997 Sep 1;87(3):415-9.
- [29] Nance PW, Bugaresti J, Shellenberger K, Sheremata W, Martinez-Arizala A. Efficacy and safety of tizanidine in the treatment of spasticity in patients with spinal cord injury. North American Tizanidine Study Group. *Neurology*. 1994 Nov 1;44(11 Suppl 9):S44-51.
- [30] Latash ML, Penn RD, Corcos DM, Gottlieb GL. Effects of intrathecal baclofen on voluntary motor control in spastic paresis. *Journal of neurosurgery*. 1990 Mar 1;72(3):388-92.
- [31] Little JW, Goldstein BA, Hammond MC, Dillingham TR, Belandres PV. Spinal cord injury rehabilitation. In *Rehabilitation of the Injured Combatant 1998* (Vol. 1, pp. 162-205). Office of the Surgeon General at TMM Publications Institute.
- [32] Barbeau H, Ladouceur M, Norman KE, Pépin A, Leroux A. Walking after spinal cord injury: evaluation, treatment, and functional recovery. *Archives of physical medicine and rehabilitation*. 1999 Feb 1;80(2):225-35.
- [33] Mills III WJ, Pozos RS. A decrease in clonus amplitude by topical anesthesia. *Electroencephalography and clinical neurophysiology*. 1985 Dec 1;61(6):509-18.
- [34] Sabbahi MA, De Luca CJ, Powers WR. Topical anesthesia: a possible treatment method for spasticity. *Arch Phys Med Rehabil*. 1981 Jul 1;62(7):310-4.
- [35] Pinder RM, Brogden RN, Speight TM, Avery GS. Dantrolene sodium. *Drugs*. 1977 Jan;13(1):3-23.
- [36] Dengler R, Neyer U, Wohlfarth K, Bettig U, Janzik HH. Local botulinum toxin in the treatment of spastic drop foot. *Journal of neurology*. 1992 Jul;239(7):375-8.
- [37] Hesse S, Krajnik J, Luecke D, Jahnke MT, Gregoric M, Mauritz KH. Ankle muscle activity before and after botulinum toxin therapy for lower limb extensor spasticity in chronic hemiparetic patients. *Stroke*. 1996 Mar;27(3):455-60.
- [38] Davidoff RA. Antispasticity drugs: mechanisms of action. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*. 1985 Feb;17(2):107-16.
- [39] Mondrup K, Pedersen E. The effect of the GABA-agonist, progabide, on stretch and flexor reflexes and on voluntary power in spastic patients. *Acta neurologica scandinavica*. 1984 Apr;69(4):191-9.
- [40] Davidoff RA. Antispasticity drugs: mechanisms of action. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*. 1985 Feb;17(2):107-16.
- [41] Mondrup K, Pedersen E. The effect of the GABA-agonist, progabide, on stretch and flexor reflexes and on voluntary power in spastic patients. *Acta neurologica scandinavica*. 1984 Apr;69(4):191-9.
- [42] Albright AL, Barron WB, Fasick MP, Polinko P, Janosky J. Continuous intrathecal baclofen infusion for spasticity of cerebral origin. *Jama*. 1993 Nov 24;270(20):2475-7.
- [43] Coffey RJ, Cahill D, Steers W, Park TS, Ordia J, Meythaler J, Herman R, Shetter AG, Levy R, Gill B, Smith R. Intrathecal baclofen for intractable spasticity of spinal origin: results of a long-term multicenter study. *Journal of neurosurgery*. 1993 Feb 1;78(2):226-32.
- [44] Stern P, Bokonjić R. Glycine therapy in 7 cases of spasticity. *Pharmacology*. 1974;12(2):117-9.
- [45] Lee A, Patterson V. A double-blind study of L-threonine in patients with spinal spasticity. *Acta neurologica scandinavica*. 1993 Nov;88(5):334-8.
- [46] Davidoff RA. Antispasticity drugs: mechanisms of action. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*. 1985 Feb;17(2):107-16.
- [47] Palusci VJ, Lazoritz S. *The shaken baby syndrome: a multidisciplinary approach*. Routledge; 2002 Feb 7.
- [48] Mai J, Pedersen E. Clonus depression by propranolol. *Acta neurologica scandinavica*. 1976 Jun;53(5):395-8.
- [49] Saint Amour THE, Hodges SC, Laakman RW, Tamas DE (1994) MRI of the spine. Raven Press, New York, pp 594–667.
- [50] Manelfe C (1992) *Imaging of the spine and spinal cord*. Raven Press, New York.
- [51] Colombo N, Berry I, Norman D (1992) Infections of the spine. In: Manelfe C (ed) *Imaging of the spine and spinal cord*. Raven Press, New York, pp 489–512.
- [52] Larde D, Mathieu D, Frija J, Gaston A, Vasile N (1982) Vertebral osteomyelitis: disc hypointensity on CT. *Am J Roentgenol* 139:963–967.
- [53] Thrush A, Enzmann D (1990) MR imaging of infectious spondylitis. *Am J Neuroradiol* 11:1171–1180.
- [54] Maiuri F, Iaconetta G, Gallicchio B, Manto A, Briganti F (1997) Spondylodiscitis: clinical and magnetic resonance diagnosis. *Spine* 22:1741–1746.
- [55] Quencer RM, Post MJD (1997) Spinal cord lesions in patients with AIDS. *Neuroimaging Clin North Am* 7:359–366.
- [56] Murphy KJ, Brunberg JA, Quint DJ et al. (1998) Spinal cord infection: myelitis and abscess formation. *Am J Neuroradiol* 19:341–348.
- [57] Chan CT, Gold WL (1998) Intramedullary abscess of the spinal cord in the antibiotic era: clinical features, microbial etiologies, trends in pathogenesis, and outcomes. *Clin Infect Dis* Sep 27:619–626.

[58] Weng TI, Shih FY, Chen WJ, Lin FY (2001) Intramedullary abscess of the spinal cord. Am J Emerg Med 19:177–178.  
[59] Chimelli L (2001) Tumors and tumorlike lesions of the spine and spinal cord. Neuroimaging Clin North Am 11:79–111.  
[60] Stumpe KD, Zanetti M, Weishaupt D, Hodler J, Bboos N, Schulthess GK von (2002) FDG positron emission tomography for differentiation of degenerative and infectious endplate abnormalities in the lumbar spine detected on MR imaging. Am J Roentgenol 179:1151–1157.  
[61] Sharif H (1992) Role of MR imaging in the management of spinal infections. Am J Roentgenol 158:133–1345.  
[62] Sharif H, Clark D, Aabed M et al. (1990) Granulomatous spinal infection: MR imaging. Radiology 177:101.

