



INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

Estimation Of herbicides In Environmental Samples

Balabharathi¹, Ramanamma¹ and Rajasekhar reddy²

1.principal and Head department of botany, yarlagadda annapoornamba
govt.degrecollege,chirala,prakasham dist,andhrapradesh,india.

1.senior lecturer in chemistry yarlagadda annapoornamba govt.degrecollege,chirala,prakasham
dist,andhrapradesh,india.

2.senior lecturer in chemistry yarlagadda annapoornamba govt.degrecollege,chirala,prakasham
dist,andhrapradesh,india.

Abstract

This approach is to estimate the residual amounts of herbicides present in environmental samples by applying electro chemical technique adsorptive stripping voltammetry. Average amounts for ten replicates founded by using carbon nano tubes paste electrodes as working electrodes. statistical concepts such as standard deviation and correlation coefficient and in all the findings in this approach all the possible errors are minimised and accuracy is maximised. Water samples of various areas are collected and investigated for pesticide residues before and after the application of pesticides.

Key words: herbicides, adsorptive stripping voltammetry, carbon nano tubes paste electrodes ,water samples.

Introduction

Herbicides are chemicals used to manipulate or control undesirable vegetation. Herbicide application occurs most frequently in row-crop farming, where they are applied before or during planting to maximize crop productivity by minimizing other vegetation. They also may be applied to crops in the fall, to improve harvesting.

Despite the fact that herbicides are useful for the control of various pests, many of them are hazardous chemicals. They are perilous because they can poison the land, the water and the air. Some pesticides do not break down for a long time. These types of pesticides are often used when something must be protected from pest attack for a long period of time, for example, protecting houses from termite attack.

Pesticides which remain in the soil or on the treated surface are also often called residual chemicals[1-7]. When residual pesticides get into the environment they can remain poisonous and active for many years. If applied incorrectly or used in the wrong place, these chemicals may spread to other land areas and possibly to the water supply.

There are good reasons (advantages) pesticides are very effective. This means that nearly all the target pests which come in contact with these pesticides are killed. Results are quick. This means the pests are killed within a very short time.

Using pesticides can be an economical (cheap) way of controlling pests. Pesticides can be applied quickly and there is not the high labour cost which might apply to other methods of control, such as removing weeds by hand.

If pesticides are not used correctly, they can affect human health or cause serious injury or death to the pesticide operator, other people or household pets. Pesticides can also directly affect other non-target animals. For example, a gardener spraying his garden to kill caterpillars will probably also kill harmless lady bird beetles and praying mantises. If pesticides are used incorrectly or applied wrongly, they may find their way into places where they are not wanted, for example, they might be washed into rivers or into the soil. In this article an electroanalytical method voltammetry supported by statistical findings was applied.

Instruments and reagents

Electro analytical determinations conducted using a model meterohm Auto Lab 101 PG stat (Netherlands). CNTPE was used as working electrode for differential pulse adsorptive stripping voltammetry and cyclic voltammetry. pH measurements were carried out with an Eutech PC_510 cyber scan. Meltzer Toledo (Japan) Xp26 delta range micro balancer were used to weigh the samples during the preparation of standard solutions. All the experiments were performed at 25°C.

All reagents used are analytical reagent grade. Double distilled water was used throughout the analysis. In the present investigation universal buffers of pH 4.0 was used as supporting electrolytes and are prepared by using 0.2 M boric acid, 0.05M citric acid and 0.1M trisodium orthophosphate solutions.

Measurements and calculations

In this standard addition method, the voltammogram of the unknown is first recorded after which a known volume of standard solution of the same electro active species is added to the cell and second voltammogram is taken. From the magnitude of the peak height, the unknown concentration of species may be calculated using the following equations.

$$C (\text{un known}) = \frac{C_s \times V}{V_t \times i_2} \times i_1$$

Result And Discussions

Well resolvable and reproducible peak obtained for each sample is useful for the analysis of water samples. The optimum pH to get well defined peak for the detection is found to be 4.0. The peak current is found to vary linearly with the concentration of the pesticide over the range $1.0 \times 10^{-5}\text{M}$ to $1.0 \times 10^{-9}\text{M}$. The lower

detection was limit found to be 1.02×10^{-9} M. The correlation coefficient and relative standard deviation (for 10 replicates) obtained using the above procedure [8-15].

Recovery experiments

A stock solution (1.0×10^{-3} M) of each sample is prepared in dimethyl formamide. In voltammetric cell, 1 mL of standard solution is taken and 9 mL of the supporting electrolyte (pH 4.0) is added to it. Then the solution is de aerated with nitrogen gas for 10 min. after obtaining the voltammogram, small additions of standard solution are added and the voltammograms are recorded under similar experimental conditions. The optimum conditions for analytical estimation at pH 4.0 are found to be pulse amplitude of 25 mV, applied potential of -0.35V and scan rate 40 mVs^{-1} .

Water samples are collected from paddy fields which sprayed by the pesticides under investigation 48 hours after spraying the pesticides. These samples were filtered through a Whatman No.41 filter paper and Aliquots of water samples were taken in a 25mL graduated tube, to it buffer solution was added and analyzed as described above. The recoveries of samples obtained in water samples ranged from 51.00 to 57.00% and the results are summarized in Table 1.0.

Table 1.0: Recoveries of herbicides in water

samples

Name of the pesticide	Amount added (mg/L)	Amount found (mg/L)	*Recovery (%)	Standard deviation
1.attrazine	4.0	2.15	53.75	0.07
2.Dicamba	4.0	2.36	59.00	0.05
3.trifluralin	4.0	2.31	57.75	0.16
4.cyanazine	4.0	2.25	56.25	0.06
5.pendimethalin	4.0	2.10	52.50	0.17

***Average of 10 replicates**

Conclusions

In this attempt statistical parameters for the determination of pesticide residues satisfactory applied to interpret the instrumental out puts without considerable errors. And during the estimations pollution arises due to heavy metal electrodes such as mercury electrodes is avoided by using carbon electrodes.

References

- 1.S.Rajasekharreddy, K.Chandramohan and NY.sreedhar,International journal of scientific and engineering research. 2,2011, 10.
2. SarvareddyRajasekhar Reddy , NeelamY.Sreedhar, Kukkambacum Chandra mohan and MaddelaNagaraju.Global journal of science frontier research-B, 12, 2012,1.
3. ThommandruRaveendranathBabu, SarvareddyRajasekharReddy,PuchakayalaSujana, J. Electrochem. Sci. Eng. 4 ,2014,1.
- 4.T.R.Babu,P. Sujana,S.R. Sekhar Reddy, J. Atoms and Molecules/ 4,2014,2.
- 5.SarvareddyRajasekhar Reddy and T. Raveendranathbabu,int. J. Nanosci. 12, 2013.
- 6.S. RajasekharReddy,T.Raveendranathbabu, B.Sreenivasulu , International Journal of Research in Pharmacy and Life Sciences , 1,2013, 1.
- 7.T.Raveendranath Babu, P.Sujana, K.Sivasankar, S.Rajasekhar Reddy, Analytical Chemistry an Indian Journal, 11, 2012, 2.
- 8.Feride Koc1, Yusuf Yigit, YavuzKursad Das, YaseminGure and CevdetYarali, Journal of Food and Drug Analysis, 16, 2008,3.
- 9.Jose Fernando Huertas-Perez and Ana María Garcia-Campana, AnalyticaChimica Acta,630, 2008,2.
10. Keith M. Moore, Susan R. Jones and Carole James, Water Research 29, 1995, 5.
- 11.G. Hoizey, F. Canas, L Binet, M. L. Kaltenbach, G. Jeunehomme, M. H. Bernard, D. Lamiable, J. Forensic Sci. 53 ,2008, 499-502.
12. G. Xu, W. Zheng, Y. Li, S. Wang, J. Zhang, Y. Yan., Int. Biodeter. Biodegr.62 ,2008, 51–56.
13. Susana de Melo Abreu; Paulo Herbert; PierluigiCaboni; Paolo Cabras,ArmindaAlves; Vincenzo Luigi Garau, Journal of Environmental Science and Health, Part B, 42, 2007, 7.

14. De Melo Abreu S, Caboni P, Cabras P, Garau VL, Alves A., Anal Chim Acta., 573-574, 2006, 291-7.
15. Anne Danion, Jean Disdier Chantal Guillard, Olivier Païssé, and Nicole Jaffrezic-Renault, Applied Catalysis B: Environmental, 62, 2006, 22.
16. Vincenzo Luigi Garau, Susana De Melo Abreu, Pierluigi Caboni, Alberto Angioni, Arminda Alves and Paolo Cabras, J. Agric. Food Chem., 57, 2009, 6.
17. André Schreiber, Yuriko Ozeki, Applied bio systems 2008.
18. Lutz Alder, Kerstin Greulich, Günther Kempe, Bärbel Vieth Mass Spectrometry Reviews, 25, 6, 2006, 838 – 865.
19. S. M. Waliszewski, G. A. Szymczyński, Fresen. J. Anal. Chem. 338 (1990) 75-76
20. N. Unceta, A. Ugarte, A. Sanchez, A. Gómez-Caballero, M. A. Goicolea, R. J. Barrio, J. Chromatogr. A 1061, 2004, 211-216.
21. S. Rajasekharreddy, K. Chandra Mohan and, NY. Sreedhar, Int. J. Sci. Eng. Res, 2, 2011, 10.
22. Ozge Surucu Gulcin Bolat Serdar Abaci, Talanta, 168, 1, 2017, 113-120.

