



MEDICINAL PROPERTIES OF NEEM- A REVIEW

Vachanaram Natharam Choudhary¹ Ashwin Somkuwar²

Siddhi Patil³

M.S College of Pharmacy Devghar, Guarapur-Kudus Rd, Dist. Palghar, Maharashtra 421312

ABSTRACT

Popular tree Neem (*Azadirachta indica*) is a member of the Meliaceae family. Many foods are prepared using bark, leaves, flowers, seeds, oil, and other materials in the Traditional System of Medicine. For pharmaceutical uses, neem has been the subject of several patent applications. The pharmacognosy, phytochemistry, and pharmacological activities that have been documented thus far are all thoroughly examined in this review. As a source of several therapeutic substances in Indian culture and a plant that thrives in tropical and semi-tropical regions, neem is the most beneficial traditional medicine. All parts of the neem tree—bark, leaves, fruits, seeds, and extracts—are utilized in traditional medicine. Its extracts are anti-inflammatory, anti-viral, anti-bacterial, anti-fungal, anthelmintic, and anti-allergic. The names "free tree of India," "wonder tree," "Nature's drug store," "Village dispensary," "Divine tree," "heal all," and "Panacea of all Diseases" are also used to neem. In current times, non-toxic herbal items should be used with specific concentration to control animal and human ailments. There is much room for this amazing plant to be used more effectively.

KEYWORDS

Neem, Therapeutic and diseases, Pharmacognosy, Pharmacological activity.

INTRODUCTION

The valuable medicinal tree *Azadirachta indica* is a member of the Meliaceae family. Due to the positive benefits neem has on human illnesses, all parts of the plant have achieved worldwide recognition. *Azadirachta indica* has been a plant in India for 4,000 years. (1) *A. indica* is sometimes referred to as the Margosa tree, and in Sanskrit, it means "to give good health" as "arista" or "nimba" and "nimbati swasthyamdadati." The "Charak-Samhita" and "Susruta-Samhita," which serve as the foundation for the Ayurvedic medical system, highlight

the biological advantages of neem. In Persian, "Azad- Darakth- E- Hind" means "Free tree of India." Neem is a plant that has a complex genetic makeup. Neem was utilized in ancient India to treat smallpox and chicken pox. It has been applied to prevent environmental problems such soil erosion, soil fertility, pesticides, and insecticides, among others. (2)



NEEM LEAVES



NEEM SEEDS

NEEM'S SCIENTIFIC CLASSIFICATION

Kingdom: Plantae

Division: Magnoliophyte

Class: Magnoliopsida

Order: Sapindales

Family: Meliaceae

Genus: *Azadirachta*

Species: *indica*

Scientific name: *Azadirachta indica*

VERNACULAR NAMES

Hindi – Nim, Nimb

English – Margosa, Neem, Indian Lilac

Konkani – Beva-rooku

Bengali – Nim, Nimgach

Marathi – Kadunimb

Gujarathi – Limba

Tamil – Veppamaram, Vempu, Veppam

Punjabi – Nimb

Malayalam – Veepu, Aryaveppu, Aruveppu, Kaippan, Veppu, Vepa

Oriya – Nimo

Telugu – Vepa, Yapachettu

Kannada – Bevinmar, Kahibevu (7)

DESCRIPTION OF BOTANY

Features at the Macroscopic Level

A huge, evergreen tree with a bitter flavor that is 12 to 18 meters tall.

Bark features: scaly, oblique furrows, thick, strong, woody, dark brown to greyish, sharp exterior surface, distinct odour, and straight trunk.

Leaves: The leaves are complex, ovate-lanceolate, attenuate, opposite, imparipinnate (5–15 leaflets), with a characteristic odour and dark green colour. They also have serrated edges and short, 3–8 cm long petioles.

Fruits: drupe-like, glabrous, green, oblong to roundish, tasty, with thin skin.

Flowers: Protandrous, bisexual, white, fragrant flowers are grouped in axillary panicles.

Seed: One single seed, greasy, gritty, and dirty brown in shade.

STANDARD USES AND ACTIONS

Bark: It is astringent, acrid, pectoral, tonic, and used to treat hyperdipsia, leprosy, skin conditions, eczema, leucoderma, pruritis, malaria, burning sensation, tumor, dyspepsia, intestinal worms, hepatopathy, lumbago, ulcers, amenorrhea, lumbago, hemorrhoids, otalgia, syphilis, and exhaustion.

Leaves: This herb has antiseptic, ophthalmic, ophthalmopathy, intestinal worms, dyspepsia, ulcer, tuberculosis, boils, eczema, malaria, and intermittent fever properties. It is also anthelmintic, insecticidal, demulcent, and refrigerant for skin conditions, leucoderma, burning feeling, leprosy, and itch.

Flowers: It is used to treat intestinal worms, colic, dyspepsia, burning feeling, ophthalmopathy, ophthalmic, stomachic, anthelmintic, and general debility.

Seeds: It is used to treat tumors, leprosy, skin conditions, odontalgia, intestinal worms, wounds, ulcers, diabetes, vulnerability, emollient, anodyne, anthelmintic, depurative, and uterine stimulant.

Oil: It is used to treat leprosy, syphilitic sores, ulcers, ringworm, scabies, fever, and chronic skin conditions. It is also an anodyne and depurative. (1,2,4,7)

HABITATION

The neem tree is believed to have originated naturally in south Asia, while its precise original place is unknown. It grows in the dry natural forests of southern India and Burma. Neem has been grown for millennia in Southern Malaysia, the drier Indonesian islands from Java eastward, Bangladesh, Myanmar, Thailand, India, Pakistan, and Sri Lanka. Neem was brought to Fiji and Mauritius in the early 1700s.

GROWTH AND DIVISION

This hardy tree thrives in drought-prone areas and saline soils. Under conditions of waterlogging, growth is slow. propagated from seeds, which have a very limited viability period and should be sown as soon as possible after maturing. Three weeks is all it takes for seeds to sprout. Stem cuttings and root suckers are also utilized in planting. Although it may be cultivated in any kind of soil, black-loam soil is the most ideal. The seedlings reach a height of 120 cm in a year. It has been discovered that leaf culture works well for rapid multiplication. It has been claimed that azadirachtin can be produced from cultures of leaves and flowers using tissue culture techniques. On a dry weight basis, it has been observed that the callus of leaves that are 20 weeks old can contain up to 2.68 percent azadirachtin, while the callus of flowers that are 12 weeks old can contain 2.46 percent.

CLIMATE

It is a drought-resistant tree that grows best in average yearly temperatures between 21 and 32°C. Neem grows best in India in temperatures between 0 and 49°C and with less than 600 mm of annual rainfall. Additionally, the plant is utilized in afforestation projects in semi-arid and dry areas. (4)

AYURVEDIC PROPERTIES

Rasa: Tikta, Kashaya

Guna: Laghu

Veerya: Sheeta

Vipaka: Katu

Doshagnata: Kaphapitashamaka

Rogagnata: Vidradhi, Granthi, Amavata, Vamana, Kaphapaittika etc.

Karma: Vranapachana, Vranashodhana, Ampachana, Juaraghna, Vishamapwara pratibandhaka, Chakshushya etc.

DOSES

Bark powder: 2 to 4 gm

Leaf juice: 10 to 20 ml;

Powder: 2-4 g.

Oil- 5 to 10 drops (11)

CHEMICAL CONSTITUENTS

More than 100 chemicals, primarily pentanortriterpenoids, tetranortriterpenoid- γ hydroxy butenolides, limonoids (or meliacins or tetranortriterpenoids), and protolimonoids (proto-meliacin) triterpenoids. Aside from a few nontriterpenoid components, hexanortriterpenoid constituents have been recorded from different regions, as listed below.

Leaves: Azadirachtin, azadirachtol, azadirachnol, deacetyl-azadirachtinol (-3- tigloylazadirachtol), azadiradione, an isomer of epoxyszadiradione, 17 epi and 17-hydroxyazadiradione, gedunin, 7-hydroxygedunin, melianone, bol nimboeinol (7-acetoxy-7-hydroxy azadiradione), nimocin, 7-deacetoxy-nimolicinol, nimolinone, nimbochalcin, and nimboctin, Diepoxytirucall-7-en-21-ol salannin, azadirachtanin, azadirone, azadiradione, epoxyazadiradione, isoazadirolide, nimbandiol, nimbinene desacetylnimbinene, nimbin, nimbocinolide, isonimbocinolide, 21, 23, 24, 25-salannin, the compounds nimbolide, nimocinolide, isonimocinolide, nimocinone, kaempferol-3-O-8-glucoside, myricetin, 3-L-arabinoside (mellictrin), 3-O-L-ghanmonide and 3-O-rutinoside, Quinolones, scopoletin, β -sitosterol and its β -D-glucoside, amino acids, β -carotene, carbohydrates, n-hexacosanol, nomacommol, protein, and vitamins are among the substances that include quercetin, its 3-galactoside, 3-O-L-thanoside, and 3-O-runinoside.

Fruits & Seeds: Kernels provide between 400 and 48.9% of the oil and are rich in tocopherol, arachidic, linoleic, margosic, myristic, oleic, palmitic, and stearic acids, gedunin, meldenin, meliatriol, nimbine, and mbinene. 6-Nimbinene, nimbidin, nimbidiol, desacetyl Nimbidic acid, 6-O-acetylnimbandiol, salannin, 3-desacetylsalannin, salannol and its acetate, salannolide, vepinine, vilasinin, 1,3-diacetylvilasinin, 1-tigloyl-3-acetyl-vilasinin, and tiglic acid (seed oil), azadirachtin, 22,23-dihydro-23-8-methorey- azadirachtin (vepaol) and its C-23 epimer (isovepaol), Azadirone, azadiradione, epoxyazadiradione, 2-dihydro-epoxyazadiradione,

18.28-diepoxyazadiradione, nimbidin, nimbidinin, salannic (nimbidic) acid, and 7-desacetyl-7-benzoyl-geduni acetylneotrichilenone are examples of compounds with this structure (seeds).

Flowers: Benzyl alcohol, β -sitosterol, thioamyl alcohol, oleic, palmitic, stearic, arachidic, behenic, colic, kaempferol, and its 3-glucoside, myricetin-3-L-arabinoside, quercetin-3-galactoside, azadirachtin, and margosene may all be found. Diepoxyazadirol, flowerine, flowerone, and O-methylazadiranolide. Triterpenoid (trichilenone acetate), flavanones, nimbaflavone, 3'-prenylnaringenin, and 4'-(2'hydroxyethyl) phenol are other components of flowers that are known to exist.

Heartwood: 4,14, α -dimethyl-3- α -ergosta-8, 24-methylene-cycloartenol, β -sitosterol and its glucoside, and tannin 24(28)-d-38-ol 12-methyl-Sa-ergosta-8, 24(28)6-desacetyl nimbinene, nimatone, nimbinene, nimbolins A and B, and dien-38-ol.

Stem bark contains the following compounds: vanilic acid, catechol, campesterol, stigmasterol, sitosterol, β -amyrin, lupeol, nimbin, nimbidin, nimbinin supiol, kulinone, kulactone, kulolactone methyl kulonate, 60-hydroxy-4-stigmasten-3-one, and 68-hydroxy-4-campesten-3-one.

Wood: fraxinellone, nimbolin, 7-deacetoxy-7-axogedunin, and gedunin A gydoeucalene, A and B melanin.

Twigs: Isomargosinoside, desacetyl isonimbinoside, desacetyl nimbinolide, and margosinoside.

Wood Oil: 24-methylene cycloartenol, β -sitosterol, and cycloeucalenol.

Trunk bark: Nimbiol, sugiol, and bosterol nimbolins A and B (trunk wood).

Root: 4-campesten-3-one, 4-stigmasten-3-one, trans-cinnamic and vanillic acids, nimbin and nimbidin, 24-methylene-cycloartenol, 24-methylene-cycloartanone, cycloeucalenol, cycloeucalenone.

In addition to aesculetin, cholesterol, Isomeldenin, meldonindiol, 4- and 6-hydroxy-7-methoxy-coumarin, 4- and 6-dihydroxy-A-homoazadirone, From different portions of the tree, 17-acetoxy-meliacin, 6-O-acetylnimbandiol, desacetylnimbin, nimocinol, isonimolicinolide, and nimolinolic acid have been discovered. (12,17)

QUANTITATIVE STANDARD

Foreign matter: Not more than 0.3 % w/w

Ash: Not more than 3.0 % w/w

Acid insoluble ash: Not more than 0.25 % w/w

Alcohol soluble extractive: Not less than 4.5 % w/w

Water soluble extractive: Not less than 5.0 % w/w (11,18)

COMPOUNDS AND PROCEDURES

Nimbarishta, Mahagandhaka vati, Panchaguna taila, Madhyam narayana taila, Guduchyadihana kvatha, Shankhpushpi taila, Jwarasamhara rasa, Mahamarichyadi taila, Pathyadi kvatha, Mahamanjishthadyarishta, Punarnavadi kvatha, and Mahasudarshana churna, Raktashodhaka vati, Nimbaharidra khanda, Nimbarishta, Mahagandhaka vati, Madhyam narayana taila, and Guduchyadihana kvatha. (11)

THERAPEUTIC SKILLS

The Table below lists the phytoconstituents' pharmacological actions that have been documented thus far for this plant.

Chemical constituents	Biological activity
Epoxyazadiradione	Human cervical cancer (22)
Sodium nimbidate	Anti-inflammatory
Nimbidin	Anti-inflammatory Antiarthritic Antipyretic Hypoglycaemic Antigastric ulcer Spermicidal Antifungal Antibacterial Diuretic
Nimbin	Spermicidal
Nimbolide	Antibacterial Antimalarial
Gedunin	Antifungal Antimalarial
Azadirachtin	Antimalarial
Mahmoodin	Antibacterial
Gallic acid, (-) epicatechin and catechin	Anti-inflammatory and Immunomodulatory
Margolone, margolonone and isomargolonone	Antibacterial
Cyclic trisulphide, cyclic tetrasulphide	Antifungal
Polysaccharides	Anti-inflammatory

Polysaccharides GIa, GIb	Antitumour
Polysaccharides GIIa, GIIIa	Anti-inflammatory
NB-II peptidoglycan	Immunomodulatory (13-15)
Glycoprotein	Immunomodulatory (19)
Proline	Alzheimer's and Parkinson's disease (20,21)

ESTIMATION BY HPTLC

Method is as per identification method, with the exception that the plate is scanned at 424 nm after drying.

The peak area under the curve is used to determine the content of the sample.

RECOGNITION THROUGH TLC

Phase of stationery: precoated plates with silica gel GF254


Mobile phase: Glacial acetic acid: toluene: ethyl acetate: 5:5:1.

Standard preparation: 5 cc of methanol is mixed with 10 milligrams of epicatechin.

Sample preparation: Reflux 5 g of medication for 8 hours [two times] with methanol. After filtering, extract the solvent. Soak in ten millilitres of methanol.

Spray reagent: Use a 10% potassium hydroxide ethanol solution and observe under a UV light source at 365 nm.

Detection: In both the standard and test, epicatechin is shown to have Rf 0.29.

PHARMACOLOGICAL ACTIVITY**SAFETY ASPECTS**

The medication may be regarded as safe when taken as directed at customary dosages.

ACKNOWLEDGEMENT

I would want to express my gratitude to my friends who have been instrumental in our growth both at the institutions and throughout my graduate and undergraduate courses. As my careers have progressed, I have also had the opportunity to interact with numerous distinguished pharmacy individuals and outstanding instructors from various fields, all of whom have had a significant impact on my work and provided guidance. These people have taught me a lot, therefore I'm grateful they shared their wisdom and experiences with us.

Appreciation should also be extended to the entire faculty and staff of the M.S. College of Pharmacy at Mumbai University.

I would like to express my gratitude to the writers of each of our chapters for their diligent efforts. In addition to their educational backgrounds in pharmacy and pharmacy practice, the writers were selected for

their works based on their commitment to these areas. Without their help, I could not have finished composing the ensuing piece.

CONCLUSION

Neem is a plant that can be used to treat a variety of illnesses. Its anti-inflammatory, wound-healing, antipyretic, antiallergic, nephroprotective, hepatoprotective, immunomodulatory, antifertility, dental hygiene, antidiabetic, cardioprotective, antimicrobial, anticancer, and pesticide properties have all been scientifically demonstrated. This review provides information on the molecules that were isolated, their pharmacological action, and the extract. For neem and its products to be used more profitably and therapeutically, a great deal of research and development needs to be done. With these advantages in mind, a medication development program exploiting neem's biological and therapeutic qualities ought to be launched. In the present period, non-toxic herbal items should be used with specific concentration to control animal and human ailments. Even if some neem herbal products have been made, there is still much room for improving the way this amazing plant is used.

REFERENCES

- 1) Ali A. Textbook of pharmacognosy. New Delhi, India: publication and information directorate. 1993; 213-5.
- 2) Kokate C, Purohit AP, Gokhale SB. Pharmacognosy. Maharashtra, India: Nirali Prakashan. 2010; 355-7.
- 3) Sunarwidhi AL, Sudarsono S, Nugroho AE. Hypoglycemic effect of combination of *Azadirachta indica* A. Juss. and *Gynura procumbens* (Lour.) Merr. ethanolic extracts standardized by rutin and quercetin in alloxan-induced hyperglycemic rats. Adv Pharm Bull. 2014; 4: 613-8.
- 4) Kumar VS, Navaratnam V. Neem. *Neem (Azadirachta indica)*: prehistory to contemporary medicinal uses to humankind. Asian Pac J Trop Biomed. 2013; 3: 505-14.
- 5) Ogbuewu IP, Odoemenam VU, Obikaonu HO, Opara MN, Emenalom OO, Uchegbu MC, et al. The growing importance of neem (*Azadirachta indica* A. Juss) in agriculture, industry, medicine and environment: a review. Res J Med Plants. 2011; 5: 230-45.
- 6) Jarvis AP, Morgan EP, VanderEsch SA, Vitali F, Hey SU, Tape A. Identification of azadirachtin in tissue culture cells of neem. Nat Prod Lett. 1997; 10: 95-8.
- 7) Anonymous. Medicinal plants of India. New Delhi: Indian Council of Medical Research. 1: 112-7.
- 8) Chandra V. Botany of neem. Annals For. 1997; 5: 182-8.
- 9) Mitra R. Bibliography on pharmacognosy of medicinal plants. Lucknow: National Botanical Research Institute. 1985; 53.
- 10) Narayana AK. Nimba –Pharmacognosy of ayurvedic drugs of Kerala, Central Research Institute. Trivendrum. 1957; 1: 25-30.
- 11) Anonymous. The ayurvedic Pharmacopoeia of India. part 1, Nimba (leaf& Stem bark). New Delhi: Department of Ayush. 2: 124-34.
- 12) Ahmed E, Nidal J, Nagib E. A review of chemical constituents and traditional usage of Neem plant (*Azadirachta indica* A. Juss.). Palestinian Med Pharm J. 2017; 2: 75-81.

- 13) Srivasatava SK, Agarwal B, Akhilesh K, Archana P. Phytochemicals of *Azadirachta indica* source of active medicinal constituents used for cure of various diseases: a review. *J Sci Res.* 2020; 64: 385-90.
- 14) Saleem S, Muhammad G, Muhammed AH, Syed NAB. A comprehensive review of phytochemical profile, bioactive for pharmaceutical and pharmacological attributes of (*Azadirachta indica* A. Juss.). *Phytother Res.* 2018; 32: 1241-72.
- 15) Damtew M. A review on chemical composition, medicinal values and other application of *Azadirachta indica* A. Juss. *agri. Biol Res.* 2022; 38: 268-72.
- 16) Jytoi D, Sharma RB. Medicinal importance of *Azadirachta indica* A. Juss.: an overview. *J Drug Deliv Ther.* 2023; 13: 159-65.
- 17) Baby AR, Freire TB, Marques GA, Rijo P, Lima FV, Carvalho JCM, et al. Juss (Neem) as a potential natural active for dermocosmetic and topical products: a narrative review. *Cosmetics.* 2022; 9: 58-61.
- 18) Anonymous. Quality control of Indian medicinal Plants Vol 3. New Delhi: Department of Ayush. 1999; 201-7.
- 19) Kundu B, Subhasis K, Sarkar AB, Baral R, Laskar S. Chemical investigation of Neem leaf glycoproteins used as immunoprophylactic agent for tumor growth restriction. *Int J Pharm Pharm Sci.* 2018; 7: 195-9.
- 20) Gladkevich A, Bosker F, Korf J, Yenkovyan K, Vahradyan H, Aghajanov M. Proline-rich polypeptides in Alzheimer's disease and neurodegenerative disorders – therapeutic potential or a mirage? *Prog Neuropsychopharmacol Biol Psychiatry.* 2007; 31: 1347-55.
- 21) Yenkovyan K, Fereshetyan K, Matinyan S, Chavushyan V, Aghajanov M. The role of monoamines in the development of Alzheimer's disease and neuroprotective effect of a proline rich polypeptide. *Prog Neuropsychopharmacol Biol Psychiatry.* 2018; 86: 76-82.
- 22) Shilpa G, Renjitha J, Saranga R, Sajin FK, Nair MS, Joy B et al. Epoxyazadiradione purified from the *Azadirachta indica* seed induced mitochondrial apoptosis and inhibition of NFκB nuclear translocation in human cervical cancer cells. *Phytother Res.* 2017; 31: 1892-902.
- 23) Sithisarn P, Supabphol R, Gritsanapan W. Antioxidant activity of Siamese neem tree (VP1209). *J Ethnopharmacol.* 2005; 99: 109-12.
- 24) Nahak G, Sahu RK. Evaluation of antioxidant activity of flower and seed oil of *Azadirachta indica* A. Juss. *J Appl. Nat Sci.* 2011; 3: 78-81.
- 25) Ghimeray AK, Jin C, Ghimine BK, Cho DH. Antioxidant activity and quantitative estimation of azadirachtin and nimbin in *Azadirachta indica* A. Juss. grown in foothills of Nepal. *Afr J Biotechnol.* 2009; 8: 3084-91.
- 26) Sri U, Ibrahim M, Kumar M. Antioxidant activity and total flavonoids content of different parts of *Azadirachta indica* A. Juss. *J Med Plants Res.* 2012; 6: 5737-42.
- 27) Dhakal S, Aryal P, Aryal S, Bashyal D, Khadka D. Phytochemical and antioxidant studies of methanol and chloroform extract from leaves of *Azadirachta indica* A. Juss. in tropical region of Nepal. *J Pharmacogn Phytochem.* 2016; 8: 203-8.
- 28) Singh A, Singh AK, Narayan G, Singh TB, Shukla VK. Effect of neem oil and Haridra on non-healing wounds. *Ayu.* 2014; 35: 398-403.

- 29) Chundran NK, Husen IR, Rubianti I. Effect of neem leaves extract (*Azadirachta indica*) on Wound Healing. AMJ. 2015; 2: 199-207.
- 30) Osunwoke EA, Olotu EJ, Allison TA, Onyekwere JC. The wound healing effects of aqueous leave extracts of *Azadirachta indica* on Wistar rats. J Nat Sci Res. 2013; 3: 181-6.

