



INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

Maximal Mappings in Fuzzy Topological Spaces

Dr. Bhaskar A Munedewadi

Associate Professor

Head of the Department of Mathematics

Government First Grade College, Navanagar,

Bagalkot-587103, Karnataka, India

Abstract: In this paper, we introduce fuzzy semi regular weakly Maximal-open and fuzzy semi regular weakly Maximal-closed functions in fuzzy topological spaces and obtain certain characterization of the semi regular weakly Maximal closed and semi regular weakly Maximal open functions.

2010 Mathematics Subject Classification: 54A40

Key words and phrases: Fuzzy semi regular weakly Maximal-open functions, fuzzy semi regular weakly Maximal-closed functions, fuzzy semi regular weakly Maximal-closed sets, fuzzy semi regular weakly Maximal-open sets.

I. INTRODUCTION

The concept of a fuzzy subset was introduced and studied by L.A.Zadeh in the year 1965. The subsequent research activities in this area and related areas have found applications in many branches of science and engineering. In the year 1965, L.A.Zadeh [1] introduced the concept of fuzzy subset as a generalization of that of an ordinary subset. The introduction of fuzzy subsets paved the way for rapid research work in many areas of mathematical science. In the year 1968, C.L.Chang [2] introduced the concept of fuzzy topological spaces as an application of fuzzy sets to topological spaces. Subsequently several researchers contributed to the development of the theory and applications of fuzzy topology. The theory of fuzzy topological spaces can be regarded as a generalization theory of topological spaces. An ordinary subset A of a set X can be characterized by a function called characteristic function

$$\begin{aligned} \mu_A: X &\rightarrow [0,1] \text{ of } A, \text{ defined by} \\ \mu_A(x) &= 1, \text{ if } x \in A. \\ &= 0, \text{ if } x \notin A. \end{aligned}$$

Thus an element $x \in X$ is in A if $\mu_A(x) = 1$ and is not in A if $\mu_A(x) = 0$. In general if X is a set and A is a subset of X then A has the following representation. $A = \{ (x, \mu_A(x)): x \in X \}$, here $\mu_A(x)$ may be regarded as the degree of belongingness of x to A , which is either 0 or 1. Hence A is the class of objects with degree of belongingness either 0 or 1 only.

Prof. L.A.Zadeh [1] introduced a class of objects with continuous grades of belongingness ranging between 0 and 1 ; he called such a class as fuzzy subset. A fuzzy subset A in X is characterized as a membership function $\mu_A : X \rightarrow [0,1]$, which associates with each point in x a real number $\mu_A(x)$ between 0 and 1 which represents the degree or grade membership of belongingness of x to A .

The purpose of this paper is to introduce a new class of fuzzy sets called fuzzy semi regular weakly Maximal-closed sets in fuzzy topological spaces and investigate certain basic properties of these fuzzy sets. Among many other results it is observed that every fuzzy closed set is fuzzy semi regular weakly Maximal-closed but not conversely. Also we introduce fuzzy semi regular weakly Maximal-open sets in fuzzy topological spaces and study some of their properties.

1. PRELIMINARIES

1.1 Definition: [1] A fuzzy subset A in a set X is a function $A: X \rightarrow [0, 1]$. A fuzzy subset in X is empty iff its membership function is identically 0 on X and is denoted by 0 or μ_ϕ . The set X can be considered as a fuzzy subset of X whose membership function is identically 1 on X and is denoted by μ_X or I_X . In fact every subset of X is a fuzzy subset of X but not conversely. Hence the concept of a fuzzy subset is a generalization of the concept of a subset.

1.2 Definition : [1] If A and B are any two fuzzy subsets of a set X , then A is said to be included in B or A is contained in B iff $A(x) \leq B(x)$ for all x in X . Equivalently, $A \leq B$ iff $A(x) \leq B(x)$ for all x in X .

1.3 Definition:[1] Two fuzzy subsets A and B are said to be equal if $A(x) = B(x)$ for every x in X . Equivalently $A = B$ if $A(x) = B(x)$ for every x in X .

1.4 Definition:[1] The complement of a fuzzy subset A in a set X , denoted by A' or $1 - A$, is the fuzzy subset of X defined by $A'(x) = 1 - A(x)$ for all x in X . Note that $(A')' = A$.

1.5 Definition:[1] The union of two fuzzy subsets A and B in X , denoted by $A \vee B$, is a fuzzy subset in X defined by $(A \vee B)(x) = \text{Max}\{\mu_A(x), \mu_B(x)\}$ for all x in X .

1.6 Definition:[1] The intersection of two fuzzy subsets A and B in X , denoted by $A \wedge B$, is a fuzzy subset in X defined by $(A \wedge B)(x) = \text{Min}\{A(x), B(x)\}$ for all x in X .

1.7 Definition:[1] A fuzzy set on X is 'Crisp' if it takes only the values 0 and 1 on X .

1.8 Definition:[2] Let X be a set and τ be a family of fuzzy subsets of X . τ is called a fuzzy topology on X iff τ satisfies the following conditions.

(i) $\mu_\phi ; \mu_X \in \tau$: That is 0 and $1 \in \tau$

(ii) If $G_i \in \tau$ for $i \in I$ then $\bigvee_{i \in I} G_i \in \tau$

(iii) If $G, H \in \tau$ then $G \wedge H \in \tau$

The pair (X, τ) is called a fuzzy topological space (abbreviated as fts). The members of τ are called fuzzy open sets and a fuzzy set A in X is said to be closed iff $1 - A$ is a fuzzy open set in X .

1.9 Remark:[2] Every topological space is a fuzzy topological space but not conversely.

1.10 Definition:[2] Let X be a fts and A be a fuzzy subset in X . Then $\bigwedge \{B : B \text{ is a closed fuzzy set in } X \text{ and } B \geq A\}$ is called the closure of A and is denoted by A or $\text{cl}(A)$.

1.11 Definition:[2] Let A and B be two fuzzy sets in a fuzzy topological space (X, τ) and let $A \geq B$. Then B is called an interior fuzzy set of A if there exists $G \in \tau$ such that $A \geq G \geq B$, the least upper bound of all interior fuzzy sets of A is called the interior of A and is denoted by A^0 .

1.12 Definition[3] A fuzzy set A in a fts X is said to be fuzzy semi open if and only if there exists a fuzzy open set V in X such that $V \leq A \leq \text{cl}(V)$.

1.13 Definition [3] A fuzzy set A in a fts X is said to be fuzzy semi-closed if and only if there exists a fuzzy closed set V in X such that $\text{int}(V) \leq A \leq V$. It is seen that a fuzzy set A is fuzzy semi open if and only if $1 - A$ is a fuzzy semi-closed.

1.14 Theorem:[3] The following are equivalent:

- (a) μ is a fuzzy semiclosed set,
- (b) μ^c is a fuzzy semi open set,
- (c) $\text{int}(\text{cl}(\mu)) \leq \mu$.
- (b) $\text{int}(\text{cl}(\mu)) \geq \mu^c$

1.15 Theorem [3] Any union of fuzzy semi open sets is a fuzzy semi open set and (b) any intersection of fuzzy semi closed sets is a fuzzy semi closed.

1.16 Remark[3]

- (i) Every fuzzy open set is a fuzzy semi open but not conversely.
- (ii) Every fuzzy closed set is a fuzzy semi-closed set but not conversely.
- (iii) The closure of a fuzzy open set is fuzzy semi open set
- (iv) The interior of a fuzzy closed set is fuzzy semi-closed set

1.17 Definition:[3] A fuzzy set μ of a fts X is called a fuzzy regular open set of X if $\text{int}(\text{cl}(\mu)) = \mu$.

1.18 Definition:[3] A fuzzy set μ of fts X is called a fuzzy regular closed set of X if $\text{cl}(\text{int}(\mu)) = \mu$.

1.19 Theorem:[3] A fuzzy set μ of a fts X is a fuzzy regular open if and only if μ^c fuzzy regular closed set.

1.20 Remark:[3]

- (i) Every fuzzy regular open set is a fuzzy open set but not conversely.
- (ii) Every fuzzy regular closed set is a fuzzy closed set but not conversely.

1.21 Theorem:[3]

- (i) The closure of a fuzzy open set is a fuzzy regular closed.
- (ii) The interior of a fuzzy closed set is a fuzzy regular open set.

1.22 Definition:[4] A fuzzy set α in fts X is called fuzzy rw -closed if $\text{cl}(\alpha) \leq \mu$ whenever $\alpha \leq \mu$ and μ is regular semi-open in X .

1.23 Definition [5]:A fuzzy set α in fts X is called fuzzy semi regular weakly Maximal closed if $\text{p-cl}(\alpha) \leq \mu$ whenever $\alpha \leq \mu$ and μ is $\text{rg}\alpha$ -open set in X .

1.24 Definition [5]:A fuzzy set α of a fts X is fuzzy semi regular weakly Maximal-open set, if it's complement α^c is a fuzzy semi regular weakly Maximal-closed in fts X .

1.25 definition [2]: Let X and Y be fts. A function $f: X \rightarrow Y$ is said to be a fuzzy continuous function if $f^{-1}(\mu)$ is fuzzy open in X for each fuzzy open set μ in Y .

1.26 definition[6]:A function $f: X \rightarrow Y$ is said to be fuzzy irresolute iff $f^{-1}(B)$ is fuzzy semi-open in X is fuzzy open in Y .

1.27 Definition[7]:A function $f: X \rightarrow Y$ is said to be fuzzy semi regular weakly Maximal irresolute if the inverse image of every fuzzy semi regular weakly Maximal-open in Y is a fuzzy semi regular weakly Maximal-open set in X .

2. FUZZY SEMI REGULAR WEAKLY MAXIMAL-OPEN FUNCTIONS AND FUZZY SEMI REGULAR WEAKLY MAXIMAL CLOSED FUNCTIONS IN FUZZY TOPOLOGICAL SPACES

Definition 2.1: Let X and Y be two fts. A function $f: (X, T_1) \rightarrow (Y, T_2)$ is called fuzzy semi regular weakly Maximal-open function if the image of every fuzzy open set in X is fuzzy semi regular weakly Maximal-open in Y .

Theorem 2.2: Every fuzzy open function is a fuzzy semi regular weakly Maximal-open function.

Proof: Let $f: (X, T_1) \rightarrow (Y, T_2)$ be a fuzzy open function and μ be a fuzzy open set in fts X . Then $f(\mu)$ is a fuzzy open set in fts Y . Since every fuzzy open set is fuzzy semi regular weakly Maximal-open, $f(\mu)$ is a fuzzy semi regular weakly Maximal-open set in fts Y . Hence f is a fuzzy semi regular weakly Maximal-open function.

The converse of the above theorem need not be true in general as seen from the following example.

Example 2.3: Let $X = Y = \{a, b, c\}$ and the functions $\alpha, \beta: X \rightarrow [0, 1]$ be defined as

$$\alpha(x) = \begin{cases} 1 & \text{if } x = a, c \\ 0 & \text{otherwise} \end{cases}$$

$$\beta(x) = \begin{cases} 1 & \text{if } x = a \\ 0 & \text{otherwise} \end{cases}$$

Consider $T_1 = \{1, 0, \alpha\}$, $T_2 = \{1, 0, \beta\}$ then (X, T_1) and (Y, T_2) are fts. Let $f: X \rightarrow Y$ be the identity function. Then this function is semi regular weakly Maximal-open function but it is not fuzzy open. Since the image of the fuzzy open set $\alpha(x)$ in X is fuzzy set β in Y which is not fuzzy open.

Remark 2.4 : If $f: (X, T_1) \rightarrow (Y, T_2)$ and $g: (Y, T_2) \rightarrow (Z, T_3)$ be two fuzzy semi regular weakly Maximal open function then their composition $g \circ f: (X, T_1) \rightarrow (Z, T_3)$ need not be fuzzy semi regular weakly Maximal-open as seen from the following example

Example 2.5: Let $X = Y = Z = \{a, b, c\}$ and the function $\alpha, \beta, \gamma, \delta: X \rightarrow [0, 1]$ be defined as

$$\alpha(x) = \begin{cases} 1 & \text{if } x = a \\ 0 & \text{otherwise} \end{cases}$$

$$\beta(x) = \begin{cases} 1 & \text{if } x = c \\ 0 & \text{otherwise} \end{cases}$$

$$\gamma(x) = \begin{cases} 1 & \text{if } x = a, c \\ 0 & \text{otherwise} \end{cases}$$

$$\delta(x) = \begin{cases} 1 & \text{if } x = a, b \\ 0 & \text{otherwise} \end{cases}$$

Consider $T_1 = \{1, 0, \alpha, \delta\}$, $T_2 = \{1, 0, \alpha\}$ and $T_3 = \{0, 1, \alpha, \beta, \gamma\}$. Let function $f: (X, T_1) \rightarrow (Y, T_2)$ and $g: (Y, T_2) \rightarrow (Z, T_3)$ be the identity functions then f and g are fuzzy semi regular weakly Maximal-open function but their composition $g \circ f: (X, T_1) \rightarrow (Z, T_3)$ is not fuzzy semi regular weakly Maximal-open as $\beta: X \rightarrow [0, 1]$ is fuzzy open in X . But $(g \circ f)(\beta) = \beta$ is not fuzzy semi regular weakly Maximal-open in Z .

Theorem 2.6 : If $f: (X, T_1) \rightarrow (Y, T_2)$ is fuzzy open function and $g: (Y, T_2) \rightarrow (Z, T_3)$ is fuzzy semi regular weakly Maximal open function then their composition $g \circ f: (X, T_1) \rightarrow (Z, T_3)$ is fuzzy semi regular weakly Maximal-open function.

Proof: Let α be fuzzy open set in (X, T_1) . since f is fuzzy open function $f(\alpha)$ is a fuzzy open set in (Y, T_2) . Since g is a fuzzy semi regular weakly Maximal-open function $g(f(\alpha)) = (g \circ f)(\alpha)$. Thus $g \circ f$ is a fuzzy semi regular weakly Maximal-open function.

Definition 2.7: Let X and Y be two fuzzy topological spaces. A function $f: (X, T_1) \rightarrow (Y, T_2)$ is called fuzzy semi regular weakly Maximal-closed function if the image of every fuzzy closed set in X is a fuzzy closed set in Y .

Theorem 2.8: Let $f: (X, T_1) \rightarrow (Y, T_2)$ be fuzzy closed function then f is a fuzzy semi regular weakly Maximal closed function.

Proof: Let α be a fuzzy closed set in (X, T_1) . since f is fuzzy closed function, $f(\alpha)$ is a fuzzy closed set in (Y, T_2) . since every fuzzy closed set is fuzzy semi regular weakly Maximal-closed, $f(\alpha)$ is a fuzzy semi regular weakly Maximal-closed set in (Y, T_2) . Hence f is fuzzy semi regular weakly Maximal-closed function.

The converse of the theorem need not be true in general as seen from the following example

Example 2.9: Let $X = Y = \{a, b, c\}$ and the functions $\alpha, \beta: X \rightarrow [0, 1]$ be defined as

$$\alpha(x) = \begin{cases} 1 & \text{if } x = a, c \\ 0 & \text{otherwise} \end{cases}$$

$$\beta(x) = \begin{cases} 1 & \text{if } x = a \\ 0 & \text{otherwise} \end{cases}$$

Consider $T_1 = \{1, 0, \alpha\}$ and $T_2 = \{1, 0, \beta\}$; then (X, T_1) and (Y, T_2) are fts. Let $f: (X, T_1) \rightarrow (Y, T_2)$ be the identity function. Then f is a fuzzy semi regular weakly Maximal-closed function but it is not a fuzzy closed function, since the image of the closed set α in X is not a fuzzy closed set in Y .

Remark 2.10: The composition of two fuzzy semi regular weakly Maximal closed functions need not be a fuzzy semi regular weakly Maximal-closed function

Example 2.11: Consider the fts $(X, T_1), (Y, T_2), (Z, T_3)$ and functions f, g defined in 2.5. the functions f and g are fuzzy semi regular weakly Maximal-closed but their composition is not fuzzy semi regular weakly Maximal-closed as $\beta: X \rightarrow [0, 1]$ is a fuzzy closed set in X but $(g \circ f)(\beta) = \beta$ is not fuzzy semi regular weakly Maximal-closed in Z .

Theorem 2.12: If a function $f: (X, T_1) \rightarrow (Y, T_2)$ and $g: (Y, T_2) \rightarrow (Z, T_3)$ be two functions then composition of two semi regular weakly Maximal-closed functions is fuzzy semi regular weakly Maximal-closed function.

Proof: Let α be a fuzzy closed set in (X, T_1) . Since f is a fuzzy closed function, $f(\alpha)$ is a fuzzy closed set in (Y, T_2) . Since g is a fuzzy semi regular weakly Maximal-closed function but $g(f(\alpha))$ is a fuzzy semi regular weakly Maximal-closed set in (Z, T_3) but $g(f(\alpha)) = (g \circ f)(\alpha)$. Thus $g \circ f$ is fuzzy semi regular weakly Maximal-closed function.

Theorem 2.13: A function $f: X \rightarrow Y$ is fuzzy semi regular weakly Maximal-closed function if for each fuzzy set δ of Y and for each fuzzy open set μ of X s.t $\mu \geq f^{-1}(\delta)$, there is a fuzzy semi regular weakly Maximal-open function set α of Y s.t $\delta \leq \alpha$ and $f^{-1}(\alpha) \leq \mu$.

Proof: Suppose that f is fuzzy semi regular weakly Maximal-closed function. Let δ be a fuzzy subset of Y and μ be a fuzzy open set of X s.t $f^{-1}(\delta) \leq \mu$. Let $\alpha = 1 - f(1 - \mu)$ is fuzzy semi regular weakly Maximal-open set in fts Y . note that $f^{-1}(\delta) \leq \mu$. Which implies $\delta \leq \alpha$ and $f^{-1}(\alpha) \leq \mu$. For the converse, Suppose that μ is a fuzzy closed set in X , Then $f^{-1}(1 - f(\mu)) \leq 1 - \mu$ And $1 - \mu$ is fuzzy open by hypothesis, there is a fuzzy semi regular weakly Maximal-open set α of Y s.t $(1 - f(\mu)) \leq \alpha$. And $f^{-1}(\alpha) \leq 1 - \mu$. Therefore $\mu \leq 1 - f^{-1}(\alpha)$. Hence $1 - \alpha \leq \mu$, $f(1 - f^{-1}(\alpha)) \leq 1 - \alpha$. Which implies $f(\mu) = 1 - \alpha$. Since $1 - \alpha$ is fuzzy semi regular weakly Maximal-closed $f(\mu)$ is fuzzy semi regular weakly Maximal-closed and thus f is fuzzy semi regular weakly Maximal-closed.

Lemma 2.14: Let $f: (X, T_1) \rightarrow (Y, T_2)$ be fuzzy irresolute function and α be fuzzy rg α open in Y then $f^{-1}(\alpha)$ is fuzzy rg α open in X .

Proof: Let α be fuzzy $rg\alpha$ -open in Y . To prove $f^{-1}(\alpha)$ is fuzzy $rg\alpha$ open in X that is to prove $f^{-1}(\alpha)$ is both fuzzy semi-open and fuzzy semi-closed in X . Now α is fuzzy semi-open in Y . Since f is fuzzy irresolute function, $f^{-1}(\alpha)$ is fuzzy semi-open in X . Now α is fuzzy semi-closed in Y as fuzzy $rg\alpha$ open set is fuzzy semi-closed then $1 - \alpha$ is fuzzy semi-open in Y . Since f is fuzzy irresolute function, $f^{-1}(1 - \alpha)$ is a fuzzy semi open in X . But $f^{-1}(1 - \alpha) = 1 - f^{-1}(\alpha)$ is fuzzy semi-open in X and so $f^{-1}(\alpha)$ is semi closed in X . Thus $f^{-1}(\alpha)$ is both fuzzy semi-open and fuzzy semi-closed in X and hence $f^{-1}(\alpha)$ is fuzzy $rg\alpha$ open in X .

Theorem 2.15: If a function $f: (X, \mathcal{T}_1) \rightarrow (Y, \mathcal{T}_2)$ is fuzzy irresolute function and fuzzy semi regular weakly Maximal-closed and α is fuzzy semi regular weakly Maximal-closed set of X , then $f(\alpha)$ is a fuzzy semi regular weakly Maximal-closed set in Y .

Proof: Let α be a fuzzy closed set of X . Let $f(\alpha) \leq \mu$. Where μ is fuzzy $rg\alpha$ open in Y . Since f is fuzzy irresolute function, $f^{-1}(\mu)$ is a fuzzy $rg\alpha$ open in X , by lemma 2.14 and $\alpha \leq f^{-1}(\mu)$. Since α is a fuzzy semi regular weakly Maximal-closed set in X , $P\text{-cl}(\alpha) \leq f^{-1}(\mu)$. Since f is a fuzzy semi regular weakly Maximal-closed set contained in the fuzzy $rg\alpha$ open set μ , which implies $\text{cl}(f(P\text{-cl}(\alpha))) \leq \mu$ and hence $\text{cl}(f(\alpha)) \leq \mu$. Therefore $f(\alpha)$ is a fuzzy semi regular weakly Maximal closed set in Y .

Corollary 2.16: If a function $f: (X, \mathcal{T}_1) \rightarrow (Y, \mathcal{T}_2)$ is fuzzy irresolute function and fuzzy closed and α is a fuzzy semi regular weakly Maximal-closed set in X , then $f(\alpha)$ is a fuzzy semi regular weakly Maximal-closed set in Y .

Proof : The proof follows from the theorem 2.15 and the fact that every fuzzy closed function is a fuzzy semi regular weakly Maximal-closed function.

Theorem 2.17: Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two functions s.t

- (i) if f is fuzzy continuous function and surjective, then g is fuzzy semi regular weakly Maximal closed function.
- (ii) if g is fuzzy semi regular weakly Maximal-irresolute function and injective then f is fuzzy semi regular weakly Maximal-closed function.

Proof :

(i) Let μ be a fuzzy closed set in Y , since f is fuzzy continuous function, $f^{-1}(\mu)$ is a fuzzy closed set in X . Since $g \circ f$ is a fuzzy semi regular weakly Maximal closed function, $(g \circ f)(f^{-1}(\mu))$ is a fuzzy semi regular weakly Maximal-closed set in Z . but $(g \circ f)(f^{-1}(\mu)) = g(\mu)$ as f is surjective thus g is fuzzy semi regular weakly Maximal closed function.

(ii) Let β be a fuzzy closed set of X then $(g \circ f)(\beta)$ is a fuzzy semi regular weakly Maximal closed set in Z . Since $(g \circ f)$ is a fuzzy semi regular weakly Maximal-closed function. Since g is fuzzy semi regular weakly Maximal-irresolute function $g^{-1}((g \circ f)(\beta))$ is fuzzy semi regular weakly Maximal-closed in Y . But $g^{-1}((g \circ f)(\beta)) = f(\beta)$ as g is injective. Thus f is fuzzy semi regular weakly Maximal-closed function.

References:

- [1] L.A.Zadeh, Fuzzy sets, Information and control, 8 (1965) 338-353.
- [2] C.L.Chang, Fuzzy topological spaces, JI. Math. Anal. Appl., 24(1968), 182-190.
- [3] K.K.Azad, On fuzzy semi continuity, fuzzy almost continuity and fuzzy weakly continuity. JI. Math. Anal. Appl. 82 No. 1 (1981), 14-32.
- [4] S. S. Benchalli, R. S. Wali and Basavaraj M. Ittanagi on fuzzy rw-closed sets and fuzzy rw-open sets in fuzzy topological spaces Int. J. of Mathematical Sciences and Applications, Vol. 1, No. 2, May 2011.
- [5] R.S.Wali and Vivekananda Dembre; Fuzzy semi regular weakly Maximal Maximal-closed sets and Fuzzy semi regular weakly Maximal Maximal-open sets in Topological Spaces. Volume 3, No. 3, March 2016 Journal of Global Research in Mathematical Archives.
- [6] Munkerjee, M.N.Sinha,, S.P.Irresolute and almost open functions between fts, fuzzy sets and systems ,29(1989)381-388.
- [7] R. S. Wali and Vivekananda Dembre; fuzzy semi regular weakly Maximal-continuous functions and fuzzy semi regular weakly Maximal Maximal-irresolute in fts. International Journal of Statistics and Applied Mathematics 2016; 1(1): 01-07.
- [8] A.Vadivel & K.vairamamanickam, $rg\alpha$ -Closed sets & $rg\alpha$ -open sets in Topological Spaces, Int J of math ,Analysis Vol 3, (2009)37,1803-1819.
- [9] Thakur S.S. and Bajpai Pandey Jyoti "Intuitionistic Fuzzy $rg\alpha$ -closed sets", International Journal of Fuzzy system and Rough System 4(1), 67-73.
- [10] R.S.Wali and Vivekananda Dembre; On Pre Generalized Pre Regular Weakly Closed Sets in Topological Spaces ;Journal of Computer and Mathematical Sciences, Vol.6(2),113-125, February 2015.